

Document made available under the Patent Cooperation Treaty (PCT)

International application number: PCT/US05/006588

International filing date: 02 March 2005 (02.03.2005)

Document type: Certified copy of priority document

Document details: Country/Office: US
Number: 60/549,832
Filing date: 02 March 2004 (02.03.2004)

Date of receipt at the International Bureau: 16 August 2005 (16.08.2005)

Remark: Priority document submitted or transmitted to the International Bureau in compliance with Rule 17.1(a) or (b)



World Intellectual Property Organization (WIPO) - Geneva, Switzerland
Organisation Mondiale de la Propriété Intellectuelle (OMPI) - Genève, Suisse



BANNER & WITCOFF, LTD.
INTELLECTUAL PROPERTY LAW

1001 G STREET, N.W.
WASHINGTON, DC 20001-4597

TEL: 202.824.3000
FAX: 202.824.3001
www.bannerwitcoff.com

August 11, 2005

BY FEDERAL EXPRESS

The International Bureau of WIPO
34, chemin des Colombettes
1211 Geneva 20, Switzerland

Attn: Dominique Delmas

Re: CHIRON CORPORATION
PCT Application No. PCT/US2005/006588
Our Ref.: 002441.00124:PP21431.004

Dear Sir:

Further to our letter dated August 1, 2005, we now enclose a certified copy of priority document for U.S. Serial No. 60/549,832 for the above-identified application.

Respectfully submitted,

Rebecca M. Hale

SCS/mh

CHICAGO, IL
WASHINGTON, DC
BOSTON, MA
PORTLAND, OR

FedEx International Air Waybill

Express

From **08/11/05** Sender's FedEx Account Number **0200 0635 2**

Date **08/11/05** Sender's Name **Margaret Hoffman** (202) 224-3438

Company **Banner Wireless Ltd**

Address **1001 G Street NW**

City **Washington** State **DC**

Country **USA** ZIP **20001-4517**

2 Your Internal Billing Reference **002441-00124/0750**

3 To Recipient's Name **Dominique Delphine**

Company **Internationale Bureau d'IMP**

Address **34 chemin des Colonnelles**

City **Geneve** State **20**

Country **Suisse** ZIP **1211**

Recipient's Tax ID number for Customs purposes **ALL shipments can be subject to Customs charges.**

4 Shipment Information **Total Packages 1** **Total Weight 1.0** **kg** **1.0** **lb** **1.0** **oz**

Commodity Description **Test Documents** **Commodity Code** **Country of Manufacture** **Value for Customs**

For U.S. Export Only: Check One ☐ No SBT required per Description ☐ SBT required per Description

For U.S. Export Only: Check One ☐ No SBT required per Description ☐ SBT required per Description

For U.S. Export Only: Check One ☐ No SBT required per Description ☐ SBT required per Description

For U.S. Export Only: Check One ☐ No SBT required per Description ☐ SBT required per Description

FedEx Tracking Number **8277 9494 7450 0402**

5 Express Package Service ☐ FedEx Intl. Priority ☐ FedEx Intl. Economy

6 Packaging ☒ FedEx Envelope ☐ Other Pkg. ☐ FedEx Pkg. Box

7 Special Handling ☐ HOLD at FedEx Location ☐ SATURDAY Delivery

8a Payment Bill transportation charges to: ☐ Sender Acct. No. ☐ Recipient ☐ Third Party

8b Payment Bill duties and taxes to: ☐ Sender Acct. No. ☐ Recipient ☐ Third Party

9 Required Signature **Signature** **Date**

10 FedEx Tracking Number **8277 9494 7450 0402**

11 FedEx Tracking Number **8277 9494 7450 0402**

12 FedEx Tracking Number **8277 9494 7450 0402**

13 FedEx Tracking Number **8277 9494 7450 0402**

14 FedEx Tracking Number **8277 9494 7450 0402**

15 FedEx Tracking Number **8277 9494 7450 0402**

16 FedEx Tracking Number **8277 9494 7450 0402**

17 FedEx Tracking Number **8277 9494 7450 0402**

18 FedEx Tracking Number **8277 9494 7450 0402**

19 FedEx Tracking Number **8277 9494 7450 0402**

20 FedEx Tracking Number **8277 9494 7450 0402**

Not all services and options are available to all destinations.

Package up to 150 lbs. / 68 kg for packages over 150 lbs. / 68 kg, use the FedEx Express Service Int. Air Waybill.

Package up to 150 lbs. / 68 kg for packages over 150 lbs. / 68 kg, use the FedEx Express Service Int. Air Waybill.

Package up to 150 lbs. / 68 kg for packages over 150 lbs. / 68 kg, use the FedEx Express Service Int. Air Waybill.

Package up to 150 lbs. / 68 kg for packages over 150 lbs. / 68 kg, use the FedEx Express Service Int. Air Waybill.

Package up to 150 lbs. / 68 kg for packages over 150 lbs. / 68 kg, use the FedEx Express Service Int. Air Waybill.

Package up to 150 lbs. / 68 kg for packages over 150 lbs. / 68 kg, use the FedEx Express Service Int. Air Waybill.

Package up to 150 lbs. / 68 kg for packages over 150 lbs. / 68 kg, use the FedEx Express Service Int. Air Waybill.

Package up to 150 lbs. / 68 kg for packages over 150 lbs. / 68 kg, use the FedEx Express Service Int. Air Waybill.

Package up to 150 lbs. / 68 kg for packages over 150 lbs. / 68 kg, use the FedEx Express Service Int. Air Waybill.

Package up to 150 lbs. / 68 kg for packages over 150 lbs. / 68 kg, use the FedEx Express Service Int. Air Waybill.

Package up to 150 lbs. / 68 kg for packages over 150 lbs. / 68 kg, use the FedEx Express Service Int. Air Waybill.

Package up to 150 lbs. / 68 kg for packages over 150 lbs. / 68 kg, use the FedEx Express Service Int. Air Waybill.

Package up to 150 lbs. / 68 kg for packages over 150 lbs. / 68 kg, use the FedEx Express Service Int. Air Waybill.

Package up to 150 lbs. / 68 kg for packages over 150 lbs. / 68 kg, use the FedEx Express Service Int. Air Waybill.

Package up to 150 lbs. / 68 kg for packages over 150 lbs. / 68 kg, use the FedEx Express Service Int. Air Waybill.

Package up to 150 lbs. / 68 kg for packages over 150 lbs. / 68 kg, use the FedEx Express Service Int. Air Waybill.

Package up to 150 lbs. / 68 kg for packages over 150 lbs. / 68 kg, use the FedEx Express Service Int. Air Waybill.

FedEx International Air Waybill

Express

From **Sender's FedEx Account Number** 0200 0655 2

Date **08/11/85**

Sender's Name **Margaret Platt** Phone **(202) 284 310**

Company **Painter & Witall Ltd**

Address **1001 G Street NW**

City **Washington** State **DC**

Country **USA** ZIP **20001-4517**

Postal Code **20001-4517**

2 Your Internal Billing Reference **002441-0124/0750**

3 To Recipient's Name **Prinimique, Belin**

Company **Int'l. Airline Bureau (IABP)**

Address **34 Chemin des Colonnettes**

City **Geneve** State **20**

Country **Switzerland** ZIP **1211**

Postal Code **1211**

Recipient's Tax ID number for Customs purposes

4 Shipment Information

Total Packages **11** Total Weight **44** lbs. **kg** DIM **1** / **1** / **1** in. **cm**

Commodity Description **REQUIRED**

Country of Manufacture **REQUIRED**

Value for Customs **REQUIRED**

FedEx Tracking Number **8277 9494 7450** Form ID No. **0402**

5 Express Package Service

1 ☐ FedEx Intl. Priority 6 ☐ FedEx Intl. First

3 ☒ FedEx Intl. Economy 7 ☐ FedEx Intl. Extra Hours

6 Packaging 1 ☐ FedEx Envelope 2 ☐ FedEx Box

7 Special Handling 3 ☐ SATURDAY Delivery

8a Payment Bill transportation charges to:

8b Payment Bill duties and taxes to:

9 Required Signature

Signature

Recipient's Signature

Origin Station ID

Destination Station ID

UNSA Routing

Handling Units

Total Volume (cm)

Forms Attached

CI ☐ SED ☐ CO

Time

Date

Time

Date

Time

Not all services and options are available to all destinations.

Packages up to 150 lbs./68 kg

For packages over 150 lbs./68 kg use the FedEx Express Service Int'l. Air Waybill.

These unique line numbers with special printing provided by FedEx are for tracking and delivery confirmation only.

FedEx Tracking Number

FedEx Tracking Number

FedEx Tracking Number

FedEx Tracking Number

FedEx Tracking Number

FedEx Tracking Number

FedEx Tracking Number

FedEx Tracking Number

FedEx Tracking Number

FedEx Tracking Number

FedEx Tracking Number

FedEx Tracking Number

FedEx Tracking Number

FedEx Tracking Number

FedEx Tracking Number

FedEx Tracking Number

FedEx Tracking Number

FedEx Tracking Number

FedEx Tracking Number

01 SRC

Express

[illegible]

Not all services and options are available to all destinations.

Not all services and options are available to all destinations.

Express

Origin Station ID.		Destination Station ID.		USPS Routing		Handling Units	
						Total Volume (cm)	
Received At: 1	Drop Stop 2	On-Cut Stop 3	Drop Box 4	World Service Center 5	Station	Forms Attached:	<input type="checkbox"/> CI <input type="checkbox"/> SED <input type="checkbox"/> CO
Base Charges:	Declared Value Chrg.	Other	DDM	DDI	Card Auth.		
Exps. #	Exps. #	Date	Time	Exps. #	Date	Time	

Non-Negotiable International Air Waybill • ©194-2000 Federal Express Corporation

FedEx
Tracking
Number

8277 9494 7450

I.D. No.

0402

PART 155925
Rev. Data 10/00
©1994-2000 Fede
PRINTED IN U.S.A.

47201 SRC

9 Required Signature

8b Payment Bill dates and taxes to:

Enter FedEx Acct. No. below. Cash ☐

☐ Sender Acct. No. in Section 1 will be billed. Check ☐

☐ Recipient ☐ Third ☐ Credit ☐

☐ Party ☐ Unique ☐

8b	Payment Bill duties and taxes to:	FedEx cannot estimate customs charges
	Enter FedEx Acct No below: <input type="text"/>	Cash

☐ **Section Act No. in Section with be listed.**

☐ **Enter full Act No. or Credit Card No. below.**

☐ **Receipt**
☐ **Third**
☐ **Credit**

☐ **Fairly**
☐ **Card**

☐ **Cash**

☐ **Cheq/**

☐ **Cheque**

☐ **Total Transportation**

Specify Currency

8a Payment Bill transportation charges to

☐ **HOLD at FedEx Location** ☐ **SATURDAY Delivery**
Available to select locations
Available for FedEx Int'l. Priority only

7 Special Handling

☐ **FedEx Envelope** 
☐ **FedEx Pak** 
☐ **Other Pkg.** 

☐ **10kg Box** 
☐ **25kg Box** 

☐ **Includes FedEx Box**
☐ **FedEx 10kg and 25kg boxes and customer pkg.**

5 Packaging

NEW **FedEx Int'l. Extra Hours** Available for select locations

FedEx Int'l. Economy **FedEx Envelope and FedEx Pak** **not available**

These unique brown boxes with special yellow and green labels are available for FedEx Int'l. Priority and FedEx Int'l. Economy.

☐ **FedEx Intl. Priority**


☐ **FedEx Intl. First**
Available to select locations.
Higher rates apply.

Express Package Service

Ex tracking number

8277 9494 7450

01107



THE UNITED STATES OF AMERICA

TO ALL TO WHOM THESE PRESENTS SHALL COME:

UNITED STATES DEPARTMENT OF COMMERCE

United States Patent and Trademark Office


August 09, 2005

THIS IS TO CERTIFY THAT ANNEXED HERETO IS A TRUE COPY FROM THE RECORDS OF THE UNITED STATES PATENT AND TRADEMARK OFFICE OF THOSE PAPERS OF THE BELOW IDENTIFIED PATENT APPLICATION THAT MET THE REQUIREMENTS TO BE GRANTED A FILING DATE UNDER 35 USC 111.

APPLICATION NUMBER: 60/549,832

FILING DATE: March 02, 2004

**By Authority of the
Under Secretary of Commerce for Intellectual Property
and Director of the United States Patent and Trademark Office**



N. Woodson
N. WOODSON
Certifying Officer

PROVISIONAL APPLICATION FOR PATENT COVER SHEET

This is a request for filing a PROVISIONAL APPLICATION FOR PATENT under 37 CFR 1.53(c).

Express Mail Label No. EL 987 060 923 US

INVENTOR(S)

Given Name (first and middle (if any))	Family Name or Surname	Residence (City and either State or Foreign Country)
Guido	Grandi	Milano, Italy

Additional inventors are being named on the second separately numbered sheets attached hereto

TITLE OF THE INVENTION (500 characters max)

Immunogenic Compositions For Chlamydia Pneumoniae

Direct all correspondence to:

CORRESPONDENCE ADDRESS

☒ Customer Number:

27476

OR

☐ Firm or
Individual Name

Address

Address

City

State

Zip

Country

Telephone

Fax

ENCLOSED APPLICATION PARTS (check all that apply)

☒ Specification Number of Pages 113



CD(s), Number

☒ Drawing(s) Number of Sheets 6



Other (specify) Tables (8 sheets)

☐ Application Date Sheet. See 37 CFR 1.76

METHOD OF PAYMENT OF FILING FEES FOR THIS PROVISIONAL APPLICATION FOR PATENT

☐ Applicant claims small entity status. See 37 CFR 1.27.

☒ A check or money order is enclosed to cover the filing fees.

☒ The Director is hereby authorized to charge filing fees or credit any overpayment to Deposit Account Number: 03-1664

☐ Payment by credit card. Form PTO-2038 is attached.

FILING FEE
Amount (\$)

\$160

The invention was made by an agency of the United States Government or under a contract with an agency of the United States Government.

☐ No.

☐ Yes, the name of the U.S. Government agency and the Government contract number are:

[Page 1 of 2]

Respectfully submitted,

SIGNATURE

Rebecca M. Hale

TYPED or PRINTED NAME Rebecca M. Hale

TELEPHONE (510) 923-3179

Date

March 2, 2004

REGISTRATION NO. 45,680

(if appropriate)

Docket Number: 21431.001

USE ONLY FOR FILING A PROVISIONAL APPLICATION FOR PATENT

This collection of information is required by 37 CFR 1.51. The information is required to obtain or retain a benefit by the public which is to file (and by the USPTO to process) an application. Confidentiality is governed by 35 U.S.C. 122 and 37 CFR 1.14. This collection is estimated to take 8 hours to complete, including gathering, preparing, and submitting the completed application form to the USPTO. Time will vary depending upon the individual case. Any comments on the amount of time you require to complete this form and/or suggestions for reducing this burden, should be sent to the Chief Information Officer, U.S. Patent and Trademark Office, U.S. Department of Commerce, P.O. Box 1450, Alexandria, VA 22313-1450. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO: Mail Stop Provisional Application, Commissioner for Patents, P.O. Box 1450, Alexandria, VA 22313-1450.

If you need assistance in completing the form, call 1-800-PTO-9199 and select option 2.

PROVISIONAL APPLICATION COVER SHEET
Additional Page

PTO/SB/16 (08-03)

Approved for use through 07/31/2006. OMB 0651-0032

U.S. Patent and Trademark Office; U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

Docket Number 21431.001

INVENTOR(S)/APPLICANT(S)		
Given Name (first and middle [if any])	Family or Surname	Residence (City and either State or Foreign Country)
Giulio	Ratti	

[Page 2 of 2]

Number 2 of 2

WARNING: Information on this form may become public. Credit card information should not be included on this form. Provide credit card information and authorization on PTO-2038.

IMMUNOGENIC COMPOSITIONS FOR *CHLAMYDIA PNEUMONIAE*

All documents cited herein are incorporated by reference in their entirety.

Field

- 5 The invention is in the field of immunology and vaccinology. In particular, it relates to immunogenic compositions comprising combinations of immunogenic molecules from *Chlamydia pneumoniae*.

Background Art

- 10 The bacteria of the genus *Chlamydia* (and *Chlamydophila*, according to the recently proposed but still controversial re-classification of *Chlamydiaceae* (Bush *et al* (2001) Int J Syst Evol Microbiol 51: 203-20; Everett *et al* (1999) Int J Syst Bacteriol 49: Pt2 415-40; Schachter *et al* (2001) Int J Syst Evol Microbiol 51: 249, 251-3) are obligate
15 intracellular parasites of eukaryotic cells, which have a unique biphasic life cycle involving two pleiomorphic developmental forms: an extracellular, metabolically inert, spore-like, infectious form (the elementary bodies, EBs) and an intracellular, non-infectious, replicative form (the reticulate bodies, RBs) which remains contained in a specialized cytoplasmic compartment (the *Chlamydial* inclusion). The EBs are
20 responsible for the initial attachment to host cell surface and the establishment of the cytoplasmic inclusion where EBs can differentiate to RBs and thus initiate the replicative stage. Eventually RBs revert to infectious EB forms able to start new replicative cycles in neighbouring host cells.

- 25 As *Chlamydia* infection is an intracellular infection, the currently accepted paradigm is that effective anti-*Chlamydial* immunisation would require both an adequate T-cell response and high serum levels of neutralising antibodies and that "an ideal vaccine should induce long lasting (neutralising) antibodies and a cell mediated immunity that can quickly respond upon exposure to *Chlamydia*". Several sometimes contradictory studies have indicated that both CD4+ and CD 8 positive T cells have a role in
30 *Chlamydial* clearance (Loomis and Starnback (2002) Curr Opin Microbiol 5: 87-91). Indeed, there now appears to be a prevailing consensus that specific CD4+ T cells and B cells are critical to the complete clearance of intracellular *Chlamydia* and for mediating recall immunity to *Chlamydia* infection (see Igiertseme, Black and Caldwell (2002) Biodrugs 16: 19-35 and Igiertseme *et al* (1999) Immunology 98: 510-519).

- 35 Whilst it is now possible to carry out searches of the whole *Chlamydia pneumoniae* genome, there is still insufficient information available on parallel proteome characterisation. By way of example, while sequence data is available for many of the *Chlamydia pneumoniae* antigens, there is insufficient characterisation of the
40 *Chlamydia* antigens in terms of their immunological and/or biological function. By way of example, whilst applications such as WO 99/28475 and WO 99/27105 disclose sequence information, there is no characterisation of these sequences in terms of their immunological and/or biological function. In contrast, WO 02/02404 provides information on the immunogenicity and immunoaccessibility of certain *Chlamydia*
45 proteins and highlights that (i) current genomic annotations and/or (ii) predictions based on cellular location and/or cellular function based on *in-silico* analyses may not always be accurate.

- Applicants have recently engaged in a whole-genome search (Montigiani *et al* (2002) Infection and Immunity 70:368-379) for possible vaccine candidates among proteins
50 potentially associated with the outer membrane of *C.pneumoniae*. For this study,

5 mouse antisera was prepared against over 100 recombinant His-tagged or Glutathione-S-transferase (GST) fusion proteins encoded by genes predicted by in silico analyses to be peripherally located in the *Chlamydial* cell. From this screening study, 53 recombinant proteins derived from the genome of *Chlamydia* (*Chlamydophila pneumoniae* (CPn) were described which induced mouse antibodies, capable of binding, in a FACS assay, to the surface of purified CPn cells.

10 The scope of the Montigiani study (*ibid*) was restricted to checking if polyclonal antisera produced in mice against the recombinantly expressed antibodies to CPn antigens were capable of binding to the surface of the CPn cells. No studies were carried out to test whether antisera against the recombinant FACS positive antigens were capable of interfering with EB *in vitro* infectivity of host cells – that is, whether the murine antibodies raised against the recombinantly expressed antigens could inhibit CPn infectivity *in vitro* to an extent greater than 50%, a property that common practice qualifies such antigens as “neutralising”.

20 Indeed, so far, only few *C. pneumoniae* antigens with ‘neutralizing’ properties have been described in the literature: notably, a protein identified as 76-kDa-homolog protein (Perez-Melgosa *et al* (1994) Infect Immunity 62: 880-6), the surface-exposed outer membrane proteins MOMP (Wolf *et al* (2001) Infect Immun 69: 3082-91), PorB (Kawa *et al* (2002) J Immunol 168 : 5184-91 and Kubo *et al* (2000) Mol Microbiol 38 : 772-80), and very recently also the Pmp21 member of the *Chlamydia*-specific polymorphic family of outer membrane proteins (A.Szczepek, personal communication). All these proteins were in fact selected in the earlier FACS-based screening study (Montigiani *et al* (2002) *ibid*). It can be however noted that outer membrane antigens, as it is the case for MOMP and PorB, could possibly present some kind of practical problems for a recombinant vaccine development project. For instance both MOMP and PorB are integral membrane proteins which appear to require a native conformation to maintain neutralizing epitopes which are discontinuous and conformation-dependent. The production of such proteins may require special processing steps (refolding) which could be undesirable in the preparation of an hypothetical vaccine. Other general problems may arise from the extent of allelic variation, and from regulated proteins which are not always expressed in all *Chlamydial* cell or all *Chlamydial* isolates.

35 Thus, it is desirable to provide improved compositions capable of eliciting an immune response upon exposure to *Chlamydia pneumoniae* proteins. It is also desirable to provide improved compositions comprising one or more combinations of two or more selected CPn proteins with complementary immunological and/or biological profiles capable of providing immunity against *Chlamydial* induced disease and/or infection (such as in prophylactic vaccination) or (b) for the eradication of an established chronic *Chlamydial* infection (such as in therapeutic vaccination).

45 Description of the Invention

The present invention provides compositions comprising a first biological molecule from a *Chlamydia pneumoniae* bacterium and a second biological molecule from a *Chlamydia pneumoniae* bacterium. The term “biological molecule” includes proteins, antigens and nucleic acids. The compositions may also comprise further biological molecules preferably also from *Chlamydia pneumoniae*. That is to say, the compositions may comprise two or more biological molecules (eg. 3, 4, 5, 6, 7, 8 etc.)

at least two of which are from a *Chlamydia pneumoniae* bacterium (eg. 3, 4, 5, 6, 7, 8 etc.). Such compositions include those comprising (i) two or more different *Chlamydia pneumoniae* proteins; (ii) two or more different *Chlamydia pneumoniae* nucleic acids, or (iii) mixtures of one or more *Chlamydia pneumoniae* protein and one or more *Chlamydia pneumoniae* nucleic acid.

In one preferred embodiment the first and second biological molecules are from different *Chlamydia pneumoniae* species (for example, from different *Chlamydia pneumoniae* serovars) but they may be from the same species. The biological molecules in the compositions may be from different serogroups or strains of the same species. The first biological molecule is preferably selected from the group consisting of SEQ ID Nos 1-76. More preferably, it is selected from the group consisting of SEQ IDs 1-41 and/or SEQ ID Nos 42-76. It is preferably a purified or isolated biological molecule. The second biological molecule is preferably selected from the group consisting of SEQ ID Nos 1-76. More preferably, it is selected from the group consisting of SEQ IDs 1-41 and/or SEQ ID Nos 42-76. It is preferably a purified or isolated biological molecule. Specific compositions according to the invention therefore include those comprising: two or more biological molecules selected from the group consisting of SEQ ID Nos 1-41; one or more biological molecules selected from the group consisting of SEQ IDs 1-41 combined with one or more biological molecules selected from the group consisting of SEQ IDs 42-76. One or both of the first and second biological molecules may be a *Chlamydia pneumoniae* biological molecule which is not specifically disclosed herein, and which may not have been identified, discovered or made available to the public or purified before this patent application was filed.

A specific composition of the present invention may comprise a combination of *Chlamydia pneumoniae* antigens, said combination consisting of two, three, four, five or all six *Chlamydia pneumoniae* antigens of a first antigen group, said first antigen group consisting of: (1) pmp2; (2) pmp10; (3) Enolase; (4) OmpH-like protein; and (5) the products of CPn specific genes CPn0759 and CPn0042. These antigens are referred to herein as the 'first antigen group'.

Preferably, the composition of the invention comprises a combination of *Chlamydia pneumoniae* antigens, said combination selected from the group consisting of: (1) pmp2 and pmp10; (2) pmp2 and Enolase; (3) pmp2 and OmpH-like protein; (4) pmp2 and CPn0759; (5) pmp2 and CPn0042; (6) pmp10 and Enolase; (7) pmp10 and OmpH-like protein; (8) pmp10 and CPn0759; (9) pmp10 and CPn0042; (10) Enolase and OmpH-like protein (11) Enolase and CPn0759; (12) Enolase and CPn0042; (13) OmpH-like protein and CPn0759 (14) OmpH-like protein and CPn0042; (15) CPn0759 and CPn0042; (16) pmp2 and pmp10 and Enolase; (17) pmp2 and pmp10 and OmpH-like protein; (18) pmp2 and pmp10 and CPn0759; (19) pmp2 and pmp10 and CPn0042; (20) pmp2 and Enolase and OmpH-like protein; (21) pmp2 and Enolase and CPn0759; (22) pmp2 and Enolase and CPn0042; (23) pmp2 and OmpH-like protein and CPn0759; (24) pmp2 and OmpH-like protein and CPn0042; (25) pmp2 and CPn0759 and CPn0042; and (26) pmp10 and Enolase and OmpH-like protein; (27) pmp10 and Enolase and CPn0759; (28) pmp10 and Enolase and CPn0042; (29) Enolase and OmpH-like protein and CPn0759; (30) Enolase and OmpH-like protein and CPn0042; (31) OmpH-like protein and CPn0759 and CPn0042.

Preferably, the composition of *Chlamydia pneumoniae* antigens consists of pmp2, pmp10, Enolase, OmpH-like protein and CPn0759.

5 Preferably, the composition of *Chlamydia pneumoniae* antigens consists of pmp2, pmp10, Enolase, OmpH-like protein and CPn0042.

Preferably, the composition of *Chlamydia pneumoniae* antigens consists of pmp2, pmp10, Enolase, OmpH-like protein and CPn0759 and CPn0042.

10 The invention also provides for a slightly larger group of 12 *Chlamydia pneumoniae* antigens that are particularly suitable for immunisation purposes, particularly when used in combinations. (This second antigen group includes the six *Chlamydia pneumoniae* antigens of the first antigen group). These 12 *Chlamydia pneumoniae* antigens form a second antigen group of (1) pmp2; (2) pmp10; (3) Enolase; (4) OmpH-like protein; (5) CPn0759; (6) CPn0042; (7) ArtJ; (8) HtrA; (9) AtoS; (10) OmcA; (11) CPn0498; and (12) CPn0525. These antigens are referred to herein as the 'second antigen group'.

20 The invention therefore provides a composition comprising a combination of *Chlamydia pneumoniae* antigens, said combination selected from the group consisting of two, three, four, five, six, seven, eight, nine, ten, eleven, or twelve *Chlamydia pneumoniae* antigens of the second antigen group. Preferably, the combination is selected from the group consisting of two, three, four or five *Chlamydia pneumoniae* antigens of the second antigen group. Still more preferably, the combination consists of six *Chlamydia pneumoniae* antigens of the second antigen group. Each of the *Chlamydia pneumoniae* antigens of the first and second antigen group are described in more detail below.

(1) Pmp10 (CPn0449)

30 One example of a pmp10 protein is set forth as SEQ ID NO: 1 below (GenBank Accession No.GI:14195016). Preferred pmp10 proteins for use with the invention comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 1; and/or (b) which is a fragment of at least *n* consecutive amino acids of SEQ ID NO: 1, wherein *n* is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These pmp2 proteins include variants (e.g. allelic variants, homologs, orthologs, paralogs, mutants, etc.) of SEQ ID NO: 1. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 1. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 1. Other fragments omit one or more domains of the protein (e.g. omission of a signal peptide, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain).

SEQ ID No 1

50 1 MKSQFSWLVL SSTLACFTSC STVFAATAEN IGPSDSFDGS TNTGYTPKN TTTGIDYTLT
61 GDITLQNLGD SAALTKGCFD DTESLSFAG KGYSLSFLNI KSSAEGAALS VTDDKNLSLT
121 GFSSLTFLAA PSSVITTPSG KGAVKCGDGL TFDNNGTILF KQDYCEENG AISTKNLSLK
181 NSTGSISFEG NKSSATGKKG GAICATGTVD ITNNTAPTIF SNNIAEAAGG AINSTGNCTI

241 TGNTSLVFSE NSVTATAGNG GALSGDADVT ISGNQSVTFS GNQAVANGGA IYAKKLTLAS
301 GGGGGISFSN NIVQGTAGN GGAISILAAG ECSSAEAGD ITFNGNAIVA TTPQTTKRNS
361 IDIGSTAKIT NLRAISGHSI FFYDPITANT AADSTDITNL NKADAGNSTD YSGSIVFSGE
421 KLSSEDAKVA DNLSTLTKQP VTLTAGNLVL KRGVTLDTKG FTQTAGSSVI MDAGTTLKAS
481 TEEVTLTGLS IPVDSLGEKG KVVIAASAAS KVALSGPIL LLDNQGNAYE NHDLGKTQDF
541 SFVQLSALGT ATTTDVPAPV TVATPTHYGY QGTWGMTWVD DTASTPKTKT ATLAWTNTGY
601 LPNPERQGPL VPNSLWGSFS DIQAIQGVIE RSALTLCSDR GFWAAGVANF LDKDKKGEKR
661 KYRHKSGGYA IGGAAQTCSE NLISFAFCQL FGSDKDFLVA KNHTDTYAGA FYIQHITECS
721 GFIGCLLDKL PGWSHKLPLV LEGQLAYSHV SNDLTKKYTA YPEVKGSWGN NAFNMLGAS
781 SHSYPEYLHC FDTYAPYIKL NLTYIRQDSF SEKGTEGRSF DDSNLFNLISL PIGVKFEKFS
841 DCNDFSYDLT LSYVPDLIRN DPKCTTALVI SGASWETIAN NLARQALQVR AGSHYAFSPM
901 FEVLGQFVFE VRGSSRIYNV DLGGKQFQ

(2) *Pmp2 = Polymorphic Outer Membrane Protein G Family (CPn 0013)*

One example of a pmp2 protein is disclosed as SEQ ID NO^s: 139 and 140 in WO 02/02606. {GenBank accession number: gi|4376270|gb|AAD18172.1 'CPn0013'; SEQ ID NO: 2 below}. Preferred pmp2 proteins for use with the invention comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 2; and/or (b) which is a fragment of at least *n* consecutive amino acids of SEQ ID NO: 1, wherein *n* is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These pmp2 proteins include variants (e.g. allelic variants, homologs, orthologs, paralogs, mutants, etc.) of SEQ ID NO: 2. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 1. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 2. Other fragments omit one or more domains of the protein (e.g. omission of a signal peptide, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain).

SEQ ID No 2

1 MKIPLRFLLI SLVPTLSMSN LLGAATTEEL SASNSFDGTT STTSFSSKTS
51 SATDGTNYVF KDSVVIENVP KTGETQSTSC FKNDAAAGDL NFLGGGFSFT
101 FSNIDATTAS GAAIGSEAN KVTLSGFSFA LSFLKSPAST VTNGLGAINV
151 KGNLSLLDND KVLIQDNFST GDGGAINCAG SLKIANNKSL SFIGNSSSTR
201 GGAIHTKNLT LSSGGETLFQ GNTAPTAAGK GGAIAIADSG TLSISGDSGD
251 IIFEGNTIGA TGTVSHSAID LGTSAKITAL RAAQGHITIYF YDPITVTGST
301 SVADALNINS PDTGDNKEYT GTIVFSGEKL TEAEAKDEKN RTSKLLQNVA
351 FKNGTVVLKG DVVLSANGFS QDANSKLIMD LGTSLVANTE SIELTNLEIN
401 IDSLRNGKKI KLSAATAQKD IRIDRPVULA ISDESFYQNG FLNEDHSYDG
451 ILELDAGKDI VISADRSID AVQSPYGYQG KWTINWSTDD KKATVSWAQK
501 SFNPTAEQEA PLVPNLLWGS FIDVRSFQNF IELGTEGAPY EKRFFWAGIS
551 NVLHRSGREN QRKFRHVSGG AVVGASTRMP GGDTLSLGFA QLFARDKDYF
601 MNTNFAKTYA GSLRLQHDAS LYSVVSILLG EGGLREILLP YVSKTLPCSF
651 YGQLSYGHTD HRMKTESLPP PPPTLSTDHT SWGGYVWAGE LGTRVAVENT
701 SGRGFFQEYT PFVKVQAVYA RQDSFVELGA ISRDFSDSLH YNLAIPLGIK
751 LEKRFAEQYY HVVAMYSPPV CRSNPKCTTT LLSNQGSWKT KGSNLARQAG
801 IVQASGFRSL GAAAELEFNGF GFEWRGSSRS YNVDAGSKIK F*

(3) *Enolase (Cpn0800)*

One example of an 'Eno' protein is disclosed as SEQ ID NO^s: 93 and 94 in WO 02/02606. {GenBank accession number: gi|4377111|gb|AAD18938.1| 'Cpn0800'; SEQ ID NO: 3 below}. Preferred Eno proteins for use with the invention comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 2; and/or (b) which is a fragment of at least *n* consecutive amino acids of SEQ ID NO: 2, wherein *n* is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40,

50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These Eno proteins include variants (e.g. allelic variants, homologs, orthologs, paralog, mutants, etc.) of SEQ ID NO: 3. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 3. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 3. Other fragments omit one or more domains of the protein (e.g. omission of a signal peptide, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain).

10 SEQ ID No 3

1 MFEAVIADIQ AREILDSRGY PTLHVKVTTT TGSVGEARVP SGASTGKKEA
51 LEFRDTSDFR YQKGVLQAV KNVKEILFPL VKGCSVYEQS LIDSLMMDSD
151 GSPNKETLGA NAILGVSLAT AHAAAATLRR PLYRYLGGCF ACSLPCPMN
151 LINGGMHADN GLEFQEFMIR PIGASSIKEA VMGADVFT LKLLHERGL
201 STGVGDEGGF APNLANSEEA LELLLLAIEK AGFTPGKDIS LALDCAASSF
251 YNVKTGTIDG RHYEEQIAIL SNLCDRYPID SIEDGLAED YDGWALLTEV
301 LGEKVQIVGD DLFVTNPELI LEGISNGLAN SVLIKPNQIG TLTEVYAIK
351 LAQMAGYTTI ISHRSGETTD TTIADLAVAF NAGQIKTGSL SRSERVAKYN
401 RLMEIEEELG SEAFITDSNV FSYEDSEE*

(4) *OmpH-like outer membrane protein (CPn0301)*

One example of 'OmpH-like' protein is disclosed as SEQ ID NO: 77 & 78 in WO 02/02606. {GenBank accession number: gi|4376577|gb|AAD18450.1| 'CPn0301'; SEQ ID NO: 4 below}. Preferred OmpH-like proteins for use with the invention comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 4; and/or (b) which is a fragment of at least n consecutive amino acids of SEQ ID NO: 3, wherein n is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These OmpH-like proteins include variants (e.g. allelic variants, homologs, orthologs, paralog, mutants, etc.) of SEQ ID NO: 4. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 4. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more; preferably 19 or more, to remove the signal peptide) from the N-terminus of SEQ ID NO: 4. Other fragments omit one or more domains of the protein (e.g. omission of a signal peptide as described above, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain).

40 SEQ ID No 4

1 MKLLFSTFL LVLGSTSAAH ANLGYVNLKR CLEESDLGKK ETEELEAMKQ
51 QFVKNAEKIE EELTSIYNKL QDEDYMESLS DSASEELRKK FEDLSGEYNA
101 YQSQQYQSIN QSNVKRIQKL IQEVKIAAES VRSKEKLEAI LNEEAVLAIA
151 PGTDKTTEII AILNESFKKQ N*

(5) *CPn0042 (Hypothetical)*

One example of hypothetical protein is set forth as SEQ ID NO: 5 below. GenBank accession number: gi|4376296|gb|AAD18195.1. Preferred hypothetical proteins for use with the invention comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 5; and/or (b) which is a fragment of at least n consecutive amino acids of SEQ ID NO: 5, wherein n is 7 or

more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These Hypothetical proteins include variants (e.g. allelic variants, homologs, orthologs, paralogs, mutants, etc.) of SEQ ID NO: 5. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 5. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more; preferably 19 or more, to remove the signal peptide) from the N-terminus of SEQ ID NO: 5. Other fragments omit one or more domains of the protein (e.g. omission of a signal peptide as described above, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain).

SEQ ID No 5

1 MEEVSEYLQQ VENQLESCSK RLTKMETFAL GVRLEAKEEI ESIILSDVFN RFEVLCRDIE
61 DMLSRVEEIE RMLRMAELPL LPIKEALTKA FVQHNSCKEK LTKVEPYFKE SPAYLTSEER
121 LQSLNQTLQR AYKESQKVSG LESEVRACRE QLKDQVRQFE TQGVSLIKEE ILFVTSTFRT
181 KFSYHSFRLH VPCMRLYEEY YDDIDLERTR ARWMAMSERY RDAFQAFQEM LKEGLVEEAA
241 ALRETEYWLY REERKSKKKH

(6) CPn0795 (Hypothetical)

One example of hypothetical protein is disclosed as SEQ ID NO^s: 63 & 64 in WO 02/02606. {GenBank accession number: gi|4377106|gb|AAD18933.1| 'CPn0795'; SEQ ID NO: 6 below}. Preferred hypothetical proteins for use with the invention comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 6; and/or (b) which is a fragment of at least *n* consecutive amino acids of SEQ ID NO: 6, wherein *n* is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These Hypothetical proteins include variants (e.g. allelic variants, homologs, orthologs, paralogs, mutants, etc.) of SEQ ID NO: 6. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 6. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more; preferably 19 or more, to remove the signal peptide) from the N-terminus of SEQ ID NO: 6. Other fragments omit one or more domains of the protein (e.g. omission of a signal peptide as described above, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain).

SEQ ID No 6

1 MKDLGTLGGT SSTAKTVSPD GKVIMGRSQI ADGSHWAFMC HTDFSSNNVL
51 FDLNNTYKTL RENGRLNSI FNLQNMMLQR ASDHEFTEFG RSNIALGAGL
101 YVNALQNLPS NLAAQYFGIA YKIRPKYRLG VFLDHNPFSSH VPNNFNVSHN
151 RLWMGAFIGW QSDALGSSV KVSFGYQKQK ATITREQLN TEAGSGESHF
201 EGVAQAIEGR YGKSLGGHVR VQPFLGLQFV HITRKEYTEN AVQFPVHYDP
251 IDYSTGVVYL GIGSHIALVD SLHVGTTRMG M EQNFAAHTDR FSGSIASIGN
301 FVFEKLDVTH TRFAEMRVN YELPYLQSLN LILRVNQQL QGVMGFSSDL
351 RYALGF*

(7) ArtJ arginine periplasmic-binding protein (CPn 0482)

One example of 'ArtJ' protein is disclosed as SEQ ID NO^s: 73 & 74 in WO 02/02606. {GenBank accession number: gi|4376767|gb|AAD18622.1| 'CPn0482'; SEQ ID NO: 7 below}. Preferred ArtJ proteins for use with the invention comprise an amino acid

sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 7; and/or (b) which is a fragment of at least n consecutive amino acids of SEQ ID NO: 7, wherein n is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These ArtJ proteins include variants (e.g. allelic variants, homologs, orthologs, paralogs, mutants, etc.) of SEQ ID NO: 7. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 7. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 7. Other fragments omit one or more domains of the protein (e.g. omission of a signal peptide, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain). The ArtJ protein may be bound to a small molecule like arginine or another amino acid.

15 SEQ ID No 7

1 MIKQIGRFFR AFIFIMPLSL TSCSKIDRN RIWIVGTNAT YPPFEYVDAQ
51 GEVVGFDIDL AKAISEKLGK QLEVREFAFD ALILNLKKHR IDAILAGMSI
201 TPSRQKEIAL LPYYGDEVQE LMVVSKRSL TPVLPLTQYS SVAVQTGTFTQ
151 EHYLLSQPGI CVRSFDSTLE VIMEVRYGKS PVAVLEPSVG RVVLKDFPNL
201 VATRLELPPE CWVLGCGLGV AKDRPEEIQT IQQAITDLKS EGVQSLTKK
WQLSEVAYE*

25 (8) HtrA DO Serine Protease (CPn0979)

One example of an 'HtrA' protein is disclosed as SEQ ID NO⁸: 111 & 112 in WO 02/02606. {GenBank accession number: gi|4377306|gb|AAD19116.1| 'CPn0979'; SEQ ID NO: 8 below}. Preferred HtrA proteins for use with the invention comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 8; and/or (b) which is a fragment of at least n consecutive amino acids of SEQ ID NO: 8, wherein n is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These HtrA proteins include variants (e.g. allelic variants, homologs, orthologs, paralogs, mutants, etc.) of SEQ ID NO: 8. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 8. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more; preferably at least 16 to remove the signal peptide) from the N-terminus of SEQ ID NO: 8. Other fragments omit one or more domains of the protein (e.g. omission of a signal peptide as described above, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain). In relation to SEQ ID NO: 8, distinct domains are residues: 1-16; 17-497; 128-289; 290-381; 394-485; and 394-497.

45 SEQ ID No 8

1 MITKQLRSWL AVLVGSSLLA LPLSGQAVGK KESRVSELPQ DVLLKEISGG
51 FSKVATKATP AVVYIESFPK SQAVTHPSPG RRGPYENPFD YFNDEFNRF
101 FGLPSQREKP QSKEAVRG TG FLVSPDGYIV TNNHVVEDTG KIHVTLHDGQ
151 KYPATVIGLD PKTDLAVIKI KSQNLPLYLSF GNSDHLKVG D WAIAGNPFPG
201 LQATVTVGVI SAKGRNQLHI ADFEDFIQTD AAINPGNSGG PLLNIDGQVI
50 251 GVNTAIVSGS GGYIGIGFAI PSIMANRIID QLIRDGQVTR GFLGVTLQPI
301 DAELAACYKL EKVGALVTD VVGKSPADKA GLKQEDVIA YNGKEVDSL
351 MFRNAVSLMN PDTRIVLKVV REGKVIEIPV TVSQAPKEDG MSALQRVGIR
401 VQNLTPETAK KLGIAPETKG ILIISVEPGS VAASSGIAPG QLILAVNRQK
451 VSSIEDLNRT LKDSNNENIL LMVSQGDVIR FIALKPPE*

(9) AtoS two-component regulatory system sensor histidine kinase protein (CPn0584)

One example of 'AtoS' protein is disclosed as SEQ ID NO^s: 105 & 106 in WO 02/02606. {GenBank accession number: gi|4376878|gb|AAD18723.1| 'CPn0584'; SEQ ID NO: 9 below}. Preferred AtoS proteins for use with the invention comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 9; and/or (b) which is a fragment of at least *n* consecutive amino acids of SEQ ID NO: 9, wherein *n* is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These AtoS proteins include variants (e.g. allelic variants, homologs, orthologs, paralogs, mutants, etc.) of SEQ ID NO: 9. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 9. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 9. Other fragments omit one or more domains of the protein (e.g. omission of a signal peptide, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain).

SEQ ID No 9

20
1 MNVPDSKNLH PPAYELLEIK ARITQSYKEA SAILTAIPDG ILLSETGHF
51 LICNSQAREI LGIDENLEIL NRSFTDVLDPD TCLGFSIQEA LESLKVPKTL
101 RLSLCKESKE KEVELFIRKN EISGYLFIQI RDRSDYKQLE NAIERYKNIA
25 151 ELGKMTATLA HEIRNPLSGI VGFASILKKE ISSPRHQML SSIISGTRSL
201>NNLVSSMLEY TKSQPLNLKI INLQDFFSSL IPLLSVSFPN CKFVREGAQP
251 LFRSIDPDRM NSVVWNLVKN AVETGNSPIT LTLHTSGDIS VTNPGTIPSE
301 IMDKLFTPF TTKREGNGLG LAEAQKIIRL HGGDIQLKTS DSAVSFFIII
351 PELLAALPKE RAAS*

(10) OmcA 9kDa-cysteine-rich lipoprotein(CPn0558)

One example of 'OmcA' protein is disclosed as SEQ ID NO^s: 9 & 10 in WO 02/02606. {GenBank accession number: gi|4376850|gb|AAD18698.1| 'CPn0558', 'OmcA', 'Omp3'; SEQ ID NO: 10 below}. Preferred OmcA proteins for use with the invention comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 10; and/or (b) which is a fragment of at least *n* consecutive amino acids of SEQ ID NO: 10, wherein *n* is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These OmcA proteins include variants (e.g. allelic variants, homologs, orthologs, paralogs, mutants, etc.) of SEQ ID NO: 10. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 10. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more; preferably 18 or more to remove the signal peptide) from the N-terminus of SEQ ID NO: 10. Other fragments omit one or more domains of the protein (e.g. omission of a signal peptide as described above, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain). The protein may be lipidated (e.g. by a *N*-acyl diglyceride), and may thus have a *N*-terminal cysteine.

SEQ ID No 10

1 MKKAVLIAAM FCGVVSLS CRIVDCCFED PCAPSSCNPC EVIRKKERSC
51 GGNACGSYVP SCSNPGSTE CNSQSPQVKG CTSPDGRCKQ *

(11) CPn0498 (Hypothetical)

One example of a hypothetical protein is set forth as SEQ ID NO: 11 below.
(GenBank Accession No. GI:4376784; AAD18638.1). Preferred hypothetical proteins for use with the invention comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 11; and/or (b) which is a fragment of at least n consecutive amino acids of SEQ ID NO: 11, wherein n is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These Hypothetical proteins include variants (e.g. allelic variants, homologs, orthologs, paralog, mutants, etc.) of SEQ ID NO: 11. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 11. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more; preferably 18 or more to remove the signal peptide) from the N-terminus of SEQ ID NO: 11. Other fragments omit one or more domains of the protein (e.g. omission of a signal peptide as described above, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain). The protein may be lipidated (e.g. by a N -acyl diglyceride), and may thus have a N-terminal cysteine.

SEQ ID No 11

1 MNRRKARWV ALFAMTALIS VGCCPWSQAK SRCSDKYIP VVNRLLLEVC LPEAENVEDL
61 IESSSAWVLT PEERFSGELV SICQVKDEHA FYNDLSLLHM TQAVPSYSAT YDCAVVFGGP
121 LPALRQLDF LVREWQRGVR FKKIVFLCGE RGRYQSIEEQ EHFDSRYNP FPTEENWESG
181 NRVTPSSEEE IAKFVWMQML LPRAWRDSTS GVRVTFLAK PEENRVVANR KDTLLLFESY
241 QEAFPGRVLF VSSQFFIGLD ACRVGQFFKG ESYDLAGPGF AQGVLYKHYA PRICLHTLAE
301 WLKETNGCLN ISEGCFF

(12) CPn 0525 (hypothetical)

One example of 'Cpn0525' protein is disclosed as SEQ ID NO^s: 117 & 118 in WO 02/02606. {GenBank accession number: gi|4376814|gb|AAD18665.1| 'CPn0525', SEQ ID NO: 12 below}. Preferred hypothetical proteins for use with the invention comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 12; and/or (b) which is a fragment of at least n consecutive amino acids of SEQ ID NO: 12, wherein n is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These OmcA proteins include variants (e.g. allelic variants, homologs, orthologs, paralog, mutants, etc.) of SEQ ID NO: 12. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 12. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more; preferably 18 or more to remove the signal peptide) from the N-terminus of SEQ ID NO: 12. Other fragments omit one or more domains of the protein (e.g. omission of a signal peptide as described above, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain).

SEQ ID No 12

5
1 MHDALLSILA IQELDIKIR LMRVKKEHQK ELAKVQSLKS DIRRKVQEKE
51 LEMENLKTQI RDGENRIQEI SEQINKLENQ QAAVKKMDEF NALTQEMTTA
101 NKERRSLEHQ LSDLMDKQAG GEDLIVSLKE SLASTENSSS VIEKEIFESI
151 KKINEEGKAL LEQRTCLKHA TNPELLSIYE RLLNNKKDRV VVPIENRVCS
201 GCHIVLTPQH ENLVRKKDRL IFCEHCSRIL YWQESQVNAQ ENSTAKRRRR
251 RAAV*

10 **Third Antigen Group**

15 The immunogenicity of other *Chlamydia pneumoniae* antigens may be improved by combination with two or more *Chlamydia pneumoniae* antigens from either the first antigen group or the second antigen group. Such other *Chlamydia pneumoniae* antigens include a third antigen group consisting of (1) LcrE, (2) DnaK, (3) Omp85 homolog, (4) Mip-like; (5) OmcB (6) MurG (7) Cpn0186 and (8) fliY. These antigens are referred to herein as the "third antigen group".

(13) *LcrE low calcium response E protein (CPn0324)*

20 One example of a 'LcrE' protein is disclosed as SEQ ID NO^s: 29 & 30 in WO 02/02606. {GenBank accession number: gi|4376602|gb|AAD18473.1| 'CPn0324'; SEQ ID NO: 13 below}. Preferred LcrE proteins for use with the invention comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 13; and/or (b) which is a fragment of at least *n* consecutive amino
25 acids of SEQ ID NO: 13, wherein *n* is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These LcrE proteins include variants (e.g. allelic variants, homologs, orthologs, paralogs, mutants, etc.) of SEQ ID NO: 13. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 13. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15,
30 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 13. Other fragments omit one or more domains of the protein (e.g. omission of a signal peptide, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain).

35 **SEQ ID No 13**

40
1 MAASGGTGGL GGTQGVNLAA VEAAAKADA AEVVASQEGS EMNMIQQSQD
51 LTNPAAATRT KKKEEFQTL ESRKKGEAGK AEKKSESTEE KPDTDLADKY
101 ASGNSEISGO ELRGLRDAIG DDASPEDILA LVQEKIKDPA LQSTALDYLV
151 QTTTPSQGKL KEALIQARNT HTEQFGRTAI GAKNILEFASQ EYADQLNVSP
201 SGLRSLYLEV TGDTHCDQL LSMLQDRYTY QDMAIVSSFL MKGMATELKR
251 QGPYVPSAQL QVLMTETRNL QAVLTSYDYF ESRVPILLDS LKAEGIQTPS
301 DLNFVKVAES YHKIINDKFP TASKVEREVR NLIGDDVDSV TGVNLNFFSA
45 351 LRQTSSRLFS SADKRQQLGA MIANALDAVN INNEDYPKAS DFPKPYPWS*

(14) *DnaK heat-shock protein 70 (chaperone) (CPn0503)*

50 One example of 'DnaK' protein is disclosed as SEQ ID NO^s: 103 & 104 in WO 02/02606. {GenBank accessionnumber: gi|4376790|gb|AAD18643.1| 'CPn0503'; SEQ ID NO: 14 below. Preferred DnaK proteins for use with the invention comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 14; and/or (b) which is a fragment of at least *n* consecutive amino acids of SEQ ID NO: 14, wherein *n* is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30,

35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These DnaK proteins include variants (e.g. allelic variants, homologs, orthologs, paralogs, mutants, etc.) of SEQ ID NO: 14. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 14. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 14. Other fragments omit one or more domains of the protein (e.g. omission of a signal peptide, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain).

10 SEQ ID No 14

1 MSEHKKSSKI IGIDLGTNS CVSVMEGGQA KVTSSSEGTR TTPSIVAFKG
51 NEKLVGIPAK RQAVTNPEKT LGSTKRFIGR KYSEVASEIQ TVPYTVTSGS
101 KGDAVFEVDG KQYTPPEIGA QILMKMKETA EAYLGGETVTE AVITVPAYFN
151 DSQRASTKDA GRIAGLDVCR IIEPTAAAL AYGIDKVGDK KIAVFDLGGG
201 TFDISILEIG DGVFEVLSTN GDTLLGGDDF DEVIIKWMIE EFKKQEGIDL
251 SKDNMALQRL KDAAEKAKIE LSGVSSTEIN QPFITMDAQG PKHLALTLTR
301 AQFEKLAASL IERTKSPCIK ALSDAKLSAK DIDDVLLVGG MSRMFAVQET
351 VKELFGKEPN KGVNPDEVVA IGAAIQGGVL GGEVKDVLIL DVIPLSLGIE
401 TLGGVMTTLV ERNTTIPTQK KQIFSTAADN QPAVTIVVLQ GERPMADNK
451 EIGRFDLTDI PPAPRGHPQI EVSFDIDANG IFHVSADVA SGKEQKIRIE
501 ASSGLQDEI QRMVRDAEIN KEEDKKRREA SDAKNEADSM IFRAEKAIKD
551 YKEQIPETLV KEIERIENV RNALKDDAPI EKIKEVTEDL SKHMOKIGES
601 MQSQSASAAA SSAANAKGGP NINTEDLKKH SFSTKPPSNN GSSEDHIEEA
651 DVEIIDNDDK*

(15) Omp85 homolog (Cpn0300)

One example of an Omp85 Homolog protein is disclosed as SEQ ID NO^s: 147 & 148 in WO 02/02606. {GenBank accession number: gi|4376576|gb|AAD18449.1| 'CPn0300'; SEQ ID NO: 15 below}. Preferred Omp85 proteins for use with the invention comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 15; and/or (b) which is a fragment of at least *n* consecutive amino acids of SEQ ID NO: 15, wherein *n* is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These DnaK proteins include variants (e.g. allelic variants, homologs, orthologs, paralogs, mutants, etc.) of SEQ ID NO: 15. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 15. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 15. Other fragments omit one or more domains of the protein (e.g. omission of a signal peptide, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain).

SEQ ID No 15

1 MLIMRNKVIIL QISILALIQT PLTLFSTEKV KEGHVVDSDI TIITEGENAS
51 NKHPLPKLKT RSGALFSQLD FDEDLRILAK EYDSVEPKVE FSEGKTNIAL
101 HLIAPPSIRN IHISGNQVVP EHKILKTLQI YRNDLFEREK FLKGLDDLRT
151 YYLKRGYFAS SVDYSLEHNQ EKGHIDVLIK INEGPCGKIK QLTFSGISRS
201 EKSDIQEFIQ TKQHSITTSW FTGAGLYHPD IVEQDSLAI NYLHNNGYAD
251 AIVNSHYDL DKGNIILYMD IDRGSRYTLG HVHIQGFVFL PKRLIEKQSQ
301 VGPNDLYCPD KIWDGAHKIK QTYAKYGYIN TNVDVLFIP ATRPIYDVTY
351 EVSEGSYPYK GLIKITGNTH TKSDVILHET SLFPGDTFNR LKLEDTEQRL
401 RNTGYFQSVS VYTVRSQDLP MGNADQYRDI FVEVKETTTG NLGLFLGFSS
451 LDNLFGGIEL SESNFDLFGA RNIFSKGFRG LRGGGEHLFL KANFGDKVTD

501 YTLKWTKPHF LNTPWILGIE LDKSINRALS KDYAVQTYGG NVSTTYILNE
551 HLKYGIFYRG SQTSLHEKRR FLLGPNIDSN KGFVSAAGVN LNYDSVDSPP
601 TPTTGIRGGV TFEVSGLGTT YHFTKLSLNS SIYRKLTRKG ILKIKGEAQF
651 IKPYSNTTAE GVPVSEFFFL GGETTVRGYK SFIIGPKYSA TEPQGGLSSL
701 LISEEFQYPL IRQPNISAFV FLDSGFVGLQ EYKISLKDRL SSAGFGLRFD
751 VMNNVPVMLG FGWPFRTTET LNGEKIDVSQ RFFFALGGMF *

(16) Mip-like FKBP-type peptidyl-prolyl cis-trans (CPn0661)

10 One example of a Mip-like protein is disclosed as SEQ ID NO^s: 55 & 56 in WO
02/02606. {GenBank accession number: gi|4376960|gb|AAD18800.1| 'CPn0661';
SEQ ID NO: 16 below}. Preferred Mip-like proteins for use with the invention
comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%,
70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%,
15 99.5% or more) to SEQ ID NO: 16; and/or (b) which is a fragment of at least *n*
consecutive amino acids of SEQ ID NO: 16, wherein *n* is 7 or more (e.g. 8, 10, 12, 14,
16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These mip-
like proteins include variants (e.g. allelic variants, homologs, orthologs, paralogs,
mutants, etc.) of SEQ ID NO: 16. Preferred fragments of (b) comprise an epitope from
20 SEQ ID NO: 16. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3,
4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino
acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ
ID NO: 16. Other fragments omit one or more domains of the protein (e.g. omission
25 of a signal peptide, of a cytoplasmic domain, of a transmembrane domain, or of an
extracellular domain).

SEQ ID No 16

30 1 MNRRWNVLVA TVALALSVAS CDVRSKDKDK DQGSLEVEYKD NKDTNDIELS
51 DNQKLSRTFG HLLARQLRKS EDMFFDIAEV AKGLQAEVC KSAPLTETET
101 EEKMAEVQKL VFEKSKENL SLAEKFLKEN SKNAGVVEVQ PSKLQYKIIK
151 EGAGKAISGK PSALLHYKGS FINGQVFSSS EGNNEPILLP LGQTIPGFAL
201 GMQGMKEGET RVLVTHPDLA YGTAGQLPPN SLLIFEINLI QASADEVAAB
35 251 PQEGNQGE*

(17) OmcB 60 kDa Cysteine rich OMP (CPn0557)

40 One example of an OmcB protein is disclosed as SEQ ID NO^s: 47 & 48 in WO
02/02606. {GenBank accession number: gi|4376849|gb|AAD18697.1| 'CPn0557';
SEQ ID NO: 17 below}. Preferred OmcB proteins for use with the invention
comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%,
70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%,
45 99.5% or more) to SEQ ID NO: 17; and/or (b) which is a fragment of at least *n*
consecutive amino acids of SEQ ID NO: 17, wherein *n* is 7 or more (e.g. 8, 10, 12, 14,
16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These
OmcB proteins include variants (e.g. allelic variants, homologs, orthologs, paralogs,
mutants, etc.) of SEQ ID NO: 17. Preferred fragments of (b) comprise an epitope from
SEQ ID NO: 17. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3,
50 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino
acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ
ID NO: 17. Other fragments omit one or more domains of the protein (e.g. omission
of a signal peptide, of a cytoplasmic domain, of a transmembrane domain, or of an
extracellular domain)..

SEQ ID No 17

1 MSKLIRRVVT VLALTSMA SC FASGGIEAAV AESLITKIVA SAETKPAPVP
51 MTAKKVRLVR RNKQPVEQKS RGAFCDKEFY PCEEGRCQPV EAQQESCYGR
5 LYSVKVNDCC NVEICQSVPE YATVGSPYPI EILAIGKKDC VDVVITQQLP
101 CEAEFVSSDP ETTPSDGKL VWKIDRLGAG DKCKITVWVK PLKEGCCFTA
151 ATVCACPELR SYTKCGQPAI CIKQEGPDCA CLRCPCVYKI EVVNTGSAIA
201 RNVTVDNVPV DGYSHASGQR VLSFNLGDMR PGDKKVFTVE FCPQRRGQIT
251 NVATVTYCGG HKCSANVTV VNEPCVQVNI SGADWSYVCK PVEYSISVSN
10 PGDLVLHDVV IQDTLPSTVT VLEAPGGEIC CNKVWRIKE MCPGETLQFK
351 LVVKAQVPGR FTNQAVTSE SNCCTCTSCA ETTHWKGLA ATHMCVLDTN
401 DPICVGENTV YRICVTNRGS AEDTNVSLIL KFSKELQPIA SSGPTKGTIS
451 GNTVVFDALP KLGSKESEVF SVTLKGIAPG DARGEAILSS DTLTSPVSDT
501 ENTHVY*

(18) MurG peptidoglycan transferase protein (CPn0904)

One example of a 'MurG' protein is disclosed as SEQ ID NO^s: 107 & 108 in WO 02/02606. {GenBank accession number: gi|4377224|gb|AAD19042.1| 'CPn0904';
20 SEQ ID NO: 18 below}. Preferred MurG proteins for use with the invention comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 18; and/or (b) which is a fragment of at least *n* consecutive amino acids of SEQ ID NO: 18, wherein *n* is 7 or more (e.g. 8, 10, 12, 14,
25 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These MurG proteins include variants (e.g. allelic variants, homologs, orthologs, paralogs, mutants, etc.) of SEQ ID NO: 18. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 18. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino
30 acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 18. Other fragments omit one or more domains of the protein (e.g. omission of a signal peptide as described above, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain). The MurG may be lipidated e.g. with undecaprenyl.

SEQ ID No 18

1 MMKKIRKVAL AVGGSGGHIV PALSVKEAFS REGIDVLLLG KGLKNHPSLQ
51 QGISYREIPS GLPTVLNPIK IMSRTLSLCS GYLKARKELK IFDPDLVIGF
40 GSYHSLPVLV AGLSHKIPLF LHEQNLVPGK VNQLFSRYAR GIGVNFSPVT
101 KHFRCPAEV FLPKRSFSLG SPMMKRCTNH TPTICVVGGS QGAQILNTCV
151 PQALVKLVNK YPNMYVHHIV GPKSDVMKVQ HVYNRGEVLC CVKPFEEQLL
201 DVLLAADLVI SRAGATILEE ILWAKVPGIL IPYPGAYGHQ EVNAKFFVDV
251 LEGGTMILEK ELTEKLLVEK VTFALDSHNR EKQRNSLAAY SQQRSTKTFH
301 AFICECL*

(19) CPn0186 (Hypothetical)

One example of a hypothetical protein is set forth as SEQ ID NO: 19 below}.
(GenBank Accession No. GI:4376456; AAD18339.1). Preferred hypothetical
50 proteins for use with the invention comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 19; and/or (b) which is a fragment of at least *n* consecutive amino acids of SEQ ID NO: 19, wherein *n* is 7 or
55 more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These hypothetical proteins include variants (e.g. allelic variants,

homologs, orthologs, paralogs, mutants, *etc.*) of SEQ ID NO: 19. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 19. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 19. Other fragments omit one or more domains of the protein (e.g. omission of a signal peptide, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain).

SEQ ID No 19

10

15

```

1  MSSPVNNTPS APNIPAPPT TPGIPTTKPR SSFIEKVIIV AKYILFAIAA TSGALGTILG
61  LSGALTPGIG IALLVIFVVS MVLGLILKD SISGGEERRL REEVSRTSE NQRLTVITTT
121 LETEVKDLKA AKDQLTEIE AFRNENGNLK TTAEDLEEQV SKLSEQLEAL ERINQLIQAN
181 AGDAQEISSE LKKLISGWDS KVVQINTSI QALKVLLGQE WVQEAQTHVK AMQEIQALQ
241 AEILGMHNQS TALQSVENL LVQDQALTRV VGELLESENK LSQACSALRQ EIEKLAQHET
301 SLQQRIDAML AQEQNLAEQV TALEKMKQEA QKAESEFIAC VRDRTFGRRE TPPPTTPVVE
361 GDESQEEDEG GTPPVSQPSS PVDRTATGQG

```

(20) FltY Glutamine Binding Protein (CPn0604)

One example of a hypothetical protein is set forth as SEQ ID NO⁵: 11 & 12 in WO 02/02606. {GenBank accession number: gi|4376900|gb|AAD18743.1| 'CPn0604'; SEQ ID NO: 20 below}. Preferred hypothetical proteins for use with the invention comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 20; and/or (b) which is a fragment of at least *n* consecutive amino acids of SEQ ID NO: 20, wherein *n* is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These hypothetical proteins include variants (e.g. allelic variants, homologs, orthologs, paralogs, mutants, *etc.*) of SEQ ID NO: 20. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 20. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 20. Other fragments omit one or more domains of the protein (e.g. omission of a signal peptide, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain).

SEQ ID No 20

40

```

1  MKIKFSWKVN FLICLLAVGL IFGCSRVRK EVLVGRDATW FPKQFGIYTS
51  DTNAFLNDLV SEINYKENLN INIVNQDWH LFENLDDKKT QGAFSTVLPT
101 LEMLEHYQFS DPILLTGPVL VVAQDSPYQS IEDLKGRLIG VYKFDSSVLV
151 AQNIPDAVIS LYQHVPIALE ALTSNCYDAL LAPVIEVTAI IETAYKGRLE
201 IISKPLNADG LRLAILKGTN GDLLEGFNAG LVKTRRSKY DAIKQRYRLP

```

The immunogenicity of other *Chlamydia pneumoniae* antigens may be improved by combination with two or more *Chlamydia pneumoniae* antigens from either the first antigen group or the second antigen group or the third antigen group. Such other *Chlamydia pneumoniae* antigens include a fourth antigen group consisting one or more members of the PMP family. These antigens are referred to herein as the "fourth antigen group". Each of the *Chlamydia pneumoniae* antigens of the fourth antigen group is described in more detail below.

Fourth Antigen Group

(21) Polymorphic Membrane Proteins (PMP)

A family of twenty one *Chlamydia pneumoniae* genes encoding predicted polymorphic membrane proteins (PMP) have been identified (*pmp1* to *pmp21*).

15 *Pmp1* (CPn0005)

One example of a *Pmp1* protein is set forth as SEQ ID NO^s: 41 & 42 in WO 02/02606. {GenBank accession number: gi|4376260|gb|AAD18163.1 'CPn0005'; SEQ ID NO: 21 below}.

20 *SEQ ID No 21*

```

1  MRFSLCGFPL VFSFTLLSVF DTSLSATTTIS LTPEDSFHGD SQNAERSYNV
51  QAGDVYSLTG DVSIENVDNS ALNKACFNVT SGSVTFAGNH HGLYFNMISS
101 GTTKEGAVLC CQDPQATARF SGFSTLSFIQ SPGDIKEQGC LYSKNALMLL
151 NNYVVRFEQN QSKTKGGAIS GANVTIVGNY DSVSFYQNA TFGGAIHSSG
201 PLQIAVNQAE IRFAQNTAKN GSGGALYSDG DIDIDQNAVY LFRNEALTT
251 AIGKGGAVCC LPTSGSSTPV PIVTFSDNKQ LVFERNH SIM GGGAIYARKL
301 SISSGGPTLF INNISYANSQ NLGGAIATDT GGEISLSAEK GTITFQGNRT
351 SLPFLNGIHL LQNAKFLKLO ARNGYSIEFY DPITSEADGS TQLNINGDPK
401 NKEYTGITLF SGEKSLANDP RDFKSTIPQN VNLSAGYLVI KEGAEVTVSK
451 FTQSPGSHLV LDLGTKLIAS KEDIAITGLA IDIDSLSSSS TAAVIKANTA
501 NKQISVTDSE ELISPTGNAY EDLRMRNSQT FPLLSLEPGA GGSVTVTAGD
551 FLPVSPHYGF QGNWKLAWTG TGNKVGEFFW DKINYKPRPE KEGNLVFNIL
601 WGNVDVRSLS MQVQETHASS LQTDRLGLWID GIGNFFHVSA SEDNIRYRHN
35 651 SGGYVLNVNN EITPKHYTSM AFSQLFSRDK DYAVSNNEYR MYLGSYLYQY
701 TTSLGNIFRY ASRNPVN NVG ILSRRFLQNP LMIFHFLCAY GHATNDMKT
751 YANFPMVKNS WRNNCWAIEC GGSMPLLVFE NGRLFQGAIP FMKLQLVYAY
801 QGDFKETTAD GRRFSNGSLT SISVPLGIRF EKLALSQDVL YDFSFSYIPD
40 851 IFRKDPSCFA ALVISGDSWL VPAAHVSRHA FVGSGTG RYH FNDYTELLCR
901 GSIECRPHAR NYNINCGSKF RF*

```

Pmp 4 (CPn0017)

One example of a *Pmp 4* protein is designated SEQ ID NO: 22. The sequence for *pmp4* protein can be found at AE001587.1 GI:4376271.

Pmp 6 (CPn0444)

One example of a *Pmp 6* protein is set forth as SEQ ID NO^s 31 & 32 in WO 02/02606. {GenBank accession number: gi|4376727|gb|AAD18588.1| 'CPn0444'; SEQ ID NO: 23 below}.

SEQ ID No 23

```

55 1  MKYSLPWLLT SSALVFSLHP LMAANTDLSS SDNYENGSSG SAAFTAKETS
51  DASGTTYTLT SDVSITNVSA ITPADKSCFT NTGGALSFVG ADHSLVLQTI
101 ALTHDGAAIN NNTALSFSFG FSSLLIDSAP ATGTSGGKGA ICVTNTEGGT

```

5
10
15
20
25

```

151 ATFTDNASVT LQKNTSEKDG AAVSAYSIDL AKTTTAALLD QNTSTKNGGA
201 LCSTANTTVQ GNSGTVTFSS NTATDKGGGI YSKEKDSTLD ANTGVVTFKS
251 NTAKTGGAWS SDDNLALTGN TQVLFQENKT TGSAAQANNP EGCGGAICCY
301 LATATDKTGL AISQNOEMSF TSNTTTANGG AIYATKCTLD GNTTTLTFDQN
351 TATAGCGGAI YTETEDFSLK GSTGTVTFFST NTAKTGGALY SKGNSSLTGN
401 TNLLFSGNKA TGPSNSSANQ EGCGGAILAF IDSGSVSDKT GLSIANNQEV
451 SLTSNAATVS GGAIYATKCT LTGNGSLTFD GNTAGTSGGA IYTETEDFTL
501 TGSTGTVTFFS TNTAKTGGAL YSKGNNSLSG NTNLLFSGNK ATGPSNSSAN
551 QEGCGGAILS FLESASVSTK KGLWIEDNEN VLSGNTATV SGGAIYATKC
601 ALHGNTTLTF DGNTAETAGG AIYTETEDFT LTGSTGTVTF STNTAKTAGA
651 LHTKNTSFT KNKALVFSGN SATATATTTT DQEGCGGAIL CNISEDIAT
701 KSLTLTENES LSFINTAKR SGGGIYAPKC VISGESINF DGNTAETSGG
751 AIYSKNLSIT ANGPVSFTNN SGGKGGAIIYI ADSGELSLEA IDGDITFSGN
801 RATEGTSTPN SIHLGAGAKI TKLAAAPGHT IYFYDPITME APASGGTIEE
851 LVINPVVKAI VPPPQPKNGP IASVPVVPVA PANPNTGTIV FSSGKLPSQD
901 ASIPANTTTI LNQKINLAGG NVVLKEGATL QVYSFTQQPD STVFMDAGTT
951 LETTTTNNND GSIDLKNLSV NLDALDGKRM ITIAVNSTSG GLKISGDLKF
1001 HNNEGSFYDN PGLKANLNL PFLDLSSTSGT VNLDDFNPPI SSMAAPDYGY
1051 QGSWTLVPKV GAGGKVTILVA EWQALGYTPK PELRATLVNP SLWNAVYNIH
1101 SIQOEIATAM SDAPSHPGIW IGGIGNAFHQ DKQKENAGFR LISRGYIVGG
1151 SMTTPQEYTF AVAFSQLEFGK SKDYVVS DIK SQVYAGSLCA QSSYVPLHS
1201 SLRRHVLSKV LPELPGETPL VLHGQVSYGR NHHNMTTKLA NNTQGSQDWD
1251 SHSFAVEVGG SLPVDLNYRY LYSYSPYVKL QVSVNQKGF QEVAEDRIF
1301 DASHLVNVI PMGLTFKHES AKPPSALLLT LGYAVDAYRD HPHCLTSLTN
1351 GTSWSTFATN LSRQAFFAEA SGHLKLLHGL DCFASGSCEL RSSRSYNAN
1401 CGTRYSF*

```

Pmp 7 (CPn0445)

30 One example of a Pmp 7 protein is set forth as SEQ ID NO^s 153 & 154 in WO 02/02606. {GenBank accession number: gi|4376728|gb|AAD18589.1| 'CPn0445'; SEQ ID NO: 24 below}.

SEQ ID No 24

35
40
45
50
55

```

1 MKSSVSWLFF SSIPLFSSLS IVAAEVTLDS SNNSYDGSNG TTFTVFSTTD
51 AAAGTTYSL SDVSFQNAGA LGIPLASGCF LEAGGDLTFQ GNQHALKFAF
101 INAGSSAGTV ASTSAADKNL LFNDFSRLSI ISCPSSLSP TGQCALKSVG
151 NLSLTGNSQI IFTQNFSSDN GGVINTKNFL LSGTSQFASF SRNQFTGKQ
201 GGVVYATGTI TIENSPGIVS FSQNLAKGSG GALYSTDNCS ITDNFQVIFD
251 GNSAWEEAQA QGGAICCTTT DKTVTLTGK NLSFTNTAL TYGGAISGLK
301 VSISAGGPTL FQSNISGSSA GQGGGGAINI ASAGELALSA TSGDITFNNN
351 QVTNGSTSTR NAINIIDTAK VTSIRAATGQ SIYFYDPITN PGTAASTDTL
401 NLNLADANSE IEYGAIVFS GEKLSPTKA IAAVNTSTIR QPAVLARGDL
451 VLRDGVTVTF KDLTQSPGSR ILMGGTTLA AKEANLSLNG LAVNLSLDG
501 TNKAALKTEA ADKNISLSGT IALIDTEGSF YENHNLSAS TYPLLELTTA
551 GANGTITLGA LSTLTLOEPE THYGYQGNWQ LSWANATSSK IGSINWTRTG
601 YIPSPERKSN LPLNSLWGNF IDIRSINQLI ETKSSGEPFE RELWLSGIAN
651 FFYRDSMPTR HGFHRISGGY ALGITATTPA EDQLTFAFCQ LFARDRNHIT
701 GKNHGDYGA SLYFHHTGL FDIANFLWGK ATRAPWVLSE ISQIPLSFD
751 AKFSYLHTDN HMKTYTNDNS IIKGSRWDA FCADLGASLP FVISVPYLLK
801 EVEPFVKVQY IYAHQQDFYE RHAEGRAFNK SELINVEIPI GVTFERDSKS
851 EKGTYDLTLM YILDAYRRNP KCQTSLIASD ANWMAYGTNL ARQGFSVRAA
901 NHFQVNPHEME IFGQFAFEVR SSSRNYNTNL GSKFCF*

```

Pmp 8 (CPn0446)

60 One example of a Pmp 8 protein is set forth as SEQ ID NO^s 45 & 46 in WO 02/02606. {GenBank accession number: gi|4376729|gb|AAD18590.1| 'CPn0446'; SEQ ID NO: 25 below}.

SEQ ID No 25

5 1 MKIPLHKLII SSTLVTPILL SIATYGADAS LSPTDSFDGA GGSTFTPKST
51 ADANGTNYVL SGNVYINDAG KGTALTGCCF TETTGDLTFT GKGYSFNFNT
101 VDAGSNAGAA ASTTADKALT FTGFSNLSFI AAPGTTVASG KSTLSSAGAL
151 NLTDNGTILF SQNVSNNEANN NGGAITTKTL SISGNTSSIT FTSNSAKKLG
201 GAIYSSAAAS ISGNTGQLVF MNKGETGGG ALGFEASSSI TQNSSLFFSG
251 NTATDAAGKG GAIYCEKTGE TPTLTISGNK SLTFAENSSV TQGGAIKAHG
10 301 LDLSAAGPTL FSNNRCGNTA AGKGAIAIA DSGSLSLSAN QGDITFLGNT
351 LTSTSAPTST RNAIYLGSAA KITNLRAAQ QSIYFYDPIA SNTTASDVL
401 TINQPDNSP LDYSGTIVFS GEKLSADEAK AADNFTSILK QPLALASGTL
451 ALKGNVELDV NGFTQTEGST LLMQPGTKLK ADTEAISLTK LVVDLSALEG
501 NKSVSIETAG ANKTITLTSP LVFQDSSGNF YESHTINQAF TQPLVVFATA
15 551 TAASDIYIDA LLTSPVQTPE PHYGYQGHWE ATWADTSTAK SGTMTVWTTG
601 YNPNPERRAS VVPDSLWASF TDIRTLLQIM TSQANSIYQQ RGLWASGTAN
651 FFHKDKSGTN QAFRHKSJGY IVGGSAAEDFS ENIFSVAFQ LFGKDKDLFI
701 VENTSHNYLA SLYLQHRAFL GGLPMPSPFGS ITDMLKDIP ILNAQLSYSY
751 TKNDMDTRYT SYPEAQGSWT NNSGALELGG SLALYLPKEA PFFQGYFPFL
20 801 KFOAVYSRQQ NFKESGAEAR AFDDGDLVNC SIPVGIRLEK ISEDEKNNFE
851 ISLAYIGDVY RKNPRSRTSL MVSGASWTSI CKNLARQAFL ASAGSHLTLS
901 PHVELSGEAA YELRGSATY NVDCGLRYSF *

Pmp 9 (CPn0447)

25 One example of a Pmp 9 protein is set forth as SEQ ID NO^s 33 & 34 in WO 02/02606. {GenBank accession number: gi|4376731|gb|AAD18591.1| 'CPn0447'; SEQ ID NO: 26 below}.

SEQ ID No 26

30 1 MKSSLHWFLI SSSLALPLSL NFSAAFAVVE INLGPTNSFS GPGTYTPPAQ
51 TTNADGTIYN LTGDSVITNA GSPTALTASC FKETGNNLSF QGHGYQFLLQ
101 NIDAGANCTF TNAANKLLS FSGFSYLSLI QTTNATTGTG AIKSTGACSI
151 QSNYSCYFGQ NFSNDNGGAL QGSSISLSLN PNLTFKAKKA TQKGGALYST
35 201 GGITINNTLN SASFSENTAA NNGGAIYEA SSFISNNKAI SFINNSVTAT
251 SATGGAIYCS STSAPKPVLT LSDNGELNFI GNTAITSGGA IYTDNLVLSS
301 GGPTLFKNNS AIDTAAPLGG AIAIADSGSL SLSALGGDIT FEGNTVVKGA
351 SSSQTTTRNS INIGNTNAKI VQLRASQGN IYFYDPITTS ITAALSDALN
401 LNGPDLAGNP AYQGTIVFSG EKLSEAEAAE ADNLKSTIQQ PLTLAGGQLS
40 451 LKSGVTLVAK SFSQSPGSTL LMDAGTTLET ADGITINNLV LNVDLSLKETK
501 KATLKATQAS QVTTLSSGSL LVDPSCNVYE DVSWNNPQVF SCLTLTADDP
551 ANIHITDLAA DPLEKNPIHW GYQGNWALS QEDTATKSKA ATLTWTKTGY
601 NNPERRGTL VANTLWGSFV DVRSIQQLVA TKVRQSQETR GIWCEGISNF
651 FHKDSTKINK GFRHISAGYV VGATTTLASD NLITAAFCQL FGKDRHFIN
701 KNRASAYAAS LHLQHLATLS SPSLLRYLPG SESEQPVLFQ AQISYIYSKN
45 751 TMKTYYTQAP KGESSWYNDG CALELASSLP HTALSHEGLE HAYFPFIKVE
801 ASYIHQDSFK ERNTTLVRSF DSGDLINVS PIGITFERFS RNERASYEAT
851 VIYVADVRYK NPDCTALLI NNTSWKTTGT NLSRQAGIGR AGIFYAFSPN
901 LEVTSNLSME IRGSSRSYNA DLGKGFQF*

50 **Pmp 11 (CPn0451)**

One example of a Pmp 11 protein is set forth as SEQ ID NO^s 115 & 116 in WO 02/02606. {GenBank accession number: gi|4376733|gb|AAD18593.1| 'CPn0451'; SEQ ID NO: 27 below}.

55 **SEQ ID No 27**

60 1 MKTSIPWVLV SSVLAFSCHL QSLANEELLS PDDSFNGNID SGTFTPKTSA
51 TTYSLTGDVF FYEPGKGTPL SDSCFKQTTD NLTFGLNGHS LTFGFIDAGT
101 HAGAAASTTA NKNLTFSGFS LLSFDSSPST TVTTGQGTLS SAGGVNLENI
151 RKLVVAGNFS TADGGAIKGA SFLLTGTSGD ALFSNNSSST KGGAIATTAG
201 ARIANNTGYV RFLSNIASTS GGAIIDDEGTS ILSNNKFLYF EGNAAKTTGG
251 AICNTKASGS PELIISNNKT LIFASNVAET SGGAIHAKKL ALSGGGFTEF

5
10
15

```

301 LRNNVSSATP KGGAISIDAS GELSLSAETG NITFVRNTLT TTGSTDTPKR
351 NAINIGSNGK FTELRAAKNH TIFFYDPITS EGTSSDVLKI NNGSAGALNP
401 YQGTILFSGE TLTADELKVA DNLKSSFTQP VLSGGKLLL QKGVTLSTS
451 FSQEAGSLLG MDSGTTLSTT AGSITITNLG INVDSLGLKQ PVSLTAKGAS
501 NKVIVSGKLN LIDIEGNIYE SHMFSDQLF SLLKITVDAD VDTNVDISSL
551 IPVPAEDPNS EYGFQGWNV NWTDTATNT KEATATWTKT GFVPSPERKS
601 ALVCNTLWGV FTDIRSLQQL VEIGATGMEH KQGFVSSMT NFLHKTGDEN
651 RKGFRTSGG YVIGGSAHTP KDDLFTFAFC HLFARDKDCF IAHNNSRTYG
701 GTLFFKHSHS LQPQNYLRLG RAKFSESAIE KFPREIPLAL DVQVSFSDSD
751 NRMETHYTSL PESEGSWSNE CIAGGIGLDL PFVLSNPHEL FKTFIPQMKV
801 EMVYVQNSF FESSSDGRGF SIGRLNLSI PVGAKFVQGD IGDSYTYDLS
851 GFFVSDVYRN NPQSTATLVM SPDSWKIRGG NLSRQAFLLR GSNVYVNSN
901 CELFGHYAME LRGSSRYNV DVGTKLRF*

```

Pmp 12 (CPn0452)

One example of a Pmp 12 protein is set forth as SEQ ID NO^s 51 & 52 in WO 02/02606. {GenBank accession number: gi|4376735|gb|AAD18594.1 'CPn0452'; SEQ ID NO: 28 below}.

SEQ ID No 28

25
30

```

1  MTILRNFLTC SALFLALPAA AQVVYLHESD GYNGAINNKS LEPKITCYPE
51  GTSYIFLDDV RISNVKHDQE DAGVFINRSG NLFFMGNRCN FTFHNLMTG
101 FGAAISNRVG DTTLTLSNFS YLAFTSAPLL PQQGAIYSL GSVMIENSEE
151 VTFCGNYSW SGAAIYTPYL LGSKASRPV NLSGNRYLVF RDNVSQGYGG
201 AISTHNLTLT TRGPSCFENN HAYHDVNSNG GAIAIAPGGS ISISVKSDDL
251 IFKGNASQD GNTIHNSIHL QSGAQFKNLR AVSESGVYFY DPISHSESHK
301 ITDLVINAPE GKETYEGTIS FSGLCLDDHE VCAENLTSTI LQDVTLAGGT
351 LSLSDGVTLQ LHSFKQEASS TLTMSPGTTL LSCGDARVQN LHILIEDTDN
401 FVPVRIRAEK KDALVSLEKL KVAFEAYWSV YDFPQFKEAF TIPPLELLGP
451 SFDSLLLGET TLERTQVTE NDAVRGFWSL SWEYPPSLD KDRRITPTKK
501 TVFLTWNPEI TSTP*

```

Pmp 13 (CPn0453)

One example of a Pmp 13 protein is set forth as SEQ ID NO^s 3 & 4 in WO 02/02606. {GenBank accession number: gi|4376736|gb|AAD18595.1 'CPn0453'; SEQ ID NO: 29 below}.

SEQ ID No 29

45
50
55
60

```

1  MKTSIRKFLI STTLAPCFAS TAFTVEVIMP SENFDGSSGK IFPYTTLSDP
51  RGTLCIFSGD LYIANLDNAI SRTSSSCFSN RAGALQILGK GGVFSFLNIR
101 SSADGAALSS VITQNPCLP LSFSGFSQMI FDNCESTSD TSASNVPHPA
151 SAIYATTPML FTNNDLILFQ YNRSAGFGAA IRGTSITEN TKKSLFNGN
201 GSISNGGALT GSAAINLINN SAPVIFSTNA TGIYGGAIYL TGGSMITSGN
251 LSGVLVFNNS SRSGGAIYAN GNVTFSSNSD LTFQNNATSP QNSLPAPTTP
301 PTPPAVTPLL GYGGAIFCTP PATPPPTGVS LTISGENSVT FLENIASEQG
351 GALYGGKISI DSNKSTIFLG NTAGKGAIA IPESGELSLS ANQGDILFNK
401 NLSITSGTPT RNSIHFGKDA KPATLGATQG YTLYFYDPIT SDDLASAASAA
451 ATVVVNPKAS ADGAYSGTIV FSGETLTATE AATPANATST LNQKLELEGG
501 TLALRNGATL NVHNFQDEK SVVIMDAGTT LATTNGANNT DGAITLNLKV
551 INLDSLDTGK AAVVNVQSTN GALTISGTLG LVKNSQDCCD NHGMFNKDLQ
601 QVPILELKAT SNTVTTDFD LGTNGYQQSP YGQGTWEFT IDTTHTVTG
651 NWKKTGYLPH PERLAPLIPN SIWANVIDLR AVSQAASADG EDVPGKQLSI
701 GTITNFFHAN HTGDASYRH MGGGYLINTY TRITPDAAALS LGFGQLFTKS
751 KDYLVGHGHS NVYFATVYSN ITKSLFGSSR FFSGGTSRVT YSRSEKVKVT
801 SYTKLPKGRS SWSNNCWLGE LEGNLPITLS SRILNLKQII PFVKAEVAYA
851 THGGIQENTP EGRIFGHGHL LNVAVPVGVR FGKNSHNRPD FYTIIVAYAP
901 DVYRHNPDCE TLLPINGATW TSIGNNLTRS TLLVQASSHT SVNDVLEIFG
951 HCGCDIRRTS RQYTLDIGSK LRF*

```


Pmp 14 (CPn0454)

One example of a Pmp 14 protein is set forth as SEQ ID NO^s 35 & 36 in WO 02/02606. {GenBank accession number: gi|4376737|gb|AAD18596.1 'CPn0454'; SEQ ID NO: 30 below}.

SEQ ID No 30

1 MPLSFKSSSF CLLACLCSAS CAFAETRLGG NRVPPITNQG BEILLTSDFV
51 CSNFLGASFS SSFINSSSNL SLLGKGLSLT FTSCQAPTNS NYALLSAAET
101 LTFKNFSSIN FTGNQSTGLG GLIYKDIVF QSIKDLIFTT NRVAYSPASV
151 TTSATPAITT VTTGASALQP TDSLTVENIS QSIKFFGNLA NFGSAISSSP
201 TAVVKFINNT ATMSFSHNFT SSGGGVIYGG SLLLFENNSG CIIFTANSCV
251 NSLKGVTSSS GTYALGSGGA ICIPGTGFEL KNNQKCTFS YNGTPNDAGA
301 IYAETCNIVG NQGALLDSN TAARNGGAIC AKVLNIQGRG PIEFSRNRAE
351 KGGAIFIGPS VGDPKQTST LTLASEGDI AFQGNMLNTK PGIRNAITVE
401 AGGEIVSLSA QGGSRLVFYD PITHSLPTTS PSNKDITINA NGASGSVVFT
451 SKGLSSTELL LPANTTTILL GTVKIASGEL KITDNAVNVN LGFATQGGSGQ
501 LTLGSGGTLG LATPTGAPAA VDFITGKLAF DPFSFLKRDF VSASVNAGTK
551 NVTLTGALVL DEHDVTDLYD MVSLQTPVAI PIAVFKGATV TKTGFPDGEI
601 ATPSHYGYQG KWSYTWSRPL LIPAPDGGFP GGPSPSANTL YAVWNSDTLV
651 RSTYILDPER YGEIVSNSLW ISFLGNQAFS DILQDVLLID HPGLSITAKA
701 LGAYVEHTPR QGHEGFSGRY GGYQAALSMN YTDHTTLGLS FGQLYGKTNA
751 NPYDSRCSEQ MYLLSFFGQF PIVTQKSEAL ISWKAAYGYS KNHLNTTYLR
801 PDKAPKSQGG WHNNSYYVLI SAEHPFLNWC LLTRPLAQAW DLSGFISAEF
851 LGGWQSKFTE TGDQLRSFSR KGKYNVSLPI GCSSQWFTPF KKAPSTLTIK
901 LAYKPDYIRV NPHNIVTVVS NQESTSISGA NLRRHGLEFVQ IHDVVDLTED
951 TQAFNLNYTFD GKNGFTNHRV STGLKSTF*

Pmp 15 (CPn0466)

One example of a Pmp 15 protein is set forth as SEQ ID NO^s 5 & 6 in WO 02/02606. {GenBank accession number: gi|4376751|gb|AAD18608.1 'CPn0466'; SEQ ID NO: 31 below}.

SEQ ID No 31

1 MRFFCFGMLL PFTFVLANEG LQLPLETYIT LSPEYQAAPQ VGFTHNQNDQ
51 LAIVGNHNDI ILDYKYYSRN GGALTCKNLL ISENIGNVFF EKNVCPNSGG
101 AIYAAQNCTI SKNQNYAFTT NLVSDNPTAT AGSLLGALF AINCSTNNL
151 GQGTFFVDNLA LNKGGALYTE TNLSIKDNKG PIIKQNRAL NSDSLGGGIY
201 SGNSLNIEGN SGAIQITSNS SGSGGGIFST QTLTISSNKK LIEISENSAF
251 ANNYGSNFPN GGGGLTTTFC TILNREGVL FNNNQSQSNG GAIHAKSIII
301 KENGFPVYFLN NTATRGGALL NLSAGSGNGS FILSADNGDI IFNNNTASKH
351 ALNPPYRNAI HSTPNMNLQI GARPGYRVLF YDPIEHELPS SFPILFNFET
401 GHTGTVLFSG EHVHQNFTDE MNFFSYLRNT SELRQGVLA V EDGAGLACYK
451 FFQRGGTLLL GQGAIVTTAG TIPTPSSTPT TVGSTITLNL IAIDLPSILS
501 FQAQAPKIWI YPTKTGSTYT EDSNPTITIS GTLTLRNSNN EDPYDSLDSL
551 HSLEKVPPLY IVDVAAQKIN SSQDLSTLN SGEHYGYQGI WSTYVWETTT
601 ITNPTSLPGA NTKHKLLYAN WSPLGYRPHP ERRGEFITNA LWQSAYTALA
651 GLHSLSSWDE EKGHAASLQG IGLLVHOKDK NGFKGFRSHM TGYSATTEAT
701 SSQSPNFSLG FAQFFSKAKE HESQNSTSSH HYFSGMCIE N TLFKEWIRLS
751 VSLAYMFTSE HTHMYQGLL EGNSQGSFHN HTLAGALSCV FLPQPHGESL
801 QIYPPFITALA IRGNLAAFQE SGDHAREFSL HRPLTDVSLP VGIRASWKNH
851 HRVPLVWLTE ISYRSTLYRQ DPELHSLKLI SQGTWTQAT PVTYNALGIK
901 VKNTMQVFFK VTLSLDYSAD ISSSTLSHYL NVASRMRF*

Pmp 16 (CPn0467)

One example of a Pmp 16 protein is set forth as SEQ ID NO^s 7 & 8 in WO 02/02606. {GenBank accession number: gi|4376752|gb|AAD18609.1| 'CPn0467'; SEQ ID NO: 32 below}.

SEQ ID No 32

```

1  MFGMTPAVYS LQTDLSLEKFA LERDEEFRTS FPLLDLSLSTL TGFSPITTFV
51 GNRHNSSQDI VLSNYKSIDN ILLLWTSAGG AVSCNNFLLS NVEDHAFFSK
101 NLAIGTGGAI ACQGACTITK NRGPLIFFSN RGLNNASTGG ETRGGAIACN
151 GDFTISQNGG TFYFVNNSVN NWGGALSTNG HCRIQSNRAP LLFFNNTAPS
201 GGGALRSENT TISDNTRPIY FKNNCGNNGG AIQTSVTVAI KNNSGSVIFN
251 NNTALSGSIN SGNGSGGAIY TTNLSIDNPN GTILFNNNYC IRDGGAICTQ
301 FLTIKNSGHV YFTNNQGNWG GALMLLDQST CLLFYAEQNI AFQNNNEVFLT
10 351 TFGRYNAIHC TPNSNLQGLA NKGYTATFFD PIEHQHPTTN PLIFNPNANH
401 QGTILFSSAY IPEASDYENN FISSKNTSE LRNGVLSIED RAGWQFYKFT
451 QKGGILKLGH AASIATTANS ETPSTSVGSQ VIINNLAJNL PSILAKGKAP
501 TLWIRPLQSS APFTEDNNPT ITLSGPLTLL NEENRDPYDS IDLSEPLQNI
15 551 HLLSLSDVTA RHINTDNFHP ESLNATEHYG YQGIWSPYVW ETITTTNNAS
601 IETANTLYRA LYANWTPLGY KVNPEYQGDG ATTPWLQSFH TMFSLRLSYN
651 RTGDSIERP FLEIQIADG LFWHQNSIPG APGFRIQSTG YSLQASSETS
701 LHQKISLGFA QFFTRTKBIG SSNNVSAHNT VSSLYVELPW FQEFATSTV
751 LAYGYGDHHL HSLHPSHQEQ AEGTCYSHTL AAAIGCSFPW QQKSYLHLSL
801 FVQAIARSH QTAFEETGDN PRKFVSQKPF YNLTPLGLQ GKWQSKFHVP
20 851 TEWTLLESYQ PVLYQQNPQI GVTLLASGGS WDILGHNYVR NALGYKVHNG
901 TALFRSLDLF LDYQGSVSSS TSTHHLQAGS TLKF*

```

Pmp 18 (CPn0471)

25 One example of a Pmp 18 protein is set forth as SEQ ID No 33 below {GenBank accession number: gi|4376753|gb|AAD18610.1|'CPn0471'.

SEQ ID No 33

```

30 1  MQNNRSLSKS SFFVGALILG KTTILLNATP LSDYFDNQAN QLTTLFPLID TLTNMTYPYSH
61  RATLFGVRDD TNQDIVLDHQ NSIESWFENF SQDGGALSCK SLAITNTKNQ ILFLNSFAIK
121 RAGAMYVNGN FDLSENHCSI IFSGNLSFPN ASNFADTCTG GAVLCSKNVT ISKNQGTAYF
181 INNKAQSSGG AIQAAINIK DNTGPCIFFN NAAGGTAGGA LFANACRIEN NSQPIYFLNN
35 241 QSGLGGAIRV HQECILTKNT GSVIFNNNFA MEADISANHS SGGAIYCISS SIKONPGIAA
301 FDNNTAARDG GAICTQSLTI QDSGPVYFTN NQGTWGGAIM LRQDGACTLF ADQGDIIIFYN
361 NRHFKDTFSN HVSVNCTNRV SLTVGASQGH SATFYDPILQ RYTIQNSIQK FNNPNEHLGT
421 ILFSSTYIPD TSTSRDDFIS HFRNHIGLYN GTLALEDRAE WKVYKFDQFG GTLRLGSRV
481 FSTTDEEQSS SSVGSVININ NLAINLPSIL GNRVAPKLWI RPTGSSAPYS EDNPNPIINLS
40 541 GPLSLLDDEEN LDPYDTADLA QPIAEVPLLY LLDVTAKHIN TDNFYPEGLN TTQHYGYQGV
601 WSPYWIETIT TSDTSSSEDTV NTLHRQLYGD WPTPTGYKVP ENKGDIALLS FWQSFHNLFA
661 TLRYQTQQGQ IAPTASGEAT RLFVHQNSNN DAKGFHMEAT GYSLGTTSTN ASNHSFGVNF
721 SQLFSNLYES HSDNSVASHT TTVALQINNP WLQERFSTSA SLAYSYSNHH IKASGYSGKI
781 QTEGKCYSTT LGAALSCSLS LQWRSRPLHF TPFQIAIAVR SNQTAFOESG DKARKFSVHK
841 PLYNLTVPLG IQSAWESKFR LPTYWNIELA YQPVLYQQNP EINVSLESSG SSWLLSGTTL
45 901 ARNAIAFKGR NQIFIFPKLS VFLDYQGSVS SSTTTHYLHA GTTFKF

```

Pmp 19 (CPn0539)

50 One example of a Pmp 19 protein is set forth as SEQ ID No 34 below {GenBank accession number: gi|4376829|gb|AAD18679.1 'CPn0539'; SEQ ID NO: 34 below}.

SEQ ID No 34

```

55 1  MKQMRLWGFL FLSSFCQVSY LRANDVLLPL SGIHSGEDLE LFTLRSSSPT KTTYSLRKDF
61  IVCDFAGNSI HKPGAFLNL KGDLPFINST PLAALTFKNI HLGARGAGLF SESNVTFKGL
121 HSLVLENNES WGGVLTTSGL LSFINTSVL CQNNISYGPQ GALLLQGRKS KALFFRDNRG
181 TILFLKNKAV NQDESHPGYG GAVSSISPGS PITFADNQEI LFQENEGELG GAIYNDQGA
241 TFENNFOQTS FFSNKASFGG AVYSRYCNLY SQWGDTLFTK NAAAKVGGAI HADYVHIRDC
301 KGSIVFEENS ATAGGAIAYN AVCDINAQGP VRFINNSALG LGGGAIYMQA TGSILRLHAN
361 QGDIEFCGNK VRSQFHSNIN STSNFTNNAI TTQCAPREFS LSANEGHRIC FYDPIISATE
60 421 NYNSLYINHQ RLLEAGGAVI FSGARLSPEH KKENKNKTSI INQPVRLCSG VLSIEGGAIL
481 AVRSFYQEGG LLALGPGSKL TTQGNSEKD KIVITNLGPN LENLDSSDPA EIRATEKASI
541 EISGVPRVYG HTESFYENHE YASKPYTTSI ILSAKKLVT A PSRPEKDIQN LIIAESEYMG

```

601 YGYQGSWEFS WSPNDTKEKK TTIASWTPTG EFSLDPKRRG SFIPITLWST FSGLNIASNI
661 VNNNYLNNSE VIPLQHLVCF GGPVYQIMEQ NPKQSSNNLL VQHAGHNVGA RIPFSFNTIL
721 SAALTQLFSS SSQQNVADKS HAQILIGTVS LNKSWQALSL RSSFSYTEDS QVMKHVFPYK
781 GTSRGSWRNY GWSGSGVMSY AYPKGIRYLK MTPFVDLQYT KLVQNPFFVET GYDPRYFSSS
841 EMTNLSLPIG IALEMRFIGS RSSLFLQVST SYIKDLRRVN PQSSASLVLN HYTWDIQGVF
901 LGKEALNITL NSTIKYKIVT AYMGISSTQR EGSNLSANAH AGLSLSF

Pmp 20 (CPn0540)

One example of a Pmp 20 protein is set forth as SEQ ID NO^s 119 & 120 in WO 02/02606. {GenBank accession number: gi|4376830|gb|AAD18680.1 'CPn0540'; SEQ ID NO: 35 below}.

SEQ ID No 35

1 MKWLPATAVF AAVLPALTAF GDPASVEIST SHTGSGDPTS DAALTGFTQS
51 STETDGTITYT IVGDITFSTF TNIPVPVVTP DANDSSSSNS KGGSSSSSGAT
101 SLIRSSNLHS DFDFTKDSVL DLYHLFFPSA SNTLNPALLS SSSSSGSSSS
151 SSSSSSGSAS AVVAADPKGG AAFYSNEANG TLFTTDSGN PGSLTLQNLK
201 MTGDGAAIYS KGPLVFTGLK NLFTGNESQ KSGGAAYTEG ALTTQAIVEA
251 VTFTGNTSAG QGGAIYVKEA TLFNALDSLK FEKNTSGQAG GGIYTESTLT
301 ISNITKSIEF ISNKASVPAP APEPTSPAPS SLINSTTIDT STLQTRAASA
351 TPAVAPVAHV TPTPISTQET AGNGGAIYAK QGISISTFKD LTFKNSASV
401 DATLTVDSST IGESGGAIFA ADSIQIQCT GTTLFSGNTA NKSGGGIYAV
451 GQVTLLEDIAN LKMTNNTCKG EGCAIYTKKA LTINNGAILT TFSGNTSTDN
501 GGAIFAVGGI TSLDLVEVRF SKNKTGNYSY PITKAASNTA PVVSSSTTAA
551 SPAVPAASAAA PVTNAAKGGG LYSTEGTLVS GITSILSFEN NECQNGGGA
601 YVTKTFQCSH SHRLQFTSNK ADEGGGLYC GDDVTLTNLT GKTLPQENSS
651 EKHGGGLSLA SGKSLTMTSL ESFCLNANTA KENGGGANVP ENIVLTFTYT
701 PTPNEPAPVQ QPVYGEALVT GNTATKSGGG IYTKNAAFSN LSSVTFDQNT
751 SSENGGALLT QKAADKTDCS FTYITNVNIT NNTATGNGGG IAGGKAHFDR
801 IDNLTQSNQ AKKGGGVYLE DALILEKVIT GSVSQNTATE SGGGIYAKDI
851 QLQALPGSFT ITDNKVETSL TTSTNLVGGG IYSSGAVTLT NISGTFGITG
901 NSVINTATSQ DADIQGGGIY ATTSLSINQC NTPILFSNNS AATKKTSTTK
951 QIAGGAIFSA AVTIENNSQP IIFLNNASAKS EATTAATAGN KDSCGGAIAA
1001 NSVTLTNPE ITFKGNIAET GGAIGCIDLT NGSPPRKVS IADNGSVLFQD
1051 NSALNRGGAI YGETIDISRT GATFIGNSSK HDGSAICST ALTLPNSQL
1101 IFENNKVTET TATTKASINN LGAAIYGNNE TSDVTISLSA ENGSIFFKNN
1151 LCTATNKYCS IAGNVKFTAI EASAGKAISF YDAVNVSTKE TNAQELKLE
1201 KATSTGTILF SGELHENKSY IPQKVTFAHG NLILGKNAEL SVVSFTQSPG
1251 TTITMGPGSV LSNHSKEAGG IAINNVIIDF SEIVPTKDNA TVAPPTLKL
1301 SRTNADSKDK IDITGTVTLL DPNGNLYQNS YLGEDRDITL FNIDNSASGA
1351 VTATNVTLQG NLGAKKGYLG TWNLDPNSSG SKIILKWTFD KYLRWPYIPR
1401 DNHFYINSIW GAQNSLVTVK QGILGNMLNN ARFEDPAFNN FWASAIGSFL
1451 RKEVSRNSDS FTYHGRGYTA AVDAKPRQEF ILGAAFSQVF GHAESVHLD
1501 NYKHKGSGHS TOASLYAGNI FYFPAIRSRP ILFQGVATYG YMQHDTTTY
1551 PSIEEKNMAN WDSIAWLFDL RFSVDLKEPQ PHSTARLTFTY TEAEYTRIRQ
1601 EKFTELDYPD RSFSACSYGN LAIPTGFSVD GALAWREIIL YNKVSAAYLP
1651 VILRNPNKAT YEVLSTKEKG NVVNLPTRN AARAEVSSQI YLGSYWTLYG
1701 TYTIDASMNT LVQMANGGIR FVF*

Pmp21 (CPn0963)

One example of a Pmp 21 protein is set forth as SEQ ID NO^s 83 & 84 in WO 02/02606. {GenBank accession number: gi|4377287|gb|AAD19099.1| 'CPn0963'; SEQ ID NO: 36 below}.

SEQ ID No 36

1 MVAKKTVRSY RSSFSHSVIV AILSAGIAFE AHSLSSELD LGVFNKQFEE
51 HSAHVEEAQT SVLKGSDDPVN PSQKESEKVL YTQVPLTQGS SGESLDLADA
101 NFLEHFQHLF EETTVFGIDQ KLVWSDLDTR NFSQPTQEPD TSNAVSEKIS
151 SDTKENRKDL ETEDPSKKSG LKEVSSDLPK SPETAVAAIS EDLEISENIS
201 ARDPLQGLAF FYKNTSSQSI SEKDSSFQGI IFSGSGANS LGFENLKAPK

5
10
15
20
25
30
35
40
45
50
55

251	SGAAVYSDRD	IVFENLVKGL	SFISCESLED	GSAAGVNIVV	THCGDVTLTD
301	CATGLDLEAL	RLVKDFSRGG	AVFTARNHEV	QNNLAGGILS	VVGNGKAIIVV
351	EKNSAEKSNQ	GAFACGSFVY	SNNENTALWK	ENQALSGGAI	SSASDIDIQG
401	NCSAIEFSGN	QSLIALGEHI	GLTDFVGGGA	LAAQGTTLTLR	NNAVVQCCKN
451	TSKTHGGAIL	AGTVDLNETI	SEVAFKQNTA	ALTGGALSAN	DKVIIANNFG
501	EILFEQNEVR	NHGGAIYCGC	RSNPKLEQKD	SGENINIIGN	SGAITFLKNK
551	ASVLEVMTQA	EDYAGGGALW	GHNVLDSNS	GNIQFIGNIG	GSTFWIGEYV
601	GGGAILSTDR	VTISNNSGDV	VFKGNKGQCL	AQKYVAPQET	APVESDASST
651	NKDEKSLNAC	SHGDHYPPKT	VEEEVPPSLI	EEHPVVSSTD	IRGGGAILAQ
701	HIFITDNTGN	LRFSGNLGGG	EESSTVGDLA	IVGGGALLST	NEVNVCSNQN
751	VVFSNDVTSN	GCDSGGAILA	KKVDISANHS	VEFVSNNGSGK	FGGAVCALNE
801	SVNITDNGSA	VSFSKNRTRL	GGAGVAAPQG	SVTICGNQGN	IAFKENFVFG
851	SENQRSGGGA	IIANSSVNIQ	DNAGDILFVS	NSTGSYGGAI	FVGSVLVASEG
901	SNPRTLITIG	NSGDILFAKN	STQTAASLSE	KDSFGGGAIY	TQNLKIVKNA
951	GNVSFYGNRA	PSGAGVQIAD	GGTVCLEAFG	GDILFEGNIN	FDGSFNAIHL
1001	CGNDSKIVEL	SAVQDKNIIF	QDAITYEENT	IRGLPKDVS	PLSAPSLIFN
1051	SKPQDDSAQH	HEGTIRFSRG	VSKIPQIAAI	QEGTLALSQN	AELWLGLKQ
1101	ETGSSIVLSA	GSILRIFDSQ	VDSSAPLPT	NKEETLVSAQ	VQINMSSPTP
1151	NKDKAVDTPV	LADIISITVD	LSSFVPEQDG	TLPLPPEIII	PKGTKLHSNA
1201	IDLKIIDPTN	VGYNHALLS	SHKDIPLISL	KTAEGMTGTP	TADASLSNIK
1251	IDVSLPSITP	ATYGTGVWS	ESKMEDGRIV	VGWQPTGYKL	NPEKQCALVL
1301	NNLWSHYTDL	RALKQEIFAH	HTIAQRMELD	FSTNVWGSGL	GVVEDCQNIQ
1351	EFDGFKHHLT	GYALGLDTQL	VEDFLIGGCF	SQFFGKTESQ	SYKAKNDVKS
1401	YMGAAYAGIL	AGPWLKGAFA	VYGNINNDLT	TDYGTGLGIST	GSWIGKGFIA
1451	GTSIDYRYIV	NPRRFISAIV	STVVPFVEAE	YVRIDLPEIS	EQGKEVRTFQ
1501	KTRFENVAIP	FGFALEHAYS	RGSRAEVNSV	QLAYVFDVYR	KGPVSLITLK
1551	DAAYSWKSYG	VDIPCKAWKA	RLSNNTWNS	YLSTYLAFNY	EWREDLIAYD
1601	FNGGIRIIF*				

30 Preferred PMP proteins for use with the invention comprise an amino acid sequence:
(a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%,
92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to one of the
polypeptide sequences set forth for the pmp proteins above and/or (b) which is a
fragment of at least n consecutive amino acids of one of the polypeptide sequences set
35 forth above wherein n is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50,
60, 70, 80, 90, 100, 150, 200, 250 or more). These PMP proteins include variants (e.g.
allelic variants, homologs, orthologs, paralogs, mutants, etc.) of the polypeptide
sequences set forth above. Preferred fragments of (b) comprise an epitope from one
of the polypeptide sequences set forth above. Other preferred fragments lack one or
40 more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-
terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or
more) from the N-terminus of one of the polypeptide sequences set forth above.
Other fragments omit one or more domains of the protein (e.g. omission of a signal
peptide, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular
45 domain).

Fifth Antigen Group

The immunogenicity of other *Chlamydia pneumoniae* antigens may be improved by
combination with two or more *Chlamydia pneumoniae* antigens from either the first
50 antigen group or the second antigen group or the third antigen group or the fourth
antigen group. Such other *Chlamydia pneumoniae* antigens include a fifth antigen
group consisting one or more cell surface exposed proteins. These antigens are
referred to herein as the "fifth antigen group". Each of the *Chlamydia pneumoniae*
antigens of the fifth antigen group is described in more detail below.
55

(37) PorB Outer Membrane Protein B (CPn0854)

One example of a PorB protein is set forth as SEQ ID NO^s: 67 & 68 in WO 02/02606. {GenBank accession number: gi|4377170|gb|AAD18992.1| 'CPn0854'; SEQ ID NO: 37 below}. Preferred PorB proteins for use with the invention comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 37; and/or (b) which is a fragment of at least *n* consecutive amino acids of SEQ ID NO: 37, wherein *n* is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These PorB proteins include variants (e.g. allelic variants, homologs, orthologs, paralogs, mutants, etc.) of SEQ ID NO: 37. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 37. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 37. Other fragments omit one or more domains of the protein (e.g. omission of a signal peptide, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain).

SEQ ID No 37

20 1 MNSKMLKHLR LATLSFSMFF GIVSSPAVYA LGAGNPAAPV LPGVNPEQTG
 51 WCAFQLCNSY DLFAALAGSL KRGFYGDYVF SESAHITNVP VITSVTTSQT
 101 GTTPTTITSTT KNVDFDLNNS SISSSCVFAT IALQETSPAA IPLLDIAFTA
 151 RVGGLKQYR LPLNAYRDFI SNPLNAEVEV TDGLIEVQSD YGIVWGLSLQ
 201 KVLWKDGVSF VGVSAHYRHG SSPINYITVY NKANPEIYFD ATDGNLSYKE
25 251 WSASIGISTY LNDYVLPYAS VSGNITSRKA PSDSFTLEK QFTNFKFKIR
 301 KITNFDVRNF CFGTTCISN NFYYSVEGRW GYQRATNITS GLQF*

(38) 76kDa Protein Homolog (CPn0728)

One example of a 76kDa Protein Homolog protein is set forth as SEQ ID NO^s: 13 & 14 in WO 02/02606. {GenBank accession number: gi|4377033|gb|AAD18867.1| 'CPn0728'; SEQ ID NO: 38 below}. Preferred 76kDa proteins homologs for use with the invention comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 38; and/or (b) which is a fragment of at least *n* consecutive amino acids of SEQ ID NO: 21, wherein *n* is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These 76kDa protein homologs include variants (e.g. allelic variants, homologs, orthologs, paralogs, mutants, etc.) of SEQ ID NO: 38. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 38. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 38. Other fragments omit one or more domains of the protein (e.g. omission of a signal peptide, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain).

SEQ ID No 38

50 1 MVNPIGPGPI DETERTPPAD LSAQGLEASA ANKSAAEQRI AGAEAKPKES
 51 KTDSVERWSI LRSVAVNAMS LADKLGIASS NSSSTSRSA DVDSTTATAP
 101 TPPPPTFDDY KTQAQTAYDT IFTSTSLADI QAALVSLQDA VTNIKDTAAT
 151 DEETAIAAEW ETKNADAVKV GAQITELAKY ASDNQAILDS LGKLTFSDDL
 201 QAALLQSVAN NNKAAELLKE MQDNPVVPGK TPAIAQSLVD QTDATATQIE
 251 KDGNAIRDAY FAGQNASGAV ENAKSNNSIS NIDSAKAAIA TAKTQIAEAQ
 301 KKFPDSPILQ EAEQMVIAE KDLKNIKPAD GSDVPNPGTT VGGSKQGGSS

5

351 IGSIRVSMML DDAENETASI LMSGFRQMIH MFNTENPDSQ AAQQLAAQA
401 RAKAAGDDS AAAALADAQK ALEAALGKAG QQQGILNALG QIASAAVUSA
451 GVPPAAASSI GSSVKQLYKT SKSTGSDYKT QISAGYDAYK SINDAYGRAR
501 NDATRDVINN VSTPALTRSV PRARTEARGP EKTDQALARV ISGNSRTLGD
551 VYSQVSALQS VMQIIQSNPQ ANNEEIROKL TSAVTKPPQF GYPYVQLSND
601 STQKFIKLE SLFAEGSRTA AEIKALSFET NSLFIQQVLV NIGSLYSGYL
651 Q*

(39) *OmpA conserved outer membrane protein (CPn0695)*

- 10 One example of an OmpA conserved outer membrane protein protein is set forth as
SEQ ID NO^s: 59 & 60 in WO 02/02606. {GenBank accession number:
gi|4376998|gb|AAD18834.1| 'CPn0695'; SEQ ID NO: 39 below}. Preferred ompA
15 proteins for use with the invention comprise an amino acid sequence: (a) having 50%
or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%,
95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 39; and/or (b) which is a
fragment of at least *n* consecutive amino acids of SEQ ID NO: 39, wherein *n* is 7 or
20 more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200,
250 or more). These hypothetical proteins include variants (e.g. allelic variants,
homologs, orthologs, paralogs, mutants, etc.) of SEQ ID NO: 39. Preferred fragments
25 of (b) comprise an epitope from SEQ ID NO: 39. Other preferred fragments lack one
or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-
terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or
more) from the N-terminus of SEQ ID NO: 39. Other fragments omit one or more
domains of the protein (e.g. omission of a signal peptide, of a cytoplasmic domain, of
a transmembrane domain, or of an extracellular domain).

SEQ ID No 39

30 1 MKKLLKSALL SAAFAGSVGS LQALPVGNPS DPSLLIDGTI WEGAAGDPCD
51 PCATWCDAIS LRAGFYGDYV FDRILKVDAP KTFSGMAKPT GSAAANYTTA
101 VDRPNPAYNK HLHDAEWF^{TN} AGFIALNIWD RFDVFCTLGA SNGYIRGNST
151 AFNLVGLFGV KGTTVNANEL PNVSLSNGVV ELYTDTSTFSW SVGARGALWE
201 CGCATLGAEF QYAQSKPKVE ELNVICNVSQ FSVNKP^{KGYK} GVAFFPLPTDA
251 GVATATG^{TKS} ATINYHEWQV GASLSYRLNS LVPYIGVQWS RATFDADNIR
301 IAQPKLPTAV LNLTAWNPSL LGNATALSTT DSFSDFMQIV SCQINKPKSR
351 KACGVTVGAT LVDADKWSLT AEARLINERA AHVSGQFRF*

(40) *PepA (CPn0385)*

- 40 One example of a PepA protein protein is set forth as SEQ ID NO^s: 99 & 100 in WO
02/02606. {GenBank accession number: gi|4376664|gb|AAD18529.1 'CPn0385';
SEQ ID NO: 40 below}. Preferred PepA proteins for use with the invention comprise
an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%,
80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more)
45 to SEQ ID NO: 40; and/or (b) which is a fragment of at least *n* consecutive amino
acids of SEQ ID NO: 40, wherein *n* is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30,
35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These PepA proteins include
variants (e.g. allelic variants, homologs, orthologs, paralogs, mutants, etc.) of SEQ ID
NO: 40 Preferred fragments of (b) comprise an epitope from SEQ ID NO: 40. Other
50 preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15,
20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5,
6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 40. Other
fragments omit one or more domains of the protein (e.g. omission of a signal peptide,
of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain).

SEQ ID No 40

5
10
15
20
25
30
35
40
45

```

1  MVLPHAQASG RNRVKADAIV LPFWHFKDAK NAASFEEFE PSYLPALENF
51 QGKTGEIELL YSSPKAKEKR IVLLGLGKNE ELTSDVVFQT YATLTRVLRK
101 AKCSTVNII PTISELRISA EEFLVGLSSG ILSLNYDYPR YNKVDRNLET
151 PLSKVTVIGI VPKMADAIFR KEAAIFEGVY LTRDLVNRNA DEITPKKLAE
201 VALNLGKEFP SIDTKVLGKD AIAKEKMGLL LAVSKGSCVD PHFIVVRYQG
251 RPKSKDHTVL IGKGVTFDSG GLDLKPGKSM LTMKEDMAGG ATVLGILSAL
301 AVLELPINVT GIIPATENAI DGASYKMGDV YVGMSGLSVE ICSTDAEGRL
351 ILADAITYAL KYCKPTRIID FATLTGAMVV SLGEEVAGFF SNNDVLAEDL
401 LEASAETSEP LWRLPLVKKY DKTLHSDIAD MKNLGSNRAG AITAALFLQR
451 FLEESSVAWA HLDIAGTAYH EKEEDRYPKY ASGFGVRSIL YYLENSLSK*

```

(41) Conserved Outer Membrane Protein (Cpn0278)

One example of a conserved outer membrane protein is set forth as SEQ ID NO: 41 below. GenBank Accession No. GI:4376552; AAD18427.1. Preferred conserved outer membrane proteins for use with the invention comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 41; and/or (b) which is a fragment of at least n consecutive amino acids of SEQ ID NO: 41, wherein n is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These conserved outer membrane proteins include variants (e.g. allelic variants, homologs, orthologs, paralogs, mutants, etc.) of SEQ ID NO: 41. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 41. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 41. Other fragments omit one or more domains of the protein (e.g. omission of a signal peptide, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain).

SEQ ID No 41

35
40

```

1  MKKLSLLVG LIFVLSSCHK EDAQNKIRIV ASPTPHAEEL ESLQEEAKDL GIKLKILPVD
61 DYRIPNRLLL DKQVDANYFQ HQAFLLDECE RYDCKGELVV IAKVHLEPQA IYSKKHSSLE
121 RLKSQKKLTI AIPVDRINAQ RALHLLIECG LIVCKGPANL NMTAKDVCCK ENRSINILEV
181 SAPLLVGSLP DVDAVIPGN FAIAANLSPK KDSLCLLEDLS VSKYTNLVVI RSEDVGSPKM
241 IKLQKLFQSP SVQHFFDTKY HGNILMTQD NG

```

Sixth Antigen Group

The immunogenicity of other *Chlamydia pneumoniae* antigens may be improved by combination with two or more *Chlamydia pneumoniae* antigens from either the first antigen group or the second antigen group or the third antigen group or the fourth antigen group or the fifth antigen group. Such other *Chlamydia pneumoniae* antigens include a sixth antigen group consisting one or more FACS positive CPn antigens. These antigens are referred to herein as the "sixth antigen group". Each of the *Chlamydia pneumoniae* antigens of the sixth antigen group is described in more detail below.

(42) Predicted Omp (CPn0020)

One example of a predicted Omp protein is set forth as SEQ ID NO^s: 91 & 92 in WO 02/02606. {GenBank accession number gi|4376272|gb|AAD18173.1: 'CPn0020'; SEQ ID NO: 42 below}. Preferred Omp proteins for use with the invention comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 42; and/or (b) which is a fragment of at least *n* consecutive amino acids of SEQ ID NO: 42, wherein *n* is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These Omp proteins include variants (e.g. allelic variants, homologs, orthologs, paralogs, mutants, etc.) of SEQ ID NO: 42. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 42. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 42. Other fragments omit one or more domains of the protein (e.g. omission of a signal peptide, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain).

SEQ ID No 42

20	1	MRRCFLFLAS FVLMGSSADA	LTHQEA VKKK	NSYLSHF KSV	SGIVTIEDGV
	51	LNTHMNLRIQ	ANKVYVENTV	GQSLKLVAHG	NVMVNYRAKT
	101	DTDSCLLTNG	RFAMYPWFLG	GSMITLTPET	IVIRKGYIST
	151	SGDYLEYSSD	SLLSIGKTTL	RVCRIPIFL	PPFSIMPMEI
	201	TGGFLGSYL	MSYSPISRKH	FSSTFFLDSF	FKHGVGMGFN
25	251	NVFNMKSYIA	HLAIDMAEA	HDRYRLHGDF	CFTHKHVNFS
	301	TVADIFPNNF	MLKNTGPTRV	DCTWNDNYFE	GYLTSSVKVN
	351	YLTIRQYPIS	IYNTGVYLEN	IVECGYLNFA	FSDHIVGENF
	401	LHKTIVPLPIG	TLSSTLGSSL	IYSDVPEIS	SRHSQLSAKL
	451	SYIQRRHIE	PFVTFITETR	PLAKNEDHYI	FSTQDAFHSL
30	501	VLSKTNPRFP	RIHAKLWTH	ILSNTESKPT	FPKTACELSL
	551	DAEWIWKHC	WDHMNIRWEW	IGNDNVAMTL	ESLHRSKYSL
	601	DVSRPIDQLL	DSPLSDHRNL	ILGKLFVRPH	PCWNYRLSLR
	651	YLEYQMILGT	KIFEHWQLYG	VYERREADSR	FFFFLKLDPK

(43) Predicted Omp (CPn0021)

One example of a predicted Omp protein is set forth as SEQ ID NO^s: 49 & 50 in WO 02/02606. {GenBank accession number gi|4376273|gb|AAD18174.1: 'CPn0021'; SEQ ID NO: 43 below}. Preferred Omp proteins for use with the invention comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 43; and/or (b) which is a fragment of at least *n* consecutive amino acids of SEQ ID NO: 43, wherein *n* is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These hypothetical proteins include variants (e.g. allelic variants, homologs, orthologs, paralogs, mutants, etc.) of SEQ ID NO: 43. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 43. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 43. Other fragments omit one or more domains of the protein (e.g. omission of a signal peptide, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain).

SEQ ID No 43

```

1  MGLFHLTLFG LLLCSLPISL VAKFPESVGH KILYISTOST QQALATYLEA
51 LDAYGDHDFV VLRKIGEDYL KQSIHSSDPQ TRKSTIIGAG LAGSSEALDV
101 LSQAMETADP LQQLLVLSAV SGHLGKTSDD LLEKALASPY PVIRLEAAYR
151 LANLKNTKVI DHLHSFIHKL PEEIQCLSAA IFLRLETEES DAYIRDLLAA
201 KKSARSATA LQIGEQQKR FLPTLRNLLT SASPDQDEAI LYALGKLDKG
251 QSYNIKKQL QKPDVDVTLA AAQALIALGK EEDALPVIKK QALEERPRAL
301 YALRHLPSSEI GIPIALPIFL KTKNSEAKLN VALALLELGC DTPKLEYIT
10 351 ERLVQPHYNE TLALSFSKGR TLQNWKRNI IVPQDPQERE RLLSTTRGLE
401 EQILTFLFRL PKEAYLPCII KLLASQKTQL ATTAISFLSH TSHQEALDLL
451 FQAAKLPGEP IIRAYADLAI YNLTKDPEKK RSLHDIYAKL IQETLLEFVDT
501 ENQRPHPSMP YLRYQVTPES RTKLMLDILE TLATSKSSED IRLLIQLMTE
15 551 GDAKNFPVLA GLLIKIVE*

```

(44) Oligopeptide Binding Protein Oppa-1 Lipoprotein (CPn0195)

One example of an oligopeptide binding protein is set forth as SEQ ID NO^s: 23 and 24 in WO 02/02606. {GenBank accession number gi|4376466|gb|AAD18348.1: 'CPn0195'; SEQ ID NO: 44 below}. Preferred oligopeptide binding proteins for use with the invention comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 44; and/or (b) which is a fragment of at least *n* consecutive amino acids of SEQ ID NO: 44, wherein *n* is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These hypothetical proteins include variants (e.g. allelic variants, homologs, orthologs, paralog, mutants, etc.) of SEQ ID NO: 44. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 44. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 44. Other fragments omit one or more domains of the protein (e.g. omission of a signal peptide, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain).

SEQ ID No 44

```

35 1  MRKISVGICI TILLSLSVVL QGCKESSHSS TSRGELAINI RDEPRSLDPR
51 QVRLLSSEISL VKHIYEGLVQ ENNLSCNIEP ALAEDYSLSS DGLTYTFKLK
101 SAFWSNGDPL TAEDFIESWK QVATQEVSGI YAFALNPIKN VRKIQEGHLS
40 151 IDHFGVHSPN ESTLVVTLES PTSHFLKLLA LPVFFPVHKS QRTLQSKSLP
201 IASGAFYPKN IKQKQWIKLS KNPHYYNQSQ VETKTITTHF IPDANTAACL
251 FNQGKLNWQG PPWGERIPQE TLSNLQSKGH LHSFDVAGTS WLTFFNINKFP
301 LNNMKLREAL ASALDKEALV STIFLGRAKT ADHLLPTNIH SYPEHQKQEM
351 AQRQAYAKKL FKEALEELQI TAKDLEHLNL IFPVSSSASS LLVQLIREQW
45 401 KESLGFAIPI VGKEFALLQA DLSSGNFSLA TGGWFADFAD PMAFLTIFAY
451 PSGVPPYAIN HKDFLEILQN IEQEQDHQKR SELVSQASLY LETFHIIEPI
501 YHDAFQFAMN KKLSNLGVSP TGVVDFRYAK EN*

```

(45) CHLPS 43 kDa Protein Homologue-1 (CPn0562)

One example of a CHLPS protein is set forth as SEQ ID NO: 45 below. GenBank Accession No. GI:4376854; AAD18702.1. Preferred CHLPS proteins for use with the invention comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 45; and/or (b) which is a fragment of at least *n* consecutive amino acids of SEQ ID NO: 45, wherein *n* is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more).

These CHLPS proteins include variants (e.g. allelic variants, homologs, orthologs, paralog, mutants, etc.) of SEQ ID NO: 45. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 45. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 45. Other fragments omit one or more domains of the protein (e.g. omission of a signal peptide, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain).

10 SEQ ID No 45

1 MSIAIAREQY AAILDMHPKP SIAMFSSEQA RTSWEKROAH PYLYRLLEII WGVVKFLILGL
61 IFFIPLGLFW VLQKICQNF I LLGAGGWIFR PICRDSNLLR QAYAARLFSA SFQDHVSSVR
121 RVCLQYDEVF IDGLELRPN AKPDRWMLIS NGNSDCLEYR TVLQGEKDWI FRIAEESSQN
181 ILIFNYPGVM KSQGNITRNN VVKSQACVR YLRDEPAGPQ ARQIVAYGYS LGASVQAEAL
241 SKEIADGSDS VRWFVVKDRG ARSTGAVAKQ FIGSLGVWLA NLTHWNINSE KRSKDLHCPE
301 LFIYKDSQGS NLIGDGLFKK ETCFAAPFLD PKNLEECGSK KIPVAQTGLR HDHILSDDVI
361 KEVAGHIQRH FDN

20

(46) YscJ (Yop translocation J protein) (CPn0828)

One example of a YscJ protein is set forth as SEQ ID NO^s: 109 and 110 in WO 02/02606. {GenBank accession number gi|4377140|gb|AAD18965.1| 'CPn0828'; SEQ ID NO: 46 below. Preferred YscJ proteins for use with the invention comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 46; and/or (b) which is a fragment of at least *n* consecutive amino acids of SEQ ID NO: 46, wherein *n* is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These YscJ proteins include variants (e.g. allelic variants, homologs, orthologs, paralog, mutants, etc.) of SEQ ID NO: 46. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 46. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 46. Other fragments omit one or more domains of the protein (e.g. omission of a signal peptide, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain).

SEQ ID No 46

40 1 MVRRSISFCL FFLMTLLCCT SCNSRSLIVH GLPGREANEI VVLLVSKGVA
51 AQKLPQAAAA TAGAATEQMW DIAVPSAQIT EALAILNQAG LPRMKGTSLL
101 DLFAKQGLVP SELQEKIRYQ EGLSEQMAST IRKMDGVVDA SVQISFTTEN
151 EDNLPLTASV YIKHRGVLDN PNSIMVSKIK RLIAAVPGL VPENVSVD
201 RAAYSBITIN GPWGLTEEID YVSVWGIIA KSSLTKFRLI FYVLILILFV
251 ISCGLLWVIW KTHTLIMTMG GTKGFFNPTP YTKNALEAKK AEGAAADKEK
301 KEDADSQGES KNAETSDKDS SDRDAPEGSN EIEGA*

(47) Hypothetical (CPn 0415)

One example of a hypothetical protein is set forth as SEQ ID NO^s: 101 and 102 in WO 02/02606. {GenBank accession number gi|4376696|gb|AAD18559.1| 'CPn0415'; SEQ ID NO: 47 below}. Preferred hypothetical proteins for use with the invention comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 47; and/or (b) which is a fragment of at

least n consecutive amino acids of SEQ ID NO: 47, wherein n is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These hypothetical proteins include variants (e.g. allelic variants, homologs, orthologs, paralog, mutants, etc.) of SEQ ID NO: 47. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 47. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 47. Other fragments omit one or more domains of the protein (e.g. omission of a signal peptide, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain).

SEQ ID No 47

```

1  MTLIFVITIV WCNAFLIKLC VIMGLQSRLO HCIEVSQNSN FDSQVKQFIY
51 ACQDKTLRQS VLKIFRYHPL LKIHDIARAV YLLMALEEGE DLGLSFLNVQ
101 QYPSGAVELF SCGGFPWKGL PYPAEHAIEFG LLLLQIAEFY EESQAYVSKM
151 SHFQQALFDH QGSVPFSLWS QENSRLLEKEK TTLSQSFLFQ LGMQIHPEYS
201 LEDPALGFWM QTRSSSAFV AASGCQSSLG AYSSGDVGVI AYGPCSGDIS
251 DCYYFGCCGI AKEFVCQKSH QTTEISFLT S TGKPHPRNTG FSYLRDSYVH
301 LPIRCKITIS DKQYRVHAAL AEATSAMTFS IFCKGKNCQV VDGPRLRSCS
351 LDSYKGPGND IMLGENDAI NIVSASPYME IFALQKKEKF WNADFLINIP
401 YKEEGVMLIF EKKVTSEKGR FFTKMN*

```

(48) Hypothetical (CPn0514)

One example of a hypothetical protein is set forth as SEQ ID NO^s: 87 and 88 in WO 02/02606. {GenBank accession number gi|4376802|gb|AAD18654.1| 'CPn0514'; SEQ ID NO: 48 below}. Preferred hypothetical proteins for use with the invention comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 48; and/or (b) which is a fragment of at least n consecutive amino acids of SEQ ID NO: 48, wherein n is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These hypothetical proteins include variants (e.g. allelic variants, homologs, orthologs, paralog, mutants, etc.) of SEQ ID NO: 48. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 48. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 48. Other fragments omit one or more domains of the protein (e.g. omission of a signal peptide, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain).

SEQ ID No 48

```

1  MSNQLQPCIS LGCVSYINSF PLSLQLIKRN DIRCVLAPPA DLLNLLIEGK
51 LDVALTSSLG AISHNLGYVP GFGIAANQRI LSVNLYAAPT FFNSPQPRIA
101 ATLESRSSIG LLKVLCRHLW RIPTPHILRF ITTKVLRQTP ENYDGLLLIG
151 DAALQHPVLP GFVTYDLASG WYDLTKLPFV FALLHSTSW KEHPLPNLAM
201 EEALQQFESS PEEVLKEAHQ HTGLPPSLLQ EYYALCQYRL GEEHYESFEK
251 FREYYGTLYQ QARL*

```

(49) Hypothetical (CPn0668)

One example of a hypothetical protein is set forth as SEQ ID NO^s: 57 and 58 in WO 02/02606. {GenBank accession number gi|4376968|gb|AAD18807.1 'CPn0668'; SEQ ID NO: 49 below}. Preferred hypothetical proteins for use with the invention comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 49; and/or (b) which is a fragment of at least *n* consecutive amino acids of SEQ ID NO: 49, wherein *n* is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These hypothetical proteins include variants (e.g. allelic variants, homologs, orthologs, paralogs, mutants, etc.) of SEQ ID NO: 49. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 49. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 49. Other fragments omit one or more domains of the protein (e.g. omission of a signal peptide, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain).

SEQ ID No 49

20
1 MKFLLYVPLL LVLVSTGCDA KPVSFEPFSG KLSTQRFEPQ HSAEEYFSQG
51 QEFLKKGNFR KALLCFGIIT HHFPRDILRN QAQYLIGVCY FTQDHPDLAD
101 KAFASYLQLP DAEYSEELFQ MKYAIQRFA QGKRKRICRL EGFPKLMNAD
25 151 EDALRIYDEI LTAFPSKDLG AQALYSKAAL LIVKNDLSEA TKTLKKLTLQ
201 FPLHILSSEA FVRLSEIYLQ QAKKEPHNLQ YLHFAKLNEE AMKKQHPNHP
251 LNEVVSANVG AMREHYARGL YATGRFYEKK KKAEEAANIYY RTAITNYPDT
301 LLVAKCQKRL DRISKHTS*

(50) Hypothetical (CPn0791)

30 One example of a hypothetical protein is set forth as SEQ ID NO^s: 123 and 124 in WO 02/02606. {GenBank accession number gi|4377101|gb|AAD18929.1 'CPn0791'; SEQ ID NO: 50 below}. Preferred hypothetical proteins for use with the invention comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 50; and/or (b) which is a fragment of at least *n* consecutive amino acids of SEQ ID NO: 50, wherein *n* is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These hypothetical proteins include variants (e.g. allelic variants, homologs, orthologs, paralogs, mutants, etc.) of SEQ ID NO: 50. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 50. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 50. Other fragments omit one or more domains of the protein (e.g. omission of a signal peptide, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain).

SEQ ID No 50

50 1 MYSCYSKGIS HNYLLHPMSR LDIFVFDSL I ANQDQNLLEE IFCSEDTVLF
51 KAYRTTALQS PLAANKLNIA RKVANYILAD NGEIDTVKLV EAIHHLSQCT
101 YPLGPHRHNE AQDREHLLKM LKALKENPKL KESIKTLFVP SYSTIQNLIR
151 HTLALNPQTI LSTIHVRQAA LTALFTYLRO DVGSCFATAP AILIHQEYPE
201 RFLKDLNDLI SSGKLSRIVN QREIAPVNL SGCIGELFKP LRILDLYPDP

5
10
15

251	LVKLSSSPGL	KKAFSAANLI	ETLGDSEAQI	QQLLSHQYLM	OKLQNVHETL
301	TANDIIKSTL	LHYYQLQEST	VRAIFFKEGL	FSKEQVAFST	QHPRELSEIQ
351	RVYHYLHAYE	EAKSAFIHDT	QNPLLKAWAY	TLATLADASQ	PTISNHIRLA
401	LGWKSEDPHS	LVSLVTHFVE	EEVENIRILV	QQCEQTYHEA	RSQLEYIEGR
451	MRNPLNNQDS	QILTMDHMRF	RQELNKALYE	WDSAQEKAKK	FLHLPEFLLS
501	FYTKQIPLYF	RSSYDAFIQE	FAHLYANAPA	GFRILFTHGR	THPNTWSPYI
551	SINEFIRFLS	EFFTSTESSEL	LGKHAVINLE	KETSRLVHNI	TAMLHTDVFO
601	EALLTRILEA	YQLPVPPSIL	NHLDQLSOTP	WVYVSGGTVD	TLLLDYFESS
651	EPLTLTEKHP	ENPHELAIFY	ADALKDLPFG	IKSYLEEGSH	SLLSSSPTHV
701	FSIIAGSPLF	REAWDNDWYS	YTWLRDVWVK	QHQDFLQDTI	LPQLSIYAFI
751	ENFCNKYALQ	HVVHDFHDFC	SDHSLTLPEL	YDKGSRFLSS	LFTKDKTVAL
801	IYIRRLLYLM	VREVPYVSEQ	QLPEVLDNVS	SYLGISSRIT	YEKFRSLIEE
851	TIPKMTLLSS	ADLRHIYKGL	LMQSYQKIYT	EEDTYLRLTT	AMRHHNLAYP
901	APLLFADSNW	PSIYFGFILN	PGTTEIDLWK	FNAYAGLQGP	LDNIQELFAT
951	SRPWTLYANP	IDYGMPEPPG	YRSRLPKKEF	*	

(51) Hypothetical (CPn0792)

20 One example of a hypothetical protein is set forth as SEQ ID NO⁵: 61 and 62 in WO 02/02606. {GenBank accession number gi|4377102|gb|AAD18930.1| 'CPn0792'; SEQ ID NO: 51 below}. Preferred hypothetical proteins for use with the invention comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 51; and/or (b) which is a fragment of at least *n* consecutive amino acids of SEQ ID NO: 51, wherein *n* is 7 or more (e.g. 8, 10, 12, 14, 25 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These hypothetical proteins include variants (e.g. allelic variants, homologs, orthologs, paralogs, mutants, etc.) of SEQ ID NO: 51. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 51. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or 30 one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 51. Other fragments omit one or more domains of the protein (e.g. omission of a signal peptide, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain).

35 **SEQ ID No 51**

40
45
50

1	MKHTFTKRVL	FFFFLVIPI	LLLNLVVVGF	FSFSAAKANL	VQVLHTRATN
51	LSIEFEKKLT	IHKFLDLRLA	NTLALKSYAS	PSAEPYAQAY	NEMMALSNITD
101	FSLCLIDPFD	GSVRTKNPGD	PFIRYLKQHP	EMKKKLSAAV	GKAFLLTTPG
151	KPLLHYLILV	EDVASWDSTT	TSGLLVSFYP	MSFLQKDLFQ	SLHITKGNIC
201	LVNKYGEVLF	CAQDSESSFV	FSLDLNLPQ	FQARSPSAIE	IEKASGILGG
251	ENLITVSINK	KRYLGLVLNK	IPIQGTYTLS	LVPVSDLIQS	ALKVPLNICF
301	FYVLAFLLMW	WIFSKINTKL	NKPLQELTFC	MEAAWRGNHN	VRFEPPQVGY
351	EFNELGNIFN	CTLLLLLSNI	EKADIDYHSG	EKLQKELGIL	SSLQSALLSP
401	DFPTFPKVT	SSQHLRRRQL	SGHPNGWTVQ	DGGDTLLGII	GLAGDIGLPS
451	YLYALSARSL	FLAYASSDVS	LQKISKDTAD	SFSKTTEGNE	AVVAMTFIKY
501	VEKDRSLELL	SLSEGAPTMF	LQRGESFVRL	PLETHQALQP	GDRLICLTGG
551	EDILKYFSQL	PIEELLKDPL	NPLNTENLID	SLTMMLNNET	EHSADGTLTI
601	LSFS*				

55 **(52) Hypothetical (CPn0820)**

One example of a hypothetical protein is set forth as SEQ ID NO⁵: 113 and 114 in WO 02/02606. {GenBank accession number gi|4377132|gb|AAD18958.1| 'CPn0820'; SEQ ID NO: 52 below}. Preferred hypothetical proteins for use with the invention comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 52; and/or (b) which is a fragment of at

least n consecutive amino acids of SEQ ID NO: 52, wherein n is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These hypothetical proteins include variants (e.g. allelic variants, homologs, orthologs, paralog, mutants, etc.) of SEQ ID NO: 52. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 52. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 52. Other fragments omit one or more domains of the protein (e.g. omission of a signal peptide, of a cytoplasmic domain, of a transmembrane domain; or of an extracellular domain).

SEQ ID No 52

1 MCNSIAMKKQ KRGFVLMELL MSFTLIALLL GTLGFWYRKI YTVQKQKERI
51 YNFYIEESRA YKQLRTLFSM SLSSSYEEPG SLFSLIFDRG VYRDPKLAGA
101 VRASLHHDTK DQRLRLICN IKDQSYFETQ RLLSHVTHV LSFQRNPDPPE
151 KLPETIALTI TREPKAYPPR TLTYQFAVGK*

(53) Hypothetical (CPn0126)

One example of a hypothetical protein is set forth as SEQ ID NO: 53 below. GenBank Accession No. GI:4376390; AAD18279.1 Preferred hypothetical proteins for use with the invention comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 53; and/or (b) which is a fragment of at least n consecutive amino acids of SEQ ID NO: 53, wherein n is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These hypothetical proteins include variants (e.g. allelic variants, homologs, orthologs, paralog, mutants, etc.) of SEQ ID NO: 53. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 53. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 53. Other fragments omit one or more domains of the protein (e.g. omission of a signal peptide, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain).

SEQ ID No 53

1 MVFSYYCMGL FFFSGAIISS GLLVSLGVGL GLSVLGVLLL LLAGLLLFKI QSMLEVPKA
61 PDLLDLEDAS ERLRVKASRS LASLPKEISQ LESYIRSAAN DLNTIKTWPH KDQRLVETVS
121 RKLERLAAQ NYMISELCEI SEILEEEHH LILAQESLEW IGKSLFSTFL DMESFLNLSH
181 LSEVRPYLAV NDRLLEITE ESWEVVSHTI NVTSAFKKAQ ILFKNNEHSR MKKKLESVQE
241 LLETFIYKSL KRSYRELGCL SEKMRIHDN PLFPWVQDQ KYAHAKNEFG BIARCLEEFE
301 KTFFWLDEEC AISYMDCWDF LNESIQNKKS RVDYDITSTK KIALKDRART YAKVLLLENP
361 TTEGKIDLQD AQRATERQSQ EFTYLETET KVRLEALQQC FSDLREATNV RQVRFNTSEN
421 ANDLKESFEK IDKERVRYQK EQLRYWETID RNEQELREEI GESLRLQNRK KGYRAGYDAG
481 RLKGLLRQWK KNLRDVEAHL EDATMDFEHE VSKSELCSVR ARLEVLEEEL MDMSPKVADI
541 BEELSYBERC ILPIRENLER AYLQYNKCE ILSKAKFFFP EDEQLLVSEA NLREVGAQLK
601 QVQKQKQERA QKFAIFEKHI QEOKSLIKEQ VRSFDLAGVG FLKSELLSIA CNLYIKAVVK
661 ESIPVDVPCM QLYYSYEDN EAVVRNRLN MTERYQNFKR SLNSIQFNGD VLLRDPVYQP
721 EGHETRLKER ELQETTLSC KLVQAQDRLS ELESRLSRR

(54) Hypothetical (CPn0794)

One example of a hypothetical protein is set forth as SEQ ID NO: 54 below. GenBank Accession No. GI:4377105; AAD18932.1. Preferred hypothetical proteins for use with the invention comprise an amino acid sequence: (a) having 50% or more

identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 54; and/or (b) which is a fragment of at least n consecutive amino acids of SEQ ID NO: 54, wherein n is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These hypothetical proteins include variants (e.g. allelic variants, homologs, orthologs, paralogs, mutants, etc.) of SEQ ID NO: 54. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 54. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 54. Other fragments omit one or more domains of the protein (e.g. omission of a signal peptide, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain).

SEQ ID No 54

1 MSLYQKWWNS QLKSLCYST VAALIFMIPS QESFADSLID LNLGLDPSVE CLSGDGAFSV
61 GYFTKAGSTP VEYQPFKYDV SKKTFTILSV ETANQSGYAY GISYDGTITV GTCSLGAGKY
121 NGAKWSADGT LTPLTGITGG TSHTEARAIS KDTQVIEGFS YDASGQPKAV QWASGATTVT
181 QLADISGGSR SSYAYAISDD GTIIVGSMES TTRKTTAVK WVNNVPTYLG TLGGDASTGL
241 YISGDGTIVV GAANTATVTN GNQESHAYMY KDNQMKD

(55) Hypothetical (CPn0796)

One example of a hypothetical protein is set forth as SEQ ID NO: 55 below. GenBank Accession No. GI:4377107; AAD18934.1. Preferred hypothetical proteins for use with the invention comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 55; and/or (b) which is a fragment of at least n consecutive amino acids of SEQ ID NO: 55, wherein n is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These hypothetical proteins include variants (e.g. allelic variants, homologs, orthologs, paralogs, mutants, etc.) of SEQ ID NO: 55. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 55. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 55. Other fragments omit one or more domains of the protein (e.g. omission of a signal peptide, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain).

SEQ ID No 55

1 MQPCLNMSIV RNSALPLPCL SRSETFKKVR SHMKFMKVLV PWIYRKDLWV TAFLLTAIPG
61 SFAHTLVDA GEPRHAAQAT GVSGDGKIVI GMKVPDDPFA ITVGFQYIDG HLQPLEAVRP
121 QCSVYPNGIT PDGTVIVGTN YAIGMGSAV KVVNGKVSEL PMLPDTLDSV ASAVSADGRV
181 IGGNRNINLG ASVAVKWEDD VITQLPSLPD AMNACVNGIS SDGSIIIVGTM VDVSWRNTAV
241 QWIGDQLSVI GTLGGTTSVA SAISTDGTVI VGGSENADSQ THAYAYKNGV MSDIGTLGGF
301 YSLAHAVSSD GSVIVGVSTN SEHRYHAFQY ADGQMVDLGT LGGPESYAQG VSGDGKVIIVG
361 RAQVPSGDWH AFLCPFQAPS PAPVHGGSTV VTSQNPRGMV DINATYSSLK NSQQQLQRLI
421 IQHSAKVESV SSGAPSFTSV KGAISKQSPA VQNDVQKGTG LSYSRQVHGN VQSQQLLTGA
481 FMDWKLASAP KCGFKVALHY GSQDALVERA ALPYTEQGLG SSVLSGFGGQ VQGRYDFNLG
541 ETVVLLQPFMG IQVLHLSREG YSEKNVRFPV SYDSVAYSAA TSFMGAHVFA SLSPKMSTAA
601 TLGVERDLNS HIDEFKGSVS AMGNFVLENS TVSVLRPFAS LAMYYDVRRQ QLVTLSVVMN
661 QQPLTGTLST VSQSSYNLSF

(56) Hypothetical (CPn0797)

One example of a hypothetical protein is set forth as SEQ ID NO: 56 below. GenBank Accession No. GI:4377108; AAD18935.1 Preferred hypothetical proteins for use with the invention comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 56; and/or (b) which is a fragment of at least n consecutive amino acids of SEQ ID NO: 56, wherein n is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These hypothetical proteins include variants (e.g. allelic variants, homologs, orthologs, paralogs, mutants, etc.) of SEQ ID NO: 56. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 56. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 56. Other fragments omit one or more domains of the protein (e.g. omission of a signal peptide, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain).

SEQ ID No 56

20 1 MSKKIKVLGH LTLCTLFRGV LCAAALSNIG YASTSQESPY QKSIEDWKGY TFTDLELLSK
61 EGWSEAHAVS GNGSRIVGAS GAGQGSVTAV IWESHLIKHL GTLGGEASSA EGISKDGEV
121 VGWSDTREGY THAFVFDGRD MKDLGTLGAT YSVARGVSGD GSIIVGVSAT ARGEDYGWQV
181 GVKWEKGKIK QLKLLPQGLW SEANAISEDG TVIVGRGEIS RNHIVAVKWN KNAVYSLGTL
241 GGSVASAEAI SANGKVIVGW STTNNGETHA FMHKDETMHD LGTLGGGFSV ATGVSADGRA
25 301 IVGFSAVKTG EIHAFYYAEG EMEDLTTLGG BEARVFDISS EGNDIIGSIK TDAGAERAYL
361 FHIHK

(76) Oligopeptide Binding Protein Oppa-2 Lipoprotein (CPn0196)

30 One example of an oligopeptide binding protein is set forth as SEQ ID NO^s: 127 and 128 in WO 02/02606. {GenBank accession number GI:4376467; AAD18349.1 'CPn0196'; SEQ ID NO: 76 below}. Preferred oligopeptide binding proteins for use with the invention comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 76; and/or (b) which is a fragment of at least n consecutive amino acids of SEQ ID NO: 76, wherein n is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These hypothetical proteins include variants (e.g. allelic variants, homologs, orthologs, paralogs, mutants, etc.) of SEQ ID NO 76. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 76. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 76. Other fragments omit one or more domains of the protein (e.g. omission of a signal peptide, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain).

SEQ ID No 76

50 1 mlrffavfis tlwlitgcs psqsskgifv vnmkemprsl dpgktrliad qtlmrhlyeg
61 lveehsqnge ikpalaesyt isedgtrytf kiknilwsng dpltaqdfvs swkeilkeda
121 ssvlyafip iknaraifdd tespenlgvr aldkrhleiq letpcahflh fltlpiffpv
181 hetlrnysts feempitcga frpvslekgl rhlhleknpmy hnkssrvklhk iivqfisnan
241 taailfkhhk ldwqgppwge pippeisasl hqddqlfslp gasttwllfn iqkqpwnnak
301 lrkalslaid kdmlltkvvyq glaepthdil hprlypgtyp erkrqneril eaqqfleeal

361 delgmtredl eketltfstf sfsygricqm lreqwkkvlk ftipivggef ftigknfleg
421 nysltvnqwt aafidpmsyl mifanpggis pyhlqdsfhq tllikitgeh kkhlnqglii
481 ealdylehch ileplchpnl rialnknkn fnlfvrtsd frfiek1

5

Seventh Antigen Group

The immunogenicity of other *Chlamydia pneumoniae* antigens may be improved by combination with two or more *Chlamydia pneumoniae* antigens from either the first antigen group or the second antigen group or the third antigen group or the fourth antigen group or the fifth antigen group or the sixth antigen group. Such other *Chlamydia pneumoniae* antigens include a seventh antigen group consisting one or more hypothetical proteins (ie proteins which, for example, have no known cellular location and/or function. These antigens are referred to herein as the "seventh antigen group". Each of the *Chlamydia pneumoniae* antigens of the seventh antigen group is described in more detail below.

15

(57) Hypothetical (CPn0331)

One example of a hypothetical protein is set forth as SEQ ID NO: 57 below. GenBank Accession No. GI:4376609; AAD18480.1. Preferred hypothetical proteins for use with the invention comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 57; and/or (b) which is a fragment of at least n consecutive amino acids of SEQ ID NO: 57, wherein n is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These hypothetical proteins include variants (e.g. allelic variants, homologs, orthologs, paralog, mutants, etc.) of SEQ ID NO: 57. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 57. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 57. Other fragments omit one or more domains of the protein (e.g. omission of a signal peptide, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain).

20

25

30

SEQ ID No 57

35

40

45

1	MAVSGGGGVQ	PSSDPGKWNP	ALQGEQAEGP	SPLKESIFSE	TKQASSAAKQ	ESLVRSGSTG
61	MYATESQINK	AKYRKAQDRS	STSPKSKLKG	TFSKMRAVQ	GFMSGFGSRA	SRVSAKRASD
121	SGEGTSLIPT	EMDVALKKGN	RISPQMGGFF	LDASGMGGSS	SDISQLSLEA	LKSSAFSGAR
181	SLSLSSSESS	SVASFGSPQK	AIEPMSEKVV	NAWTVARLGG	EMVSSLLDPN	VETSSLVVRA
241	MATGNEGMID	LSDLGQEEVS	TAMTSPRAVE	GKVKVSSSDS	PEANPTGIPN	SNTLERAKEE
301	AEKQESREQL	SEDQMMLARA	MAGLLTGAAP	QEVLSNSVWS	GPSTVFPPPK	FSGTLPTQRS
361	GDKSKHKSPG	IEKSTNHTNF	SPLREGTVKS	AEVKSPLPHE	SMYRFPKDSI	VSREEPEAVV
421	KESTAFKNPE	NSSQNFLPIA	VESVFPKESG	TGGALGSDAV	SSSYHFLAQR	GVSL LAPLPR
481	ATDDYKEKLE	AHKGPGGPPD	PLIYQYRNVA	VEPPIVLRSP	QPFSGSSRLS	VQGKPEAAASV
541	HDDGGGGNSG	GFSGDQRRGS	SGQKASRQEK	KGKKLSTDI		

(58) Hypothetical (CPn0234)

One example of a hypothetical protein is set forth as SEQ ID NO: 58 below. GenBank Accession No. gi|4376508|gb| AAD18387.1 Preferred hypothetical proteins for use with the invention comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 58; and/or (b) which is a fragment of at least *n* consecutive amino acids of SEQ ID NO: 21, wherein *n* is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These hypothetical proteins include variants (e.g. allelic variants, homologs, orthologs, paralogs, mutants, etc.) of SEQ ID NO: 58. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 58. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 58. Other fragments omit one or more domains of the protein (e.g. omission of a signal peptide, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain).

SEQ ID No 58

1 MLQSCCKALL SIVVSILAFH PIPGMGVEAK SGFLGKVKGW FSKKEIQEEA RILPVKDSLS
61 WKRYDYTSSS GFSVEFPGEF DHSGQIVEVP QSEITIRYDT YVTETHPDNT VYVSVWEYP
121 EKVDISRPEL NLQEGFSGMM QALPESQVLF MQARQIQGHK ALEFWIVCED VYFRGMLISV
181 NHTLYQVFMV YKKNPNQALD KEYEAFSQSF KITKIREPRT IPSSVKKKVS L

(59) Hypothetical (CPn0572)

One example of a hypothetical protein is set forth as SEQ ID NO: 59 below. Genbank Accession No. gi|4376866|gb| AAD18712.1. Preferred hypothetical proteins for use with the invention comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 59; and/or (b) which is a fragment of at least *n* consecutive amino acids of SEQ ID NO: 59, wherein *n* is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These hypothetical proteins include variants (e.g. allelic variants, homologs, orthologs, paralogs, mutants, etc.) of SEQ ID NO: 59. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 59. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 59. Other fragments omit one or more domains of the protein (e.g. omission of a signal peptide, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain).

SEQ ID No 59

1 MAAPINQPS TTTQITQTGQT TTTTIVGSLG EHSVTTTGSG AAAQTSQTVT LIADHEMQEI
61 ASQDGSVASF SAEHSFSTLP PETGSVGATA QSAQSAGLFS LSGRTQRRDS EISSSSDGSS
121 ISRTSSNASS GETSRAESSP DLGDLDSLGS SERAEGAEGP EGPGGLPEST IPHYDPTDKA
181 SILNFLKNPA VQKMQTKGG HFVYVDEARS SFIFVRNGDW STAESIKVSN AKTKENITKP
241 ADLEMCIKAF CVGYETIHS WTGRVKPTME ERSGATGNYN HMLSMKFKT AVVYGPWNAK
301 ESSSGYTTPSA WRRGAKVETG PIWDDVGGLK GINWKTTPAP DFSFINETPG GGAHSTSHTG
361 PGTVPVATVV PNVNVNLGGI KVDLGGINLG GITTNTVTEE GGGTITSTK STSTDDKVS
421 TSTGSQSTIE EDTIQFDDPG QGEDDNAIPG TNTPPPPGPP PNLSSRLLT ISNASLNQVL
481 QNVRQHNLTA YDSNGNSVSD LNQDLGQVVK NSENGVNFPT VILPKTTGDT DPSGQATGGV
541 TEGGGHIRNI IQRNTQSTGQ SEGATPTPQP TIAKIVTSLR KANVSSSSVL PQPQVATTIT
601 PQARTASTST TSIGTGTEST STTSTGTGTG SVSTQSTGVG TPTTTTRSTG TSATTTTSSA
661 STQTPQAPLP SGTRHVATIS LVRNAAGRSI VLQQGGRSQS FPIPPSGTGT QNMGAQLWAA
721 ASQVASTLGQ VVNQAATAGS QPSSRRSSPT SPRRK

Eight Antigen Group

The immunogenicity of other *Chlamydia pneumoniae* antigens may be improved by combination with two or more *Chlamydia pneumoniae* antigens from either the first antigen group or the second antigen group of the third antigen group or the fourth antigen group or the fifth or the sixth antigen group or the seventh antigen group. Such other *Chlamydia pneumoniae* antigens include an eighth antigen group consisting one or more FACS positive CPn antigens. These antigens are referred to herein as the "eight antigen group". Each of the *Chlamydia pneumoniae* antigens of the eight antigen group is described in more detail below.

(60) Low Calcium Response Protein H (CPn0811)

One example of a Low Calcium Response Protein H is set forth as SEQ ID NO: 60 below. Genbank Accession No. GI:4377123; AAD18949.1. Preferred low calcium response proteins for use with the invention comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 60; and/or (b) which is a fragment of at least *n* consecutive amino acids of SEQ ID NO: 60, wherein *n* is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These low calcium response proteins include variants (e.g. allelic variants, homologs, orthologs, paralogs, mutants, etc.) of SEQ ID NO: 60. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 60. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 60. Other fragments omit one or more domains of the protein (e.g. omission of a signal peptide, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain).

SEQ ID No 60

```

1 mskpsprnan qpqksasfn kktrsr lael aaqkkakadd leqvhvppte eeikkalgni
61 feglslgldl qqilglsdyl leeiytvayt fysqgkynea vglfqllaaa qpqnykymlg
121 lsscyhqlhl yneaaqgffl afdaqpndpi ppyyiadsll klqqpeesnn fldvtmdicg
181 nnpefkilke rcqimkqsie kqmagetkka ptkkpagksk tttknksgkk r

```

(61) Yop Proteins Translocation Protein T (CPn0823)

One example of a Yop Proteins Translocation Protein T is set forth as SEQ ID NO: 61 below. Genbank Accession No. GI:4377135; AAD18960.1. Preferred Yop proteins for use with the invention comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 61; and/or (b) which is a fragment of at least *n* consecutive amino acids of SEQ ID NO: 61, wherein *n* is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These Yop proteins include variants (e.g. allelic variants, homologs, orthologs, paralogs, mutants, etc.) of SEQ ID NO: 61. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 61. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 61. Other fragments omit one or more domains of the protein (e.g. omission of a signal peptide, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain).

SEQ ID No 61

5
1 mgislpelfs nlgsayldyi fqhppayvws vflllllarll pifavapflg aklfpfpiki
61 gislswlaaii fpkvladtqi tnymdnnlfy vllvkemiig ivigfvlaflp fyaaqsagsf
121 itnqqgiqgl egatslisie qtsphgilyh yfvttiifwlv gghrvisl1 lqtlevipih
181 sffpaemmsl sapiwitmik mcqlclvmti qlsapaalam lmsdlflgii nrmappqvvi
241 yllsalkafm glflftlaww fiikqidyft lawfkevpm llgsnpqvl

(62) Yop Proteins Translocation Protein J

10 One example of a Yop Proteins Translocation Protein J is set forth as SEQ ID NO: 62 below Genbank Accession No. GI:4377140; AAD18965.1. Preferred hypothetical proteins for use with the invention comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 62; and/or (b) which is a
15 fragment of at least n consecutive amino acids of SEQ ID NO: 62, wherein n is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These hypothetical proteins include variants (e.g. allelic variants, homologs, orthologs, paralog, mutants, etc.) of SEQ ID NO: 62. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 62. Other preferred fragments lack one
20 or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 62. Other fragments omit one or more domains of the protein (e.g. omission of a signal peptide, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain).

SEQ ID No 62

30
1 mvrrsisfcl fflmtllcct scnsrslivh glpgreanei vlllvskgva aqklpqaaaa
61 tagaateqmw diavpsaqit ealailnqag lprmkgtsl1 dlakqglvp selqekiryq
121 eglseqmast irkmdgvvda svqisftten ednlpltasv yikhrgvldn pnsimvskik
181 rliasavpgl vpenvsvvsd raaysditin gpwglteeid yvsvwgiila kssltkfrli
241 fyvlililfv iscgllwviw kthtlimtmg gtkgffnptp ytknaleakk aegaaadkek
301 kedadsqges knaetsdkds sdkdapegsn eiega

(63) OmpA (CPn0695)

35 One example of an OmPA encoded (MOMP) protein is set forth as SEQ ID NO: 63 below Genbank Accession No. GI:4376998; AAD18834.1. Preferred OmpA proteins for use with the invention comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%,
40 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 63; and/or (b) which is a fragment of at least n consecutive amino acids of SEQ ID NO: 63, wherein n is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These OmpA proteins include variants (e.g. allelic variants, homologs, orthologs, paralog, mutants, etc.) of SEQ ID NO: 63. Preferred fragments of (b)
45 comprise an epitope from SEQ ID NO: 63. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 63. Other fragments omit one or more domains of the protein (e.g. omission of a signal peptide, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain).

SEQ ID No 63

1 mkkllksall saafagsvgs lqalpvgngps dpsllidgti wegaagdpcd pcatwcdais
5 61 lragfygdyv fdrilkvdp ktfsmgakpt gsaaanytta vdrpnpaynk hlhdaewftn
121 agfialniwd rfdvftlga sngyirngst afnlvglfgv kgttvanel pnvslsngvv
181 elytdtsfsw svrgargalwe cgcattlgaef qyagskpkve elnvicnvsq fsvnkpkyk
241 gvafplptda gvatatgtks atinyhewqv gaslsyrlns lvpvlgvqws ratfdadnir
301 iaqpkltav lnlawnpsl lgnatalstt dsfsdfmqiv scqinkfksr kacgvtvgat
10 361 lvdadkwslt aearlinera ahvsgqfrf

(64) Hypothetical (CPn0210)

One example of a Hypothetical Protein is set forth as SEQ ID NO: 64 below Genbank
Accession No. GI:4376482; AAD18363.1. Preferred hypothetical proteins for use
with the invention comprise an amino acid sequence: (a) having 50% or more identity
15 (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%,
98%, 99%, 99.5% or more) to SEQ ID NO: 64; and/or (b) which is a fragment of at
least n consecutive amino acids of SEQ ID NO: 64, wherein n is 7 or more (e.g. 8, 10,
12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more).
These hypothetical proteins include variants (e.g. allelic variants, homologs,
20 orthologs, paralogs, mutants, etc.) of SEQ ID NO: 64. Preferred fragments of (b)
comprise an epitope from SEQ ID NO: 64. Other preferred fragments lack one or
more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-
terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or
25 more) from the N-terminus of SEQ ID NO: 64. Other fragments omit one or more
domains of the protein (e.g. omission of a signal peptide, of a cytoplasmic domain, of
a transmembrane domain, or of an extracellular domain).

SEQ ID No 64

30 1 mlvelealkr efahlkdqkp tsdqeitsly qclldhlevl lglgqdkflk atededvlfe
61 sqkaidawna lltkardvlg lgdigaiyqt ieflgaylsk vnrrafcias eihflktair
121 dlnayylldf rwplckieef vdwgndcvei akrklctfek etkelnesll reehamekcs
181 iqdlqrklsl iielhdvsl fcfsktspse eyqkdcllys rlyrllllye ytllicktstd
241 fgeqarakee firekfslll lekqikqtke lefaiakskl ergclvmrky eaaakhslds
35 301 mfeeetvksp rkdte

(65) Low Calcium Response Locus Protein H (CPn1021)

One example of a Low Calcium Response Protein H is set forth as SEQ ID NO: 65
below Genbank Accession No. GI:4377352; AAD19158.1. Preferred low calcium
response proteins for use with the invention comprise an amino acid sequence: (a)
40 having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%,
93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 65; and/or
(b) which is a fragment of at least n consecutive amino acids of SEQ ID NO: 65,
45 wherein n is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90,
100, 150, 200, 250 or more). These low calcium response proteins include variants
(e.g. allelic variants, homologs, orthologs, paralogs, mutants, etc.) of SEQ ID NO: 65.
Preferred fragments of (b) comprise an epitope from SEQ ID NO: 65. Other preferred
fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or
50 more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8,
9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 65. Other fragments
omit one or more domains of the protein (e.g. omission of a signal peptide, of a
cytoplasmic domain, of a transmembrane domain, or of an extracellular domain).

SEQ ID No 65

1 mshlnyllek iaasskedfp fpddlesyle gyvpdknial dtyqkifkis sedlekvyke
5 61 gyhayldkdy aksitvfrwl vffnfpvskf wfslgaslmh segysqalha ygvtaavlrdk
121 dpyphyayyi cytltnheee aekalemawv raqhkplyne lkeeildirk hk

Ninth Antigen Group

The immunogenicity of other *Chlamydia pneumoniae* antigens may be improved by combination with two or more *Chlamydia pneumoniae* antigens from either the first antigen group or the second antigen group or the third antigen group or the fourth antigen group or the fifth antigen group or the sixth antigen group or the seventh antigen group or the eighth antigen group. Such other *Chlamydia pneumoniae* antigens include a ninth antigen group. These antigens are referred to herein as the "ninth antigen group". Each of the *Chlamydia pneumoniae* antigens of the ninth antigen group is described in more detail below.

(66) Low Calcium Response Protein D (CPn0323)

One example of a Low Calcium Response Protein D is set forth as SEQ ID NO: 66 below Genbank Accession No. GI:4376601; AAD18472.1. Preferred low calcium response proteins for use with the invention comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 66; and/or (b) which is a fragment of at least *n* consecutive amino acids of SEQ ID NO: 66, wherein *n* is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These low calcium response proteins include variants (e.g. allelic variants, homologs, orthologs, paralogs, mutants, etc.) of SEQ ID NO: 66. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 66. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 66. Other fragments omit one or more domains of the protein (e.g. omission of a signal peptide, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain).

SEQ ID No 66

35 1 mnklfnfvsr tlggdtalnm inkssdlila lwmmgvvlmi iiplpppivd lmitinlsis
61 vfillmvalyi psalqlsvfp slllittmfr lginissrqr illkayaghv iqafgdfvvg
121 gnyvvgfiif liitiiqfiv vtkgaervae vaarfildam pgkqmaidad lragmidatq
40 181 ardakraiqk eselygamdg amkfikgdvi agivislini vggltigvam hgmldlaqaah
241 vytllsigdg lvsqipslli altagivttr vssdkntnlg keistqlvke pralllagaa
301 tlgvgffkqf plwsfsilal ifvalgilil tkksaagkkg gsgasttvg aagdgatvvg
361 dnpddysltl pvilelgkdl skliqhktks ggsfvddmip kmrqalyqdi girypgihr
421 tdspalegyd ymillnevyv vrgkipphhv ltnevednls rynlpfityk naaglpasawv
481 sedakailek aaikywtple viilhllyff hkssqeflgi qevrsmiefm ersfpdlvke
45 541 vtrliplqkl teifkrlvqe qisikdlrti leslsewaqt ekdtvlltey vrsslklyis
601 fkfsggqsai svylldepeie emirgaikqt sagsylaldp dsvnlilksm rntitptpag
661 gqppviltai dvrryvrkli etefpdiaivi syqeilpeir iqplgriqif

(67) CHLPS 43kDa Protein Homolog-1 (CPn0062)

One example of a CHLPS 43kDa Protein Homolog-1 is set forth as SEQ ID NO: 67 below Genbank Accession No. GI:4376318; AAD18215.1. Preferred CHLPS proteins for use with the invention comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 67; and/or (b) which is a

fragment of at least n consecutive amino acids of SEQ ID NO: 67, wherein n is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These CHLPS proteins include variants (e.g. allelic variants, homologs, orthologs, paralog, mutants, etc.) of SEQ ID NO: 67. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 67. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 67. Other fragments omit one or more domains of the protein (e.g. omission of a signal peptide, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain).

SEQ ID No 67

```

15      1 mmskrtski avlsilltft hsigfanans svglgtvyit sevvkpkqkg serkqakkep
      61 rarkgylvps srtlsaraqk mknssrkess ggcneisans tprsvklrrn kraeqkaakq
      121 gfsafsnltl kslpklpks qktsiherek atsrfvnesq lssarkryct pssaapslfl
      181 eteivrapve rtkelqndei hipvvqvqtn pkeqntkttk qlasqasiqq segteqslre
      241 laggaslpvl vrsnpevsq rkeellkel vaerrqckrk svrgalears ltkkvarggs
      301 vtstlrydpe kaaeiksrn ckvspeareq kyssckrdar angkqdkttp sedasqeeqq
      361 tgaglvrrkt ksquvasnqn fyrrskntni dsyltanqys csseetdwpc sscvskrrth
      421 nsisvctmvv tviamivgal iianatesqt tsdtpptpt p

```

(68) Hypothetical (CPn0169)

One example of a CHLPS 43kDa Protein Homolog-1 is set forth as SEQ ID NO: 68 below Genbank Accession No. GI:4376437; AAD18322.1. Preferred CHLPS proteins for use with the invention comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 68; and/or (b) which is a fragment of at least n consecutive amino acids of SEQ ID NO: 68, wherein n is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These CHLPS proteins include variants (e.g. allelic variants, homologs, orthologs, paralog, mutants, etc.) of SEQ ID NO: 68. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 68. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 68. Other fragments omit one or more domains of the protein (e.g. omission of a signal peptide, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain).

SEQ ID No 68

```

      1 mknvgsecsq plvmeintqp lrlncsrlv kitsfviall alvggitlta lagagilsfl
      61 pwlvlgivlv vlcalflifs ykfcpikeg vvyntdsqih qwfqkqrnk lekatenpel
      121 fgenraednn rsarsqvket lrdcdgnvlk kiyernldvl lfmnwvpktm dddvpvseds
      181 irtviscykl ikackpefrs lisellramq sglgllsrsc ryqeraktvs hkdaplfcpt
      241 hsyrrdgytl plragpryii nrai

```

(69) PmpD family (CPn0963)

One example of a PmpD protein is set forth as SEQ ID NO: 69 below Genbank Accession No. GI:4377287; AAD19099.1. Preferred PmpD proteins for use with the invention comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 69; and/or (b) which is a fragment of at

least n consecutive amino acids of SEQ ID NO: 69, wherein n is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These PmpD proteins include variants (e.g. allelic variants, homologs, orthologs, paralogs, mutants, etc.) of SEQ ID NO: 69. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 69. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 69. Other fragments omit one or more domains of the protein (e.g. omission of a signal peptide, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain).

SEQ ID No 69

```

15      1 mvakktvrsy rssfshsviv ailsagiafe ahsllhseld lgvfnkqfee hsahveeaqt
        61 svlkgdvpv psqkesekvl ytcvplttgs sgesldlada nfhlefqhlf eettvfgidq
121     121 klwvsdldtr nfsqptqepd tsnavsekis sdtkenrkd1 etedpskksq lkevssdlpk
181     181 spetavaais edleisenis ardplggla fykntssqsi sekdsffggi ifsgsgansg
241     241 lgfenlkapk sgaavysdrd ivfenlvkgl sfiscesled gsaagvniiv thcgdvtltd
20      301 catgldleal rlvkdfsrqg avftarnhev qnnlaggils vvgknkaivv eknsaeksng
        361 gafacgsfvy snnentalwk enqalsggai ssasdidigg ncsaiefsgn qslialgehi
421     421 gltdfvvgga laaggtltlr nnavvcvkn tskthggail agtvdlneti sevafkqnta
481     481 altggalsan dkviianfng eilfegnevr nhggaiycgc rsnpkleqkd sgeniniign
541     541 sgaitflknk asvlevmtqa edyagggalw ghnvllsds gniqfignig gstfwigeyv
601     601 gggailstdr vtisnnsqdv vfkgnkgqcl aqkyvapqet apvesdasst nkdekslnac
25      661 shgdhyppkt veeevppsl1 eehpvvsstd irgggailaq hifitdntgn lrfsgnlggg
        721 eesstvgdla ivgggallst nevnvcnqn vvfndvtsn gcdsggaila kkvdisanhs
781     781 vefvnsngsk fggavcalne svnitdngsa vsfsknrtrl ggagvaapqg svticgnqgn
841     841 iafkenfvfg senqrsqgga iianssvniq dnagdilfvs nstgsgyggai fvgslvaseg
901     901 snprtltitg nsqdilfakn stqtaaslse kdsfgggaiy tqnlkivkna gnvsvfygnra
30      961 psgagvqiad ggtvcleafg gdilfegnin fdgsfnaihl cgndskivel savqdkniif
1021    1021 qdaityeent irglpdkdvs plsapslifn skpqddsaqh hegtirfsrg vskipqiaai
1081    1081 qegtlalsqn aelwlaglkg etgssivlsa gsilrifdsq vdssaplpte nkeetlvsag
1141    1141 vqinmssptp nkdkavdtpv ladiisitvd lssfvpeqdg tlpplpeiii pkgtklhsna
35      1201 idlkiidptn vgyenhalls shkdiplisl ktaegmtgtp tadaslsnik idvslpsitp
        1261 atyghtgvws eskmedgrlv vgwqptgykl npekqgalvl nnlwshytdl ralkqEIFah
1321    1321 htiaqrmeld fstnvwsgsl gvvedcqnig efdgfkhhlt gyalgltdtql vedfliggcf
1381    1381 sqffgktesq sykakndvks ymgaayagil agpwlikgaf vvgminndlt tdygtlgist
1441    1441 gswigkgfia gtsidyryiv nprrfisaiv stvvpfveae yvridlpeis eggkevrtfq
40      1501 ktrfenvaip fgfalehays rgsraevnsv qlayvfdvyr kgpvsllitlk daayswksyg
        1561 vdipckawka rlsnntewns ylstylafny ewredliayd fnggiriif

```


Tenth Antigen Group

The immunogenicity of other *Chlamydia pneumoniae* antigens may be improved by combination with two or more *Chlamydia pneumoniae* antigens from either the first antigen group or the second antigen group or the third antigen group or the fourth antigen group or the fifth antigen group or the sixth antigen group or the seventh antigen group or the eighth antigen group or the ninth antigen group. Such other *Chlamydia pneumoniae* antigens include a tenth antigen group. Each of the *Chlamydia pneumoniae* antigens of the tenth antigen group is described in more detail below.

(70) OmpH-like outer membrane protein (CPn0301)

One example of 'OmpH-like' protein is disclosed as SEQ ID NO: 77 & 78 in WO 02/02606. {GenBank accession number: gi|4376577|gb|AAD18450.1| 'CPn0301'; SEQ ID NO: 70 below and SEQ ID No 4 above}. Preferred OmpH-like proteins for use with the invention comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 4; and/or (b) which is a fragment of at least n consecutive amino acids of SEQ ID NO: 3, wherein n is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These OmpH-like proteins include variants (e.g. allelic variants, homologs, orthologs, paralogs, mutants, etc.) of SEQ ID NO: 4. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 4. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more; preferably 19 or more, to remove the signal peptide) from the N-terminus of SEQ ID NO: 4. Other fragments omit one or more domains of the protein (e.g. omission of a signal peptide as described above, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain).

SEQ ID No 70

```

1  MKKLLFSTFL LVLGSTSAAH ANLGYNLKR CLEESDLGKK ETEELEAMKQ
51 QFVKNAEKIE EELTSIYNKL QDEDMESLS DSASEELRKK FEDLSGEYNA
101 YQSQYYQSIN QSNVKRIQKL IQEVKIAAES VRSKEKLEAI LNEEAVLAIA
151 PGTDKTTTEII AILNESFKKQ N*

```

(71) L7/L12 Ribosomal Protein (CPn0080)

One example of an L7/L12 Ribosomal protein is set forth as SEQ ID No 71 below {GenBank accession number: GI:4376338; AAD18233.1}. 'CPn0080'; SEQ ID NO: 71 below. Preferred L7/L12 proteins for use with the invention comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 71; and/or (b) which is a fragment of at least n consecutive amino acids of SEQ ID NO: 71, wherein n is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These L7/L12 ribosomal proteins include variants (e.g. allelic variants, homologs, orthologs, paralogs, mutants, etc.) of SEQ ID NO: 71. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 71. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more; preferably 19 or more, to remove the signal peptide) from the N-terminus of SEQ ID NO: 71. Other fragments omit one

or more domains of the protein (e.g. omission of a signal peptide as described above, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain).

SEQ ID No 71

5

1 mttesletlv eklsnltvle lsqklklee kwdvtasapv vavaagggge apvaaeptef
61 avtledvpad kkigvlkvvr evtglalkea kemteglpkt vkektksda edtvkklqda
121gakasfkg1

10

(72) AtoS two-component regulatory system sensor histidine kinase protein (CPn0584)

One example of 'AtoS' protein is disclosed as SEQ ID NO^s: 105 & 106 in WO 02/02606. {GenBank accession number: gi|4376878|gb|AAD18723.1| 'CPn0584'; SEQ ID NO: 72 below and SEQ ID No 9 above}. Preferred AtoS proteins for use with the invention comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 72; and/or (b) which is a fragment of at least *n* consecutive amino acids of SEQ ID NO: 72, wherein *n* is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These AtoS proteins include variants (e.g. allelic variants, homologs, orthologs, paralogs, mutants, etc.) of SEQ ID NO: 72. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 72. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 72. Other fragments omit one or more domains of the protein (e.g. omission of a signal peptide, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain).

SEQ ID No 72

30

1 MNVPDSKNLH PPAYELLEIK ARITQSYKEA SAILTAIPDG ILLSETGCHF
51 LICNSQAREI LGIDENLEIL NRSFTDVLPD TCLGFSIQEA LESLKVPKTL
101 RLSLCKESKE KEVELFIRKN EISGYLFIQI RDRSDYKQLE NAIERYKNIA
151 ELGKMTATLA HEIRNPLSGI VGFASILKKE ISSPRHQML SSIISGTRSL
201>NNLVSSMLEY TKSQPLNLKI INLQDFSSL IPLLSVSFPN CKFVREGAQP
251 LFRSIDPDRM NSVVWNLVKN AVETGNPIT LTLHTSGDIS VTNPGTIPSE
301 IMDKLFTPPF TTKREGNGLG LAEAQKIIRL HGGDIQLKTS DSAVSFFIII
351 PELLAALPKE RAAS*

40

(73) OmcA 9kDa-cysteine-rich lipoprotein(CPn0558)

One example of 'OmcA' protein is disclosed as SEQ ID NO: 9 & 10 in WO 02/02606. {GenBank accession number: gi|4376850|gb|AAD18698.1| 'CPn0558', 'OmcA', 'Omp3'; SEQ ID NO: 73 below and SEQ ID No 10 above}. Preferred
5 OmcA proteins for use with the invention comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 73; and/or
10 (b) which is a fragment of at least *n* consecutive amino acids of SEQ ID NO: 73, wherein *n* is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These OmcA proteins include variants (e.g. allelic variants, homologs, orthologs, paralogs, mutants, etc.) of SEQ ID NO: 73. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 73. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more; preferably 18 or more to remove the signal peptide) from the N-terminus of SEQ ID NO: 73. Other fragments omit one or more domains of the protein (e.g. omission of a signal peptide as described above, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain). The protein may be lipidated (e.g. by a *N*-acyl diglyceride), and may thus have a N-terminal cysteine.

SEQ ID No 73

1 MKKAVLIAAM FCGVVSLSSC CRIVDCCFED PCAPSSCNPC EVIRKKERSC
51 GGNACGSYVP SCSNFCGSTC CNSQSPQVKG CTSPDGRCKQ *

(74) Hypothetical (CPn0331)

One example of a hypothetical protein is set forth as SEQ ID NO: 74 below and SEQ ID No 57 above. Genbank Accession No. GI:4376609; AAD18480.1. Preferred
30 hypothetical proteins for use with the invention comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 74; and/or
35 (b) which is a fragment of at least *n* consecutive amino acids of SEQ ID NO: 74, wherein *n* is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These hypothetical proteins include variants (e.g. allelic variants, homologs, orthologs, paralogs, mutants, etc.) of SEQ ID NO: 74. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 74. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 74. Other fragments omit one or
40 more domains of the protein (e.g. omission of a signal peptide, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain).

SEQ ID NO 74

45 1 mavsggggvg pssdpqkwnp alqgeqaegp splkesifse tkqassaakq eslvrgsgtg
61 myatesgink akyrkaqdrs stspksklkg tfskmravsq gfmsgfgsra srvsakrasd
121 sgegtallpt emdvalkkgn rispemqgff ldsagmggss sdisqlslea lkssafsgar
181 slslssess svasfgsfqk aiepmseekv nawtvarlgg emvsslldpn vetsslvrra
241 matgnegmid lsdlgqeavs tamtsprave gkvkvssds peanptgipn sntleraeke
50 301 aekgesregl sedqmmlara maglltgaap qevlsnsvws gpstvfpppk fsgtlptqrs
361 gdkskhkspg iekstnhtnf splregtvks aevkslphpe smyrfpkdsi vsreepeavv
421 kestafknpe nssqnlflpia vesvfpkesg tggalgdsav sssyhflaqr gvsllaplpr
481 atddykekle ahkgpggppd pliyqyrnva veppivlrsp qpfsqssrls vqgkpeaasv
541 hddggggnsg gfsqdqrrgs sgqkasrqek kgkklstdi

(75) PmpD family (CPn0963)

One example of a PmpD protein is set forth as SEQ ID NO: 75 below Genbank Accession No. GI:4377287; AAD19099.1. Preferred PmpD proteins for use with the invention comprise an amino acid sequence: (a) having 50% or more identity (e.g. 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or more) to SEQ ID NO: 75; and/or (b) which is a fragment of at least *n* consecutive amino acids of SEQ ID NO: 75, wherein *n* is 7 or more (e.g. 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 50, 60, 70, 80, 90, 100, 150, 200, 250 or more). These hypothetical proteins include variants (e.g. allelic variants, homologs, orthologs, paralog, mutants, etc.) of SEQ ID NO: 75. Preferred fragments of (b) comprise an epitope from SEQ ID NO: 75. Other preferred fragments lack one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the C-terminus and/or one or more amino acids (e.g. 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25 or more) from the N-terminus of SEQ ID NO: 75. Other fragments omit one or more domains of the protein (e.g. omission of a signal peptide, of a cytoplasmic domain, of a transmembrane domain, or of an extracellular domain).

SEQ ID No 75

```

20      1 mvakktvrsy rssfshsviv ailsagiafe ahsllhsseld lgvfnkqfee hsahveeaqt
      61 svlkgdvpvn psqkesekvl ytvqpltggs sgesldlada nflehfhqlf eettvfgidq
      121 klwstdldtr nfsqptqepd tsnvavsekis sdtkenrkdl etedpskksk lkevssdlpk
      181 spetavaais edleisenis ardpqglaf fykntssqsi sekdsffggi ifsgsgansg
25      241 lgfenlkapk sgaavysdrd ivfenlvkgl sfiscesled gsaagvniiv thcgdvtltd
      301 catgldleal rlvkdfserg avftarnhev qnnlaggils vvgnkgaivv eknsaeksng
      361 gafacgsfvy snnentalwk enqalsggai ssasdidigg ncsaiefsgn qslialgehi
      421 gltdfvvgga laaaggtltlr nnavvqcvk ntskthggail agtvdlneti sevafkqnta
      481 altggalsan dkviannfg eilfegnevr nhggaiycgc rsnpkleqkd sgeniniign
      541 sgaitflknk asvlevmtga edyagggalw ghnvldsns gniqfignig gstfwigeyv
30      601 gggailstdr vtisnnsdgv vfkgnkgqcl aqkyvapqet apvesdasst nkdekslnac
      661 shgdhyppkt veeevppsll eehpvvsstd irgggailaq hifitdntgn lrfsgnlggg
      721 eesstvgdla ivgggallst nevnvcnqn vvfdsnvtsn gcdsggaila kkvdisanhs
      781 vefvnsngsk fggavcalne svnitdngsa vsfsknrtrl ggagvaapqg svticgnqgn
      841 iafkenfvfg senqrsggga iianssvniq dnagdilfvs nstgsyggai fvgslvaseg
35      901 snprtltitg nsqdilfakn stqtaaslse kdsfgggaiy tqnlkivkna gnvsfygnra
      961 psgagvqiad ggtvcleafg gdilfegnin fdgsfnaihl cgnndskivel savqdkniif
      1021 qdaityeent irglpdkdvs plsapslifn skpqddsahq hegtirfsrg vskipqiaai
      1081 qegtlalsqn aelwlaglkq etgssivlsa gsilrifdsq vdssaplpte nkeetlvsag
40      1141 vqinmsstpt nkdkavdtpv ladiisitvd lssfvpeqdg tlplppeiii pkgtklhsna
      1201 idlkiidptn vgyenhalls shkdiplisl ktaegmtgtp tadaslsnik idvslpsitp
      1261 atyghtgvws eskmedgrlv vgwqptgykl npekggalvl nnlwshytdl ralkqeifah
      1321 htiaqrmeld fstnvwsgsl gvvedcqnig efdgfkhhlt gyalgldtql vedfliggcf
      1381 sqffgktesq sykakndvks ymgaayaqil agpwlikgaf vygninndlt tdygtlgist
45      1441 gswigkgfia gtsidyryiv nprrfisaiv stvvpfveae yvridlpeis eggkevrtfq
      1501 ktrfenvaip fgfalehays rgsraevnsv qlayvfdvyr kgpvslitlk daayswksyg
      1561 vdipckawka rlsnntewns ylstylafny ewredliayd fnggiriif

```

Preferably the composition of the invention comprises a combination of CPn antigens selected from the group consisting of: (1) CPn0301 and CPn0080; (2) CPn 0584 and CPn 0558; and (3) CPn 0331 and CPN 0963. Preferably the composition comprises a combination of any one or more of groups (1), (2) and (3).

Even more preferably, the composition of the present invention comprises a combination of CPn antigens selected from the group consisting of (1) CPn0385, CPn0324, CPn 0503, CPn0525 and CPn 0482. Preferably the composition is administered in the presence of alum and/or cPG.

The invention thus includes a composition comprising a combination of *Chlamydia pneumoniae* antigens, said combination selected from the group consisting of two, three, four, five or six *Chlamydia pneumoniae* antigens of the first antigen group and two, three, four, five, or six *Chlamydia pneumoniae* antigens of the second antigen group. Preferably, the combination is selected from the group consisting of three, four, five or six *Chlamydia pneumoniae* antigens from the first antigen group and three, four, five or six *Chlamydia pneumoniae* antigens from the second antigen group. Still more preferably, the combination consists of six *Chlamydia pneumoniae* antigens from the first antigen group and three, four, five or six, *Chlamydia pneumoniae* antigens from the second antigen group.

The invention further includes a composition comprising a combination of *Chlamydia pneumoniae* antigens, said combination selected from the group consisting of two, three, four, five or six, *Chlamydia pneumoniae* antigens of the second antigen group and two, three, four, five, six, seven or eight *Chlamydia pneumoniae* antigens of the third antigen group. Preferably, the combination is selected from the group consisting of three, four, five or six *Chlamydia pneumoniae* antigens from the second antigen group and three, four, five, six, seven or eight *Chlamydia pneumoniae* from the third antigen group. Still more preferably, the combination consists of six *Chlamydia pneumoniae* antigens from the second antigen group and three, four, five, six, seven or eight *Chlamydia pneumoniae* antigens of the third antigen group.

There is an upper limit to the number of *Chlamydia pneumoniae* antigens which will be in the compositions of the invention. Preferably, the number of *Chlamydia pneumoniae* antigens in a composition of the invention is less than 20, less than 19, less than 18, less than 17, less than 16, less than 15, less than 14, less than 13, less than 12, less than 11, less than 10, less than 9, less than 8, less than 7, less than 6, less than 5, less than 4, or less than 3. Still more preferably, the number of *Chlamydia pneumoniae* antigens in a composition of the invention is less than 6, less than 5, or less than 4. The *Chlamydia pneumoniae* antigens used in the invention are preferably isolated, i.e., separate and discrete, from the whole organism with which the molecule is found in nature or, when the polynucleotide or polypeptide is not found in nature, is sufficiently free of other biological macromolecules so that the polynucleotide or polypeptide can be used for its intended purpose.

In either of the above combinations, preferably the composition comprises one or more *Chlamydia pneumoniae* antigens from the fourth antigen group which includes porB. Or, alternatively, in either of the above combinations, preferably the *Chlamydia pneumoniae* antigens from the fourth antigen group includes one or more members of the pmp3 family.

Other aspects of the present invention are presented in the accompanying claims and in the following description and drawings. These aspects are presented under separate section headings. However, it is to be understood that the teachings under each section are not necessarily limited to that particular section heading.

5

Before describing the present invention in detail, it is to be understood that this invention is not limited to particularly exemplified molecules or process parameters as such may, of course, vary. It is also to be understood that the terminology used herein is for the purpose of describing particular embodiments of the invention only, and is not intended to be limiting. In addition, the practice of the present invention will employ, unless otherwise indicated, conventional methods of virology, microbiology, molecular biology, recombinant DNA techniques and immunology all of which are within the ordinary skill of the art. Such techniques are explained fully in the literature. See, e.g., Sambrook, *et al.*, *Molecular Cloning: A Laboratory Manual* (2nd Edition, 1989); *DNA Cloning: A Practical Approach*, vol. I & II (D. Glover, ed.); *Oligonucleotide Synthesis* (N. Gait, ed., 1984); *A Practical Guide to Molecular Cloning* (1984); and *Fundamental Virology*, 2nd Edition, vol. I & II (B.N. Fields and D.M. Knipe, eds.).

10

15

20

25

All publications, patents and patent applications cited herein, whether *supra* or *infra*, are hereby incorporated by reference in their entirety. It must be noted that, as used in this specification and the appended claims, the singular forms "a", "an" and "the" include plural referents unless the content clearly dictates otherwise. All scientific and technical terms used in this application have meanings commonly used in the art unless otherwise specified. As used in this application, the following words or phrases have the meanings specified.

30

The term "comprising" means "including" as well as "consisting" e.g. a composition "comprising" X may consist exclusively of X or may include something additional e.g. X + Y.

35

40

The term "about" in relation to a numerical value x means, for example, $x \pm 10\%$. References to a percentage sequence identity between two amino acid sequences means that, when aligned, that percentage of amino acids are the same in comparing the two sequences. This alignment and the percent homology or sequence identity can be determined using software programs known in the art, for example those described in section 7.7.18 of *Current Protocols in Molecular Biology* (F.M. Ausubel *et al.*, eds., 1987) Supplement 30. A preferred alignment is determined by the Smith-Waterman homology search algorithm using an affine gap search with a gap open penalty of 12 and a gap extension penalty of 2, BLOSUM matrix of 62. The Smith-Waterman homology search algorithm is disclosed in Smith & Waterman (1981) *Adv. Appl. Math.* 2: 482-489.

IMMUNE RESPONSE

45

The mechanism by which the immune system controls disease includes the induction of neutralising antibodies via humoral immunity and the generation of T-cell responses via cellular immunity. As used herein, the term "immune response" against an antigen refers to the development in a host mammalian subject of a humoral and/or a cellular immune response against that antigen.

As used herein, the term "humoral immune response" refers to an immune response mediated by antibody molecules. The antibodies generated by humoral immunity are primarily effective against extracellular infectious agents.

- 5 SEQ ID Nos 1-76 in the compositions of the invention may be supplemented or substituted with an antibody that binds to the protein. This antibody may be monoclonal or polyclonal.

- 10 As used herein, the term "cell mediated immune (CMI) response" is one mediated by T-lymphocytes and/or other white blood cells. The CMI immune mechanisms are generally more effective against intracellular infections and disease because the CMI mechanisms prime T cells in a way that, when an antigen appears at a later date, memory T cells are activated to result in a CMI response that destroys target cells that have the corresponding antigen or a portion thereof on their cell surfaces, and thereby
- 15 the infecting pathogen. The CMI response is focused on the destruction of the source of infection mediated by either effector cells that destroy infected cells of the host by direct cell-to-cell contact and/or by the release of molecules, such as cytokines, that possess anti-viral activity. Thus the CMI response, which is characterised by a specific T lymphocyte cellular response, is crucial to produce resistance to diseases
- 20 caused by cancer, viruses, pathogenic and other intracellular microorganisms.

T CELLS IMPLICATED IN THE CMI RESPONSE

- At least two special types of T cells are required to initiate and/or to enhance CMI and and humoral responses. The antigenic receptors on a particular subset of T cells
- 25 which express a CD4 co-receptor can be T helper (Th) cells or CD4 T cells (herein after called T helper cells) and they recognise antigenic peptides bound to MHC class II molecules. In contrast, the antigenic receptors on a particular subset of T cells which express a CD8 co-receptor are called Cytotoxic T lymphocytes (CTLs) or CD8+ T cells (hereinafter called CD8+ T cells) and they react with antigens displayed
- 30 on MHC Class I molecules.

HELPER T CELLS

- Helper T cells or CD4+ cells can be further divided into two functionally distinct subsets: Th1 and Th2 which differ in their cytokine and effector function. Th1 and
- 35 Th2 responses have been shown to be regulated not only in a positive but also in a negative way such that Th1 cellular responses are augmented by Th1 cytokines such as IL-2, IL-12 and IFN-gamma and decreased by Th2 cytokines such as IL-4 and IL-10. In contrast, antibody responses are enhanced by Th2 cytokines such as IL-4 and IL-10 but are downregulated by Th1 cytokines such as IFN-gamma and another
- 40 cytokine IL-12 that enhances IFN-gamma and is produced by monocytes. Thus, classic Th1 cytokines such as IFN-gamma, IL-2 and IL-12 can be regarded as immune co-factors that induce an effective inflammatory response. In contrast, the classic Th2 cytokines such as IL-4 and IL-10 can be regarded as cytokines that will suppress a severe inflammatory response.
- 45

CD8+ T CELLS

CD8+ T cells may function in more than one way. The best known function of CD8+ T cells is the killing or lysis of target cells bearing peptide antigen in the context of an MHC class I molecule. Hence the reason why these cells are often termed cytotoxic T lymphocytes (CTL). However, another function, perhaps of greater protective relevance in certain infections is the ability of CD8+ T cells to secrete interferon gamma (IFN-gamma). Thus assays of lytic activity and of IFN-gamma release are both of value in measuring CD8+ T cell immune response (eg in an ELISPOT assay as set forth below). In infectious diseases there is evidence to suggest that CD8+ T cells can protect by killing an infectious agent comprising an infectious antigen at the early stages of a disease before any symptoms of disease are produced.

ENHANCED CMI RESPONSE

The present invention concerns methods, processes and compositions capable of enhancing and/or modulating the CMI response in a host subject against a target antigen. As used herein, the term "enhancing" encompasses improvements in all aspects of the CMI response which include but are not limited to a stimulation and/or augmentation and/or potentiation and/or up-regulation of the magnitude and/or duration, and/or quality of the CMI response to an antigen or a nucleotide sequence encoding an antigen of interest. By way of example, the CMI response may be enhanced by either (i) enhancing the activation and/or production and/or proliferation of CD8+ T cells that recognise a target antigen and/or (ii) shifting the CMI response from a Th2 to a Th1 type response. This enhancement of the Th1 associated responses is of particular value in responding to intracellular infections because, as explained above, the CMI response is enhanced by activated Th1 (such as, for example, IFN-gamma inducing) cells.

Such an enhanced immune response may be generally characterized by increased titers of interferon-producing CD4⁺ and/or CD8⁺ T lymphocytes, increased antigen-specific CD8+ T cell activity, and a T helper 1-like immune response (Th1) against the antigen of interest (characterized by increased antigen-specific antibody titers of the subclasses typically associated with cellular immunity (such as, for example IgG2a), usually with a concomitant reduction of antibody titers of the subclasses typically associated with humoral immunity (such as, for example IgG1)) instead of a T helper 2-like immune response (Th2).

The enhancement of a CMI response may be determined by a number of well-known assays, such as by lymphoproliferation (lymphocyte activation) assays, CD8+ T cell assays, or by assaying for T-lymphocytes specific for the epitope in a sensitized subject (see, for example, Erickson *et al.* (1993) J. Immunol. 151: 4189-4199; and Doe *et al.* (1994) Eur. J. Immunol. 24: 2369-2376) or CD8+ T cell ELISPOT assays for measuring Interferon gamma production (Miyahara *et al* PNAS(USA) (1998) 95: 3954-3959).

ENHANCED T-CELL RESPONSE

- As used herein, the term "enhancing a T-cell response" encompasses improvements in all aspects of the T-cell response which include but are not limited to a stimulation and/or augmentation and/or potentiation and/or up-regulation of the magnitude and/or duration, and/or quality of the T-cell response to an antigen (which may be repeatedly administered) or a nucleotide sequence encoding an antigen. The antigen may be a *Chlamydia* antigen, preferably a *Chlamydia pneumoniae* antigen. By way of example, the T-cell response may be enhanced by either enhancing the activation and/or production and/or distribution and/or proliferation of the induced T-cells and/or longevity of the T-cell response to T-cell inducing/modulating antigen or nucleotide sequence encoding an antigen. The enhancement of the T-cell response in a host subject may be associated with the enhancement and/or modulation of the Th1 immune response in the host subject.
- The enhancement of the T-cell response may be determined by a number of well-known assays, such as by lymphoproliferation (lymphocyte activation) assays, CD8+ T-cell cytotoxic cell assays, or by assaying for T-lymphocytes specific for the epitope in a sensitized subject (see, for example, Erickson *et al.* (1993) *J. Immunol.* 151: 4189-4199; and Doe *et al.* (1994) *Eur. J. Immunol.* 24: 2369-2376) or CD8+ T-cell ELISPOT assays for measuring Interferon gamma production (Miyahara *et al* PNAS(USA) (1998) 95: 3954-3959).

ANTIGEN

- Each disease causing agent or disease state has associated with it an antigen or immunodominant epitope on the antigen which is crucial in immune recognition and ultimate elimination or control of a disease causing agent or disease state in a host. In order to mount a humoral and/or cellular immune response against a particular disease, the host immune system must come in contact with an antigen or an immunodominant epitope on an antigen associated with that disease state.

- As used herein, the term "antigen" refers to any agent, generally a macromolecule, which can elicit an immunological response in an individual. The term "antigen" is used interchangeably with the term "immunogen". The immunological response may be of B- and/or T-lymphocytic cells. The term may be used to refer to an individual macromolecule or to a homogeneous or heterogeneous population of antigenic macromolecules. As used herein, "antigen" is used to refer to a protein molecule or portion thereof which contains one or more antigenic determinants or epitopes. As used herein, the term "antigen" means an immunogenic peptide or protein of interest comprising one or more epitopes capable of inducing a CMI response to an infectious *Chlamydia* pathogen. The antigen can include but is not limited to an auto-antigen, a self-antigen, a cross-reacting antigen, an alloantigen, a tolerogen, an allergen, a hapten, an immunogen or parts thereof as well as any combinations thereof.

EPITOPE

- As used herein, the term "epitope" generally refers to the site on an antigen which is recognised by a T-cell receptor and/or an antibody. Preferably it is a short peptide derived from or as part of a protein antigen. However the term is also intended to include peptides with glycopeptides and carbohydrate epitopes. Several different epitopes may be carried by a single antigenic molecule. The term "epitope" also includes modified sequences of amino acids or carbohydrates which stimulate

responses which recognise the whole organism. It is advantageous if the selected epitope is an epitope of an infectious agent, such as a *Chlamydia* bacterium, which causes the infectious disease.

- 5 SEQ ID Nos 1-76 in the compositions of the invention may be supplemented or substituted with molecules comprising fragments of SEQ ID Nos 1-76. Such fragments may comprise at least n consecutive monomers from the molecules and, depending on the particular sequence, n is either (i) 7 or more for protein molecules (eg. 8, 18, 20 or more), preferably such that the fragment comprises an epitope from
10 the sequence, or (ii) 10 or more for nucleic acid molecules (eg 15, 18, 20, 25, 30, 35, 40 or more).

SOURCE OF EPITOPES

- 15 The epitope can be generated from knowledge the amino acid and corresponding DNA sequences of the peptide or polypeptide, as well as from the nature of particular amino acids (e.g., size, charge, etc.) and the codon dictionary, without undue experimentation. See, e.g., Ivan Roitt, *Essential Immunology*, 1988; Kendrew, *supra*; Janis Kuby, *Immunology*, 1992 e.g., pp. 79-81. Some guidelines in determining whether a protein will stimulate a response, include: Peptide length—preferably the
20 peptide is about 8 or 9 amino acids long to fit into the MHC class I complex and about 13-25 amino acids long to fit into a class II MHC complex. This length is a minimum for the peptide to bind to the MHC complex. It is preferred for the peptides to be longer than these lengths because cells may cut peptides. The peptide may contain an appropriate anchor motif which will enable it to bind to the various class I or class II
25 molecules with high enough specificity to generate an immune response (See Bocchia, M. *et al*, Specific Binding of Leukemia Oncogene Fusion Protein Peptides to HLA Class I Molecules, *Blood* 85:2680-2684; Englehard, VH, Structure of peptides associated with class I and class II MHC molecules *Ann. Rev. Immunol.* 12:181 (1994)). This can be done, without undue experimentation, by comparing the
30 sequence of the protein of interest with published structures of peptides associated with the MHC molecules. Thus, the skilled artisan can ascertain an epitope of interest by comparing the protein sequence with sequences listed in the protein data base.

T CELL EPITOPES

- 35 Preferably one or more antigens of the present invention contain one or more T cell epitopes. As used herein, the term "T cell epitope" refers generally to those features of a peptide structure which are capable of inducing a T cell response. In this regard, it is accepted in the art that T cell epitopes comprise linear peptide determinants that assume extended conformations within the peptide-binding cleft of MHC molecules
40 (Unanue *et al.* (1987) *Science* 236: 551-557). As used herein, a T cell epitope is generally a peptide having at least about 3-5 amino acid residues, and preferably at least 5-10 or more amino acid residues. However, as used herein, the term "T cell epitope" encompasses any MHC Class I-or MHC Class II restricted peptide. The ability of a particular T cell epitope to stimulate/enhance a CMI response may be
45 determined by a number of well-known assays, such as by lymphoproliferation (lymphocyte activation) assays, CD8+ T-cell cytotoxic cell assays, or by assaying for T-lymphocytes specific for the epitope in a sensitized subject. See, e. g., Erickson *et al.* (1993) *J. Immunol.* 151: 4189-4199; and Doe *et al.* (1994) *Eur. J. Immunol.* 24: 2369-2376 or CD8+ T-cell ELISPOT assays for measuring Interferon gamma
50 production (Miyahara *et al* PNAS(USA) (1998) 95: 3954-3959).

CD8+ T-CELL EPITOPES

Preferably the antigens of the present invention comprise CD8+ T-cell inducing epitopes. A CD8+ T-cell -inducing epitope is an epitope capable of stimulating the formation, or increasing the activity, of specific CD8+ T-cells following its administration to a host subject. The CD8+ T-cell epitopes may be provided in a variety of different forms such as a recombinant string of one or two or more epitopes. CD8+ T-cell epitopes have been identified and can be found in the literature, for many different diseases. It is possible to design epitope strings to generate CD8+ T-cell response against any chosen antigen that contains such CD8+ T-cell epitopes. Advantageously, CD8+ T-cell inducing epitopes may be provided in a string of multiple epitopes which are linked together without intervening sequences so that unnecessary nucleic acid material is avoided.

T HELPER EPITOPES

Preferably the antigens of the present invention comprise helper T lymphocyte epitopes. Various methods are available to identify T helper cell epitopes suitable for use in accordance herewith. For example, the amphipathicity of a peptide sequence is known to effect its ability to function as a T helper cell inducer. A full discussion of T helper cell-inducing epitopes is given in U.S. Patent 5,128,319, incorporated herein by reference.

B CELL EPITOPES

Preferably the antigens of the present invention comprise a mixture of CD8+ T-cell epitopes and B cell epitopes. As used herein, the term "B cell epitope" generally refers to the site on an antigen to which a specific antibody molecule binds. The identification of epitopes which are able to elicit an antibody response is readily accomplished using techniques well known in the art. See, e. g., Geysen *et al.* (1984) Proc. Natl. Acad. Sci. USA 81: 3998-4002 (general method of rapidly synthesizing peptides to determine the location of immunogenic epitopes in a given antigen); U. S. Patent No. 4,708,871 (procedures for identifying and chemically synthesizing epitopes of antigens); and Geysen *et al.* (1986) Molecular Immunology 23: 709-715 (technique for identifying peptides with high affinity for a given antibody).

COMBINATION OF EPITOPES

In a preferred embodiment of the present invention, the antigen or antigen combination comprises a mixture of a CD8+ T-cell -inducing epitopes and a T helper cell-inducing epitopes.

As is well known in the art, T and B cell inducing epitopes are frequently distinct from each other and can comprise different peptide sequences. Therefore certain regions of a protein's peptide chain can possess either T cell or B cell epitopes. Therefore, in addition to the CD8+ T-cell epitopes, it may be preferable to include one or more epitopes recognised by T helper cells, to augment the immune response generated by the CD8+ T-cell epitopes.

The mechanism of enhancing a CD8+ T-cell induced response *in vivo* by T helper cell inducing agents is not completely clear. However, without being bound by theory, it is likely that the enhancing agent, by virtue of its ability to induce T helper cells, will result in increased levels of necessary cytokines that assist in the clonal expansion and dissemination of specific CD8+ T-cells. Regardless of the underlying mechanism, it

is envisioned that the use of mixtures of helper T cell and CD8+ T-cell -inducing antigen combinations of the present invention will assist in the enhancement of the CMI response. Particularly suitable T helper cell epitopes are ones which are active in individuals of different HLA types, for example T helper epitopes from tetanus (against which most individuals will already be primed). It may also be useful to include B cell epitopes for stimulating B cell responses and antibody production. Synthetic nucleotide sequences may also be constructed to produce two types of immune responses: T cell only and T cell combined with a B cell response.

10 IMMUNODOMINANT EPITOPE

When an individual is immunized with an antigen or combination of antigens or nucleotide sequence or combinations of nucleotide sequences encoding multiple epitopes of a target antigen, in many instances the majority of responding T lymphocytes will be specific for one or more linear epitopes from that target antigen and/or a majority of the responding B lymphocytes will be specific for one or more linear or conformational epitopes for the antigen or combination of antigens.. For the purposes of the present invention, then, such epitopes are referred to as "immunodominant epitopes". In an antigen having several immunodominant epitopes, a single epitope may be the most dominant in terms of commanding a specific T or B cell response.

As the Examples show, at least sixteen peptides of the present invention were recognised by IFN-gamma positive CD8+ T cell populations which were actually expanded as a result of bacterial infection.

25 ADJUVANTS

The compositions of the present invention may be administered in conjunction with other immunoregulatory agents. In particular, the compositions of the present invention may be administered with an adjuvant.

The inclusion of an adjuvant and in particular, a genetic adjuvant may be useful in further enhancing or modulating the CMI response. An adjuvant may enhance the CMI response by enhancing the immunogenicity of a co-administered antigen in an immunized subject, as well inducing a Th1-like immune response against the co-administered antigen which is beneficial in a vaccine product.

An immune response and particularly a CMI response may be refined, by the addition of adjuvants to combinations of antigens or nucleotide sequences encoding combinations of antigens which lead to particularly effective compositions for eliciting a long lived and sustained enhanced CMI response.

As used herein, the term "adjuvant" refers to any material or composition capable of specifically or non-specifically altering, enhancing, directing, redirecting, potentiating or initiating an antigen-specific immune response.

The term "adjuvant" includes but is not limited to a bacterial ADP-ribosylating exotoxin, a biologically active factor, immunomodulatory molecule, biological response modifier or immunostimulatory molecule such as a cytokine, an interleukin, a chemokine or a ligand or an epitope (such as a helper T cell epitope) and optimally combinations thereof which, when administered with an antigen, antigen composition

or nucleotide sequence encoding such antigens enhances or potentiates or modulates the CMI response relative to the CMI response generated upon administration of the antigen or combination of antigens alone. The adjuvant may be any adjuvant known in the art which is appropriate for human or animal use.

5

Immunomodulatory molecules such as cytokines (TNF-alpha, IL-6, GM-CSF, and IL-2), and co-stimulatory and accessory molecules (B7-1, B7-2) may be used as adjuvants in a variety of combinations. In one embodiment GM-CSF is not administered to subject before, in or after the administration regimen. Simultaneous production of an immunomodulatory molecule and an antigen of interest at the site of expression of the antigen of interest may enhance the generation of specific effectors which may help to enhance the CMI response. The degree of enhancement of the CMI response may be dependent upon the specific immunostimulatory molecules and/or adjuvants used because different immunostimulatory molecules may elicit different mechanisms for enhancing and/or modulating the CMI response. By way of example, the different effector mechanisms/immunomodulatory molecules include but are not limited to augmentation of help signal (IL-2), recruitment of professional APC (GM-CSF), increase in T cell frequency (IL-2), effect on antigen processing pathway and MHC expression (IFN-gamma and TNF-alpha) and diversion of immune response away from the Th1 response and towards a Th2 response (LTB) (see WO 97/02045). Unmethylated CpG containing oligonucleotides (see WO96/02555) are also preferential inducers of a Th1 response and are suitable for use in the present invention.

25 Without being bound by theory, the inclusion of an adjuvant is advantageous because the adjuvant may help to enhance the CMI response to the expressed antigen by diverting the Th2 response to a Th1 response and/or specific effector associated mechanisms to an expressed epitope with the consequent generation and maintenance of an enhanced CMI response (see, for example, the teachings in WO 97/02045).

30

The inclusion of an adjuvant with an antigen or nucleotide sequence encoding the antigen is also advantageous because it may result in a lower dose or fewer doses of the antigen/antigenic combination being necessary to achieve the desired CMI response in the subject to which the antigen or nucleotide sequence encoding the antigen is administered, or it may result in a qualitatively and/or quantitatively different immune response in the subject. The effectiveness of an adjuvant can be determined by administering the adjuvant with the antigen in parallel with the antigen alone to animals and comparing antibody and/or cellular-mediated immunity in the two groups using standard assays such as radioimmunoassay, ELISAs, CD8+ T-cell assays, and the like, all well known in the art. Typically, the adjuvant is a separate moiety from the antigen, although a single molecule (such for example, CTB) can have both adjuvant and antigen properties.

35

40

As used herein, the term "genetic adjuvant" refers to an adjuvant encoded by a nucleotide sequence and which, when administered with the antigen enhances the CMI response relative to the CMI response generated upon administration of the antigen alone.

45

Bacterial ADP-ribosylating toxins and detoxified derivatives thereof may be used as adjuvants in the invention. Preferably, the protein is derived from *E. coli* (i.e., *E. coli* heat labile enterotoxin "LT"), cholera ("CT"), or pertussis ("PT").

5 In one preferred embodiment, the genetic adjuvant is a bacterial ADP-ribosylating exotoxin.

10 ADP-ribosylating bacterial toxins are a family of related bacterial exotoxins and include diphtheria toxin (DT), pertussis toxin (PT), cholera toxin (CT), the *E. coli* heat-labile toxins (LT1 and LT2), *Pseudomonas* endotoxin A, *Pseudomonas* exotoxin S, *B. cereus* exoenzyme, *B. sphaericus* toxin, *C. botulinum* C2 and C3 toxins, *C. limosum* exoenzyme, as well as toxins from *C. perfringens*, *C. spiriforma* and *C. difficile*, *Staphylococcus aureus* EDIN, and ADP-ribosylating bacterial toxin mutants such as CRM₁₉₇, a non-toxic diphtheria toxin mutant (see, e.g., Bixler *et al.* (1989) *Adv. Exp. Med. Biol.* 251:175; and Constantino *et al.* (1992) *Vaccine*). Most ADP-ribosylating bacterial toxins are organized as an A:B multimer, wherein the A subunit contains the ADP-ribosyltransferase activity, and the B subunit acts as the binding moiety. Preferred ADP-ribosylating bacterial toxins for use in the compositions of the present invention include cholera toxin and the *E. coli* heat-labile toxins.

20 Cholera toxin (CT) and the related *E. coli* heat labile enterotoxins (LT) are secretion products of their respective enterotoxic bacterial strains that are potent immunogens and exhibit strong toxicity when administered systemically, orally, or mucosally. Both CT and LT are known to provide adjuvant effects for antigen when administered via the intramuscular or oral routes. These adjuvant effects have been observed at
25 doses below that required for toxicity. The two toxins are extremely similar molecules, and are at least about 70-80% homologous at the amino acid level.

30 Preferably the genetic adjuvant is cholera toxin (CT), enterotoxigenic *E. Coli* heat-labile toxin (LT), or a derivative, subunit, or fragment of CT or LT which retains adjuvant activity. In an even more preferred embodiment, the genetic adjuvant is LT. In another preferred embodiment, the genetic adjuvant may be CTB or LTB.

Preferably the enterotoxin is a non-toxic enterotoxin.

35 The use of detoxified ADP-ribosylating toxins as mucosal adjuvants is described in WO 95/17211 and as parenteral adjuvants in WO 98/42375. The toxin or toxoid is preferably in the form of a holotoxin, comprising both A and B subunits. Preferably, the A subunit contains a detoxifying mutation; preferably the B subunit is not mutated. Preferably, the adjuvant is a detoxified LT mutant such as LT-K63, LT-R72, and LTR192G. The use of ADP-ribosylating toxins and detoxified derivatives thereof,
40 particularly LT-K63 and LT-R72, as adjuvants can be found in the following references each of which is specifically incorporated by reference herein in their entirety (Beignon, *et al.* *Infection and Immunity* (2002) 70(6):3012 - 3019; Pizza, *et al.*, *Vaccine* (2001) 19:2534 - 2541; Pizza, *et al.*, *Int. J. Med. Microbiol* (2000) 290(4-5):455-461; Scharton-Kersten *et al.* *Infection and Immunity* (2000) 68(9):5306 -
45 5313; Ryan *et al.* *Infection and Immunity* (1999) 67(12):6270 - 6280; Partidos *et al.* *Immunol. Lett.* (1999) 67(3):209 - 216; Peppoloni *et al.* *Vaccines* (2003) 2(2):285 - 293; and Pine *et al.* *J. Control Release* (2002) 85(1-3):263 - 270). Numerical reference for amino acid substitutions is preferably based on the alignments of the A and B subunits of ADP-ribosylating toxins set forth in Domenighini *et al.*, *Mol.*

Microbiol (1995) 15(6):1165 – 1167, specifically incorporated herein by reference in its entirety.

5 By way of further example, at least one of the enterotoxin subunit coding regions may be genetically modified to detoxify the subunit peptide encoded thereby, for example wherein the truncated A subunit coding region has been genetically modified to disrupt or inactivate ADP-ribosyl transferase activity in the subunit peptide expression product (see, for example, WO 03/004055).

10 Thus, these results demonstrate that this genetic adjuvant is particularly desirable where an even more enhanced CMI response is desired. Other desirable genetic adjuvants include but are not limited to nucleotide sequences encoding IL-10, IL-12, IL-13, the interferons (IFNs) (for example, IFN-alpha, IFN-ss, and IFN-gamma), and preferred combinations thereof. Still other such biologically active factors that
15 enhance the CMI response may be readily selected by one of skill in the art, and a suitable plasmid vector containing same constructed by known techniques.

Preferred further adjuvants include, but are not limited to, one or more of the following set forth below:

20

Mineral Containing Compositions

Mineral containing compositions suitable for use as adjuvants in the invention include mineral salts, such as aluminium salts and calcium salts. The invention includes mineral salts such as hydroxides (e.g. oxyhydroxides), phosphates (e.g.
25 hydroxyphosphates, orthophosphates), sulphates, etc. {e.g. see chapters 8 & 9 of ref. Bush and Everett (2001) Int J Syst Evol Microbiol 51: 203-220}, or mixtures of different mineral compounds, with the compounds taking any suitable form (e.g. gel, crystalline, amorphous, etc.), and with adsorption being preferred. The mineral containing compositions may also be formulated as a particle of metal salt. See WO
30 00/23105.

Oil-Emulsions

Oil-emulsion compositions suitable for use as adjuvants in the invention include squalene-water emulsions, such as MF59 (5% Squalene, 0.5% Tween 80, and 0.5%
35 Span 85, formulated into submicron particles using a microfluidizer). See WO 90/14837. Complete Freund's adjuvant (CFA) and incomplete Freund's adjuvant (IFA) may also be used as adjuvants in the invention.

Saponin Formulations

40 Saponin formulations, may also be used as adjuvants in the invention. Saponins are a heterologous group of sterol glycosides and triterpenoid glycosides that are found in the bark, leaves, stems, roots and even flowers of a wide range of plant species. Saponin from the bark of the *Quillaia saponaria* Molina tree have been widely studied as adjuvants. Saponin can also be commercially obtained from *Smilax ornata*
45 (sarsapilla), *Gypsophilla paniculata* (brides veil), and *Saponaria officianalis* (soap root). Saponin adjuvant formulations include purified formulations, such as QS21, as well as lipid formulations, such as ISCOMs. Saponin compositions have been purified using High Performance Thin Layer Chromatography (HP-LC) and Reversed Phase High Performance Liquid Chromatography (RP-HPLC). Specific purified
50 fractions using these techniques have been identified, including QS7, QS17, QS18,

QS21, QH-A, QH-B and QH-C. Preferably, the saponin is QS21. A method of production of QS21 is disclosed in U.S. Patent No. 5,057,540. Saponin formulations may also comprise a sterol, such as cholesterol (see WO 96/33739).

5 Combinations of saponins and cholesterol can be used to form unique particles called Immunostimulating Complexs (ISCOMs). ISCOMs typically also include a phospholipid such as phosphatidylethanolamine or phosphatidylcholine. Any known saponin can be used in ISCOMs. Preferably, the ISCOM includes one or more of Quil A, QHA and QHC. ISCOMs are further described in EP 0 109 942, WO 10 96/11711 and WO 96/33739. Optionally, the ISCOMS may be devoid of additional detergent. See WO 00/07621.

A review of the development of saponin based adjuvants can be found in Barr *et al* (1998) Advanced Drug Delivery Reviews 32: 247-271 and Sjolander *et al* (1998) Advanced Drug Delivery Reviews (1998) 32: 321-338.

15

Virosomes and Virus Like Particles (VLPs)

Virosomes and Virus Like Particles (VLPs) can also be used as adjuvants in the invention. These structures generally contain one or more proteins from a virus optionally combined or formulated with a phospholipid. They are generally non-pathogenic, non-replicating and generally do not contain any of the native viral genome. The viral proteins may be recombinantly produced or isolated from whole viruses. These viral proteins suitable for use in virosomes or VLPs include proteins derived from influenza virus (such as HA or NA), Hepatitis B virus (such as core or capsid proteins), Hepatitis E virus, measles virus, Sindbis virus, Rotavirus, Foot-and-Mouth Disease virus, Retrovirus, Norwalk virus, human Papilloma virus, HIV, RNA-phages, Q β -phage (such as coat proteins), GA-phage, fr-phage, AP205 phage, and Ty (such as retrotransposon Ty protein p1). VLPs are discussed further in WO 25 03/024480, WO 03/024481; Niikura *et al* Virology (2002) 293:273 - 280; Lenz *et al* Journal of Immunology (2001) 5246 - 5355; Pinto, *et al* Journal of Infectious Diseases (2003) 188:327 - 338; and Gerber *et al* Journal of Virology (2001) 30 75(10):4752 - 4760; Virosomes are discussed further in, for example, Gluck *et al* Vaccine (2002) 20:B10 -B16.

Bacterial or Microbial Derivatives

35 Adjuvants suitable for use in the invention include bacterial or microbial derivatives such as:

Non-toxic derivatives of enterobacterial lipopolysaccharide (LPS)

Such derivatives include Monophosphoryl lipid A (MPL) and 3-O-deacylated MPL 40 (3dMPL). 3dMPL is a mixture of 3 De-O-acylated monophosphoryl lipid A with 4, 5 or 6 acylated chains. A preferred "small particle" form of 3 De-O-acylated monophosphoryl lipid A is disclosed in EP 0 689 454. Such "small particles" of 3dMPL are small enough to be sterile filtered through a 0.22 micron membrane (see EP 0 689 454). Other non-toxic LPS derivatives include monophosphoryl lipid A 45 mimics, such as aminoalkyl glucosaminide phosphate derivatives *e.g.* RC-529. See Johnson *et al.* (1999) *Bioorg Med Chem Lett* 9:2273-2278.

Lipid A Derivatives

Lipid A derivatives include derivatives of lipid A from *Escherichia coli* such as OM-174. OM-174 is described for example in Meraldi *et al.* Vaccine (2003) 21:2485 – 2491; Pajak, *et al* Vaccine (2003) 21:836 – 842.

Immunostimulatory oligonucleotides

Immunostimulatory oligonucleotides suitable for use as adjuvants in the invention include nucleotide sequences containing a CpG motif (a sequence containing an unmethylated cytosine followed by guanosine and linked by a phosphate bond). Bacterial double stranded RNA or oligonucleotides containing palindromic or poly(dG) sequences have also been shown to be immunostimulatory.

The CpG's can include nucleotide modifications/analogues such as phosphorothioate modifications and can be double-stranded or single-stranded. Optionally, the guanosine may be replaced with an analog such as 2'-deoxy-7-deazaguanosine. See Kandimalla, *et al* Nucleic Acids Research (2003) 31(9): 2393 – 2400; WO 02/26757 and WO 99/62923 for examples of possible analog substitutions. The adjuvant effect of CpG oligonucleotides is further discussed in Krieg Nature Medicine (2003) 9(7): 831 – 835; McCluskie, *et al* FEMS Immunology and Medical Microbiology (2002) 32:179 – 185; WO 98/40100, U.S. Patent No. 6,207,646, U.S. Patent No. 6,239,116, and U.S. Patent No. 6,429,199.

The CpG sequence may be directed to TLR9, such as the motif GTCGTT or TTCGTT. See Kalman *et al* (1999) (Nature Genetics 21: 385-389). The CpG sequence may be specific for inducing a Th1 immune response, such as a CpG-A ODN, or it may be more specific for inducing a B cell response, such as a CpG-B ODN. CpG-A and CpG-B ODNs are discussed in Blackwell, *et al* J. Immunol. (2003) 170(8):4061 – 4068; Krieg BBRC (2003) 306:948 – 953; and WO 01/95935. Preferably, the CpG is a CpG-A ODN.

Preferably, the CpG oligonucleotide is constructed so that the 5' end is accessible for receptor recognition. Optionally, two CpG oligonucleotide sequences may be attached at their 3' ends to form "immunomers". See, for example, Kandimalla, *et al* (2003) 31(part 3):664 – 658; Bhagat *et al* BBRC (2003) 300:853 – 861 and WO 03/035836.

Human Immunomodulators

Human immunomodulators suitable for use as adjuvants in the invention include cytokines, such as interleukins (*e.g.* IL-1, IL-2, IL-4, IL-5, IL-6, IL-7, IL-12, *etc.*), interferons (*e.g.* interferon- γ), macrophage colony stimulating factor, and tumor necrosis factor.

Bioadhesives and Mucoadhesives

Bioadhesives and mucoadhesives may also be used as adjuvants in the invention. Suitable bioadhesives include esterified hyaluronic acid microspheres (Singh *et al.* (2001) *J. Cont. Rel.* 70:267-276) or mucoadhesives such as cross-linked derivatives of poly(acrylic acid), polyvinyl alcohol, polyvinyl pyrrolidone, polysaccharides and carboxymethylcellulose. Chitosan and derivatives thereof may also be used as adjuvants in the invention. See for example, WO99/27960.

Microparticles

Microparticles may also be used as adjuvants in the invention. Microparticles (*i.e.* a particle of ~100nm to ~150 μ m in diameter, more preferably ~200nm to ~30 μ m in diameter, and most preferably ~500nm to ~10 μ m in diameter) formed from materials that are biodegradable and non-toxic (*e.g.* a poly(α -hydroxy acid), a polyhydroxybutyric acid, a polyorthoester, a polyanhydride, a polycaprolactone, *etc.*), with poly(lactide-co-glycolide) are preferred, optionally treated to have a negatively-charged surface (*e.g.* with SDS) or a positively-charged surface (*e.g.* with a cationic detergent, such as CTAB).

Liposomes

Examples of liposome formulations suitable for use as adjuvants are described in U.S. Patent No. 6,090,406, U.S. Patent No. 5,916,588, and EP 0 626 169.

Polyoxyethylene ether and Polyoxyethylene Ester Formulations

Adjuvants suitable for use in the invention include polyoxyethylene ethers and polyoxyethylene esters (WO99/52549). Such formulations further include polyoxyethylene sorbitan ester surfactants in combination with an octoxynol (WO01/21207) as well as polyoxyethylene alkyl ethers or ester surfactants in combination with at least one additional non-ionic surfactant such as an octoxynol (WO01/21152). Preferred polyoxyethylene ethers are selected from the following group: polyoxyethylene-9-lauryl ether (laureth 9), polyoxyethylene-9-stearyl ether, polyoxyethylene-8-stearyl ether, polyoxyethylene-4-lauryl ether, polyoxyethylene-35-lauryl ether, and polyoxyethylene-23-lauryl ether.

Polyphosphazene (PCPP)

PCPP formulations are described, for example, in Andrianov *et al* Biomaterials (1998) 19(1 - 3):109 - 115; Payne *et al* Adv. Drug. Delivery Review (1998) 31(3):185 - 196.

Muramyl peptides

Examples of muramyl peptides suitable for use as adjuvants in the invention include N-acetyl-muramyl-L-threonyl-D-isoglutamine (thr-MDP), N-acetyl-normuramyl-L-alanyl-D-isoglutamine (nor-MDP), and N-acetylmuramyl-L-alanyl-D-isoglutaminyl-L-alanine-2-(1'-2'-dipalmitoyl-*sn*-glycero-3-hydroxyphosphoryloxy)-ethylamine MTP-PE).

Imidazoquinolone Compounds

Examples of imidazoquinolone compounds suitable for use as adjuvants in the invention include Imiquimod and its homologues, described further in Stanley, "Imiquimod and the imidazoquinolones: mechanism of action and therapeutic potential" Clin Exp Dermatol (2002) 27(7):571 - 577; and Jones, "Resiquimod 3M", Curr Opin Investig Drugs (2003) 4(2):214 - 218. The invention may also comprise combinations of aspects of one or more of the adjuvants identified above. For example, the following adjuvant compositions may be used in the invention:

- (1) a saponin and an oil-in-water emulsion (WO99/11241);
- (2) a saponin (*e.g.*, QS21) + a non-toxic LPS derivative (*e.g.*, 3dMPL) (see WO 94/00153);
- (3) a saponin (*e.g.*, QS21) + a non-toxic LPS derivative (*e.g.*, 3dMPL) + a cholesterol;

(4) a saponin (e.g. QS21) + 3dMPL + IL-12 (optionally + a sterol (WO98/57659); combinations of 3dMPL with, for example, QS21 and/or oil-in-water emulsions (European patent applications 0835318, 0735898 and 0761231).

5 (5) SAF, containing 10% Squalane, 0.4% Tween 80, 5% pluronic-block polymer L121, and thr-MDP, either microfluidized into a submicron emulsion or vortexed to generate a larger particle size emulsion.

(6) RibiTM adjuvant system (RAS), (Ribi Immunochem) containing 2% Squalene, 0.2% Tween 80, and one or more bacterial cell wall components from the group consisting of monophosphorylipid A (MPL), trehalose dimycolate (TDM), and cell wall skeleton (CWS), preferably MPL + CWS (DetoxTM); and

10 (7) one or more mineral salts (such as an aluminum salt) + a non-toxic derivative of LPS (such as 3dPML).

15 Aluminium salts and MF59 are preferred adjuvants for parenteral immunisation. Mutant bacterial toxins are preferred mucosal adjuvants. The composition may include an antibiotic.

20 Preferably the compositions of the present invention are administered with alum and/or CpG sequences.

Nucleic Acid

25 The antigens or epitopes of the present invention may be administered as nucleotide sequences encoding the antigens or epitopes. As used herein, the term nucleotide sequence refers to one of more nucleotide sequences which encode one or more epitopes which are used in the compositions or combinations of the present invention. The term "nucleotide sequence (NOI)" is synonymous with the term "polynucleotide" or "nucleic acid". The NOI may be DNA or RNA of genomic or synthetic or of recombinant origin. The NOI may be double-stranded or single-stranded whether representing the sense or antisense strand or combinations thereof. For some applications, preferably, the NOI is DNA. For some applications, preferably, the NOI is prepared by use of recombinant DNA techniques (e.g. recombinant DNA). For some applications, preferably, the NOI is cDNA. For some applications, preferably, the NOI may be the same as the naturally occurring form.

35 The term "nucleic acid" includes DNA and RNA, and also their analogues, such as those containing modified backbones (e.g. phosphorothioates, etc.), and also peptide nucleic acids (PNA), etc. The invention includes nucleic acid comprising sequences complementary to those described above (e.g. for antisense or probing purposes).

40 Nucleic acid according to the invention can be prepared in many ways (e.g. by chemical synthesis, from genomic or cDNA libraries, from the organism itself, etc.) and can take various forms (e.g. single stranded, double stranded, vectors, probes, etc.). They are preferably prepared in substantially pure form (i.e. substantially free from other *Chlamydia* or host cell nucleic acids).

45 The invention provides a process for producing nucleic acid of the invention, comprising the step of amplifying nucleic acid using a primer-based amplification method (e.g. PCR).

The invention provides a process for producing nucleic acid of the invention, comprising the step of synthesising at least part of the nucleic acid by chemical means.

5 VECTOR

In one embodiment of the present invention, an antigen or antigenic combination or NOI encoding same is administered directly to a host subject. In another embodiment of the present invention, a vector comprising an NOI is administered to a host subject. Preferably the NOI is prepared and/or administered using a genetic vector. As it is well known in the art, a vector is a tool that allows or facilitates the transfer of an entity from one environment to another. In accordance with the present invention, and by way of example, some vectors used in recombinant DNA techniques allow entities, such as a segment of DNA (such as a heterologous DNA segment, such as a heterologous cDNA segment), to be transferred into a host and/or a target cell for the purpose of replicating the vectors comprising the NOI of the present invention and/or expressing the antigens or epitopes of the present invention encoded by the NOI. Examples of vectors used in recombinant DNA techniques include but are not limited to plasmids, chromosomes, artificial chromosomes or viruses. The term "vector" includes expression vectors and/or transformation vectors. The term "expression vector" means a construct capable of *in vivo* or *in vitro/ex vivo* expression. The term "transformation vector" means a construct capable of being transferred from one species to another.

NAKED DNA

25 The vectors comprising the NOI of the present invention may be administered directly as "a naked nucleic acid construct", preferably further comprising flanking sequences homologous to the host cell genome. As used herein, the term "naked DNA" refers to a plasmid comprising the NOI of the present invention together with a short promoter region to control its production. It is called "naked" DNA because the plasmids are not carried in any delivery vehicle. When such a DNA plasmid enters a host cell, such as a eukaryotic cell, the proteins it encodes are transcribed and translated within the cell.

VIRAL VECTORS

35 Alternatively, the vectors comprising the NOI of the present invention may be introduced into suitable host cells using a variety of viral techniques which are known in the art, such as for example infection with recombinant viral vectors such as retroviruses, herpes simplex viruses and adenoviruses. The vector may be a recombinant viral vectors. Suitable recombinant viral vectors include but are not limited to adenovirus vectors, adeno-associated viral (AAV) vectors, herpes-virus vectors, a retroviral vector, lentiviral vectors, baculoviral vectors, pox viral vectors or parvovirus vectors (see Kestler *et al* 1999 Human Gene Ther 10(10):1619-32). In the case of viral vectors, administration of the NOI is mediated by viral infection of a target cell.

45 TARGETED VECTOR

The term "targeted vector" refers to a vector whose ability to infect or transfect or transduce a cell or to be expressed in a host and/or target cell is restricted to certain cell types within the host subject, usually cells having a common or similar phenotype.

50

EXPRESSION VECTOR

Preferably, the NOI of the present invention which is inserted into a vector is operably linked to a control sequence that is capable of providing for the expression of the antigens or epitopes by the host cell, i.e. the vector is an expression vector. The agent
5 produced by a host cell may be secreted or may be contained intracellularly depending on the NOI and/or the vector used. As will be understood by those of skill in the art, expression vectors containing the NOI can be designed with signal sequences which direct secretion of the EOI through a particular prokaryotic or eukaryotic cell membrane.

10 FUSION PROTEINS

The *Chlamydia pneumoniae* antigens used in the invention may be present in the composition as individual separate polypeptides, but it is preferred that at least two (i.e. 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20) of the antigens are
15 expressed as a single polypeptide chain (a 'hybrid' polypeptide). Hybrid polypeptides offer two principal advantages: first, a polypeptide that may be unstable or poorly expressed on its own can be assisted by adding a suitable hybrid partner that overcomes the problem; second, commercial manufacture is simplified as only one expression and purification need be employed in order to produce two polypeptides which are both antigenically useful.

20 The hybrid polypeptide may comprise two or more polypeptide sequences from the first antigen group. Accordingly, the invention includes a composition comprising a first amino acid sequence and a second amino acid sequence, wherein said first and second amino acid sequences are selected from a *Chlamydia* bacterium, preferably a
25 *Chlamydia pneumoniae* antigen or a fragment thereof of the first antigen group. Preferably, the first and second amino acid sequences in the hybrid polypeptide comprise different epitopes.

30 The hybrid polypeptide may comprise two or more polypeptide sequences from the second antigen group. Accordingly, the invention includes a composition comprising a first amino acid sequence and a second amino acid sequence, wherein said first and second amino acid sequences are selected from a *Chlamydia pneumoniae* antigen or a fragment thereof of the second antigen group. Preferably, the first and second amino acid sequences in the hybrid polypeptide comprise difference epitopes.
35

The hybrid polypeptide may comprise one or more polypeptide sequences from the first antigen group and one or more polypeptide sequences from the second antigen group. Accordingly, the invention includes a composition comprising a first amino acid sequence and a second amino acid sequence, said first amino acid sequence selected
40 from a *Chlamydia pneumoniae* antigen or a fragment thereof from the first antigen group and said second amino acid sequence selected from a *Chlamydia* bacterium, preferably a *Chlamydia pneumoniae* antigen or a fragment thereof from the second antigen group. Preferably, the first and second amino acid sequences in the hybrid polypeptide comprise difference epitopes.
45

The hybrid polypeptide may comprise one or more polypeptide sequences from the first antigen group and one or more polypeptide sequences from the third antigen group or the fourth antigen group or the fifth antigen group or the sixth antigen group or the seventh antigen group or the eighth antigen group or the ninth antigen group or
50 the tenth antigen group. Accordingly, the invention includes a composition

- comprising a first amino acid sequence and a second amino acid sequence, said first amino acid sequence selected from a *Chlamydia pneumoniae* antigen or a fragment thereof from the first antigen group and said second amino acid sequence selected from a *Chlamydia pneumoniae* antigen or a fragment thereof from the third antigen group or the fourth antigen group or the fifth antigen group or the sixth antigen group or the seventh antigen group or the eighth antigen group or the ninth antigen group or the tenth antigen group. Preferably, the first and second amino acid sequences in the hybrid polypeptide comprise difference epitopes.
- 10 The hybrid polypeptide may comprise one or more polypeptide sequences from the second antigen group and one or more polypeptide sequences from the third antigen group or the fourth antigen group or the fifth antigen group or the sixth antigen group or the seventh antigen group or the eighth antigen group or the ninth antigen group or the tenth antigen group. Accordingly, the invention includes a composition
- 15 comprising a first amino acid sequence and a second amino acid sequence, said first amino acid sequence selected from a *Chlamydia pneumoniae* antigen or a fragment thereof from the second antigen group and said second amino acid sequence selected from a *Chlamydia pneumoniae* antigen or a fragment thereof from the third antigen group or the fourth antigen group or the fifth antigen group or the sixth antigen group or the seventh antigen group or the eighth antigen group or the ninth antigen group or the tenth antigen group. Preferably, the first and second amino acid sequences in the hybrid polypeptide comprise difference epitopes.
- 20 Hybrids consisting of amino acid sequences from two, three, four, five, six, seven, eight, nine, or ten *Chlamydia pneumoniae* antigens are preferred. In particular, hybrids consisting of amino acid sequences from two, three, four, or five *Chlamydia pneumoniae* antigens are preferred. Different hybrid polypeptides may be mixed together in a single formulation. Within such combinations, a *Chlamydia pneumoniae* antigen may be present in more than one hybrid polypeptide and/or as a non-hybrid polypeptide. It is preferred, however, that an antigen is present either as a hybrid or as a non-hybrid, but not as both.
- 25 Two-antigen hybrids for use in the invention may comprise any one of the combinations disclosed above.
- 30 Hybrid polypeptides can be represented by the formula $\text{NH}_2\text{-A-}\{-\text{X-L-}\}_n\text{-B-COOH}$, wherein: X is an amino acid sequence of a *Chlamydia pneumoniae* antigen or a fragment thereof from the first antigen group, the second antigen group or the third antigen group or the fourth antigen group or the fifth antigen group or the sixth antigen group or the seventh antigen group or the eighth antigen group or the ninth antigen group or the tenth antigen group.; L is an optional linker amino acid sequence; A is an optional N-terminal amino acid sequence; B is an optional C-terminal amino acid sequence; and n is 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14 or 15.
- 35 If a -X- moiety has a leader peptide sequence in its wild-type form, this may be included or omitted in the hybrid protein. In some embodiments, the leader peptides will be deleted except for that of the -X- moiety located at the N-terminus of the hybrid protein *i.e.* the leader peptide of X_1 will be retained, but the leader peptides of $\text{X}_2 \dots \text{X}_n$ will be omitted. This is equivalent to deleting all leader peptides and using the leader peptide of X_1 as moiety -A-.
- 40
- 45
- 50

For each n instances of $\{-X-L-\}$, linker amino acid sequence $-L-$ may be present or absent. For instance, when $n=2$ the hybrid may be $NH_2-X_1-L_1-X_2-L_2-COOH$, $NH_2-X_1-X_2-COOH$, $NH_2-X_1-L_1-X_2-COOH$, $NH_2-X_1-X_2-L_2-COOH$, *etc.* Linker amino acid sequence(s) $-L-$ will typically be short (*e.g.* 20 or fewer amino acids *i.e.* 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, 1). Examples comprise short peptide sequences which facilitate cloning, poly-glycine linkers (*i.e.* comprising Gly_n where $n = 2, 3, 4, 5, 6, 7, 8, 9, 10$ or more), and histidine tags (*i.e.* His_n where $n = 3, 4, 5, 6, 7, 8, 9, 10$ or more). Other suitable linker amino acid sequences will be apparent to those skilled in the art. A useful linker is GSGGGG (SEQ ID No 77), with the Gly-Ser dipeptide being formed from a *Bam*HI restriction site, thus aiding cloning and manipulation, and the $(Gly)_4$ tetrapeptide being a typical poly-glycine linker.

$-A-$ is an optional N-terminal amino acid sequence. This will typically be short (*e.g.* 40 or fewer amino acids *i.e.* 39, 38, 37, 36, 35, 34, 33, 32, 31, 30, 29, 28, 27, 26, 25, 24, 23, 22, 21, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, 1). Examples include leader sequences to direct protein trafficking, or short peptide sequences which facilitate cloning or purification (*e.g.* histidine tags *i.e.* His_n where $n = 3, 4, 5, 6, 7, 8, 9, 10$ or more). Other suitable N-terminal amino acid sequences will be apparent to those skilled in the art. If X_1 lacks its own N-terminus methionine, $-A-$ is preferably an oligopeptide (*e.g.* with 1, 2, 3, 4, 5, 6, 7 or 8 amino acids) which provides a N-terminus methionine.

$-B-$ is an optional C-terminal amino acid sequence. This will typically be short (*e.g.* 40 or fewer amino acids *i.e.* 39, 38, 37, 36, 35, 34, 33, 32, 31, 30, 29, 28, 27, 26, 25, 24, 23, 22, 21, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, 2, 1). Examples include sequences to direct protein trafficking, short peptide sequences which facilitate cloning or purification (*e.g.* comprising histidine tags *i.e.* His_n where $n = 3, 4, 5, 6, 7, 8, 9, 10$ or more), or sequences which enhance protein stability. Other suitable C-terminal amino acid sequences will be apparent to those skilled in the art. Most preferably, n is 2 or 3.

The invention also provides nucleic acid encoding hybrid polypeptides of the invention. Furthermore, the invention provides nucleic acid which can hybridise to this nucleic acid, preferably under "high stringency" conditions (*e.g.* 65°C in a 0.1xSSC, 0.5% SDS solution).

The NOI of the present invention may be expressed as a fusion protein comprising an adjuvant and/or a biological response modifier and/or immunomodulator fused to the antigens or epitopes of the present invention to further enhance and/or augment the CMI response obtained. The biological response modifier may act as an adjuvant in the sense of providing a generalised stimulation of the CMI response. The antigens or epitopes may be attached to either the amino or carboxy terminus of the biological response modifier.

METHODS OF MAKING

Polypeptides of the invention can be prepared by various means (*e.g.* recombinant expression, purification from cell culture, chemical synthesis, *etc.*) and in various forms (*e.g.* native, fusions, non-glycosylated, lipidated, *etc.*). They are preferably prepared in

substantially pure form (*i.e.* substantially free from other *Chlamydial* or host cell proteins).

5 The invention also provides a process for producing a polypeptide of the invention, comprising the step of culturing a host cell transformed with nucleic acid of the invention under conditions which induce polypeptide expression. The invention provides a process for producing a polypeptide of the invention, comprising the step of synthesising at least part of the polypeptide by chemical means. The invention further provides a process for producing a composition according to the invention comprising the step of bringing one or more of SEQ IDs 1-76 into combination with
10 one or more other of SEQ IDs 1-76

Strains

Preferred polypeptides of the invention comprise an amino acid sequence found in *C.pneumoniae* serovar D, or in one or more of an epidemiologically prevalent
15 serotype. Where hybrid polypeptides are used, the individual antigens within the hybrid (*i.e.* individual -X- moieties) may be from one or more strains. Where $n=2$, for instance, X_2 may be from the same strain as X_1 or from a different strain. Where $n=3$, the strains might be (i) $X_1=X_2=X_3$ (ii) $X_1=X_2 \neq X_3$ (iii) $X_1 \neq X_2 = X_3$ (iv) $X_1 \neq X_2 \neq X_3$ or (v) $X_1 = X_3 \neq X_2$, *etc.*

Heterologous host

Whilst expression of the polypeptides of the invention may take place in *Chlamydia*, the invention preferably utilises a heterologous host. The heterologous host may be prokaryotic (*e.g.* a bacterium) or eukaryotic. It is preferably *E.coli*, but other suitable
25 hosts include *Bacillus subtilis*, *Vibrio cholerae*, *Salmonella typhi*, *Salmonella typhimurium*, *Neisseria lactamica*, *Neisseria cinerea*, *Mycobacteria* (*e.g.* *M.tuberculosis*), yeasts, *etc.*

Details as to how the molecules which make up the SEQ IDs 1-76 can be produced and used can be found from the relevant international applications such as WO 00/37494, WO 02/02606 and WO 03/049762 and WO 03/068811 and these details need not be repeated here. Where the composition includes a protein that exists in different nascent and mature forms, the mature form of the protein is preferably used. For example, the mature form of the *Chlamydia pneumoniae* protein lacking the signal peptide may be
35 used

ADMINISTRATION

Compositions of the invention will generally be administered directly to a patient. Direct delivery may be accomplished by parenteral injection (*e.g.* subcutaneously, intraperitoneally, intravenously, intramuscularly, or to the interstitial space of a
40 tissue), or by rectal, oral (*e.g.* tablet, spray), vaginal, topical, transdermal (*e.g.* see WO99/27961) or transcutaneous (*e.g.* WO02/074244 and WO02/064162 intranasal (*e.g.* see WO03/028760) ocular, aural, pulmonary or other mucosal administration. The invention may be used to elicit systemic and/or mucosal immunity.

45 The compositions of the present invention may be administered, either alone or as part of a composition, via a variety of different routes. Certain routes may be favoured for certain compositions, as resulting in the generation of a more effective immune response, preferably a CMI response, or as being less likely to induce side effects, or as being easier for administration.
50

By way of example, the compositions of the present invention may be administered via a systemic route or a mucosal route or a transdermal route or it may be administered directly into a specific tissue. As used herein, the term "systemic administration" includes but is not limited to any parenteral routes of administration.

5 In particular, parenteral administration includes but is not limited to subcutaneous, intraperitoneal, intravenous, intraarterial, intramuscular, or intrasternal injection, intravenous, intraarterial, or kidney dialytic infusion techniques. Preferably, the systemic, parenteral administration is intramuscular injection.

10 In one preferred embodiment of the method, the compositions of the present invention are administered via a transdermal route. While it is believed that any accepted mode and route of immunization can be employed and nevertheless achieve some advantages in accordance herewith, the examples below demonstrate particular advantages with transdermal NOI administration. In this regard, and without being bound by theory, it is believed that transdermal administration of a composition may
15 be preferred because it more efficiently activates the cell mediated immune (CMI) arm of the immune system.

The term "transdermal" delivery intends intradermal (e.g., into the dermis or epidermis), transdermal (e.g., "percutaneous") and transmucosal administration, i.e.,
20 delivery by passage of an agent into or through skin or mucosal tissue. See, e.g., *Transdermal Drug Delivery: Developmental Issues and Research Initiatives*, Hadgraft and Guy (eds.), Marcel Dekker, Inc., (1989); *Controlled Drug Delivery: Fundamentals and Applications*, Robinson and Lee (eds.), Marcel Dekker Inc., (1987);
25 and *Transdermal Delivery of Drugs*, Vols. 1-3, Kydonieus and Berner (eds.), CRC Press, (1987). Thus, the term encompasses delivery of an agent using a particle delivery device (e.g., a needleless syringe) such as those described in U.S. Patent No. 5,630,796, as well as delivery using particle-mediated delivery devices such as those described in U.S. Patent No. 5,865,796.

30 As used herein, the term "mucosal administration" includes but is not limited to oral, intranasal, intravaginal, intrarectal, intratracheal, intestinal and ophthalmic administration.

35 Mucosal routes, particularly intranasal, intratracheal, and ophthalmic are preferred for protection against natural exposure to environmental pathogens such as RSV, flu virus and cold viruses or to allergens such as grass and ragweed pollens and house dust mites. The enhancement of the immune response, preferably the CMI response will enhance the protective effect against a subsequently encountered target antigen such as an allergen or microbial agent.

40 In another preferred embodiment of the present invention, the compositions of the present invention may be administered to cells which have been isolated from the host subject. In this preferred embodiment, preferably the composition is administered to professional antigen presenting cells (APCs), such as dendritic cells. APCs may be
45 derived from a host subject and modified *ex vivo* to express an antigen of interest and then transferred back into the host subject to induce an enhanced CMI response. Dendritic cells are believed to be the most potent APCs for stimulating enhanced CMI responses because the expressed epitopes of the antigen of interest must be acquired, processed and presented by professional APCs to T cells (both Th1 and Th2 helper
50 cells as well as CD8+ T-cells) in order to induce an enhanced CMI response.

PARTICLE ADMINISTRATION

Particle-mediated methods for delivering the compositions of the present invention are known in the art. Thus, once prepared and suitably purified, the above-described antigens or NOI encoding same can be coated onto core carrier particles using a variety of techniques known in the art. Carrier particles are selected from materials which have a suitable density in the range of particle sizes typically used for intracellular delivery from a gene gun device. The optimum carrier particle size will, of course, depend on the diameter of the target cells.

By "core carrier" is meant a carrier on which a guest antigen or guest nucleic acid (e.g., DNA, RNA) is coated in order to impart a defined particle size as well as a sufficiently high density to achieve the momentum required for cell membrane penetration, such that the guest molecule can be delivered using particle-mediated techniques (see, e.g., U.S. Patent No. 5,100,792). Core carriers typically include materials such as tungsten, gold, platinum, ferrite, polystyrene and latex. See e.g., *Particle Bombardment Technology for Gene Transfer*, (1994) Yang, N. ed., Oxford University Press, New York, NY pages 10-11. Tungsten and gold particles are preferred. Tungsten particles are readily available in average sizes of 0.5 to 2.0 microns in diameter. Gold particles or microcrystalline gold (e. g., gold powder A1570, available from Engelhard Corp., East Newark, NJ) will also find use with the present invention. Gold particles provide uniformity in size (available from Alpha Chemicals in particle sizes of 1-3 microns, or available from Degussa, South Plainfield, NJ in a range of particle sizes including 0.95 microns). Microcrystalline gold provides a diverse particle size distribution, typically in the range of 0.5-5 microns. However, the irregular surface area of microcrystalline gold provides for highly efficient coating with nucleic acids. A number of methods are known and have been described for coating or precipitating NOIs onto gold or tungsten particles. Most such methods generally combine a predetermined amount of gold or tungsten with plasmid DNA, CaCl₂ and spermidine. The resulting solution is vortexed continually during the coating procedure to ensure uniformity of the reaction mixture. After precipitation of the NOI, the coated particles can be transferred to suitable membranes and allowed to dry prior to use, coated onto surfaces of a sample module or cassette, or loaded into a delivery cassette for use in particular gene gun instruments.

The particle compositions or coated particles are administered to the individual in a manner compatible with the dosage formulation, and in an amount that will be effective for the purposes of the invention. The amount of the composition to be delivered (e. g., about 0.1 mg to 1 mg; more preferably 1 to 50 ug of the antigen or allergen, depends on the individual to be tested. The exact amount necessary will vary depending on the age and general condition of the individual to be treated, and an appropriate effective amount can be readily determined by one of skill in the art upon reading the instant specification.

HOST MAMMALIAN SUBJECT

As used herein, the term "host mammalian subject" means any member of the subphylum cordata, including, without limitation, humans and other primates, including non-human primates such as chimpanzees and other apes and monkey species; farm animals such as cattle, sheep, pigs, goats and horses; domestic mammals such as dogs and cats; laboratory animals including rodents such as mice, rats and

guinea pigs; birds, including domestic, wild and game birds such as chickens, turkeys and other gallinaceous birds, ducks, geese, and the like. The terms do not denote a particular age. Thus, both adult and newborn individuals are intended to be covered. The methods described herein are intended for use in any of the above vertebrate species, since the immune systems of all of these vertebrates operate similarly. If a mammal, the subject will preferably be a human, but may also be a domestic livestock, laboratory subject or pet animal.

The mammal is preferably a human. Where the vaccine is for prophylactic use, the human is preferably a child (e.g. a toddler or infant) or a teenager; where the vaccine is for therapeutic use, the human is preferably a teenager or an adult. A vaccine intended for children may also be administered to adults e.g. to assess safety, dosage, immunogenicity, etc.

PREVENT AND/OR TREAT

The invention also provides the use of the compositions of the invention in the manufacture of a medicament for raising an immune response in a mammal. The medicament is preferably a vaccine and to the preparation of a vaccine to prevent and/or treat an disorder associated with a *Chlamydia* bacterium. It is to be appreciated that all references herein to treatment include curative, palliative and prophylactic treatment.

The administration of antigenic combinations of the present invention or a composition comprising the NOI encoding the antigenic combinations may be for either "prophylactic" or "therapeutic" purpose. As used herein, the term "therapeutic" or "treatment" includes any of following: the prevention of infection or reinfection; the reduction or elimination of symptoms; and the reduction or complete elimination of a pathogen. Treatment may be effected prophylactically (prior to infection) or therapeutically (following infection).

Prophylaxis or therapy includes but is not limited to eliciting an effective immune response, preferably a CMI immune response and/or alleviating, reducing, curing or at least partially arresting symptoms and/or complications resulting from a T cell mediated immune disorder. When provided prophylactically, the composition of the present invention is typically provided in advance of any symptom. The prophylactic administration of the composition of the present invention is to prevent or ameliorate any subsequent infection or disease. When provided therapeutically, the composition of the present invention is typically provided at (or shortly after) the onset of a symptom of infection or disease. Thus the composition of the present invention may be provided either prior to the anticipated exposure to a disease causing agent or disease state or after the initiation of an infection or disease.

Whether prophylactic or therapeutic administration (either alone or as part of a composition) is the more appropriate will usually depend upon the nature of the disease. By way of example, immunotherapeutic composition of the present invention could be used in immunotherapy protocols to actively inducing immunity by vaccination. This latter form of treatment is advantageous because the immunity is prolonged. On the other hand a vaccine composition will preferably, though not necessarily be used prophylactically to induce an effective CMI response against subsequently encountered antigens or portions thereof (such as epitopes) related to the target antigen.

5 These uses and methods are preferably for the prevention and/or treatment of a disease caused by a *Chlamydia* (e.g. trachoma, pelvic inflammatory disease, epididymitis, infant pneumonia, arteriosclerosis, cardiovascular disease etc.). The compositions may also be effective against *C.pneumoniae*.

PROPHYLACTICALLY OR THERAPEUTICALLY OR IMMUNOLOGICALLY EFFECTIVE AMOUNT

10 The composition dose administered to a host subject, in the context of the present invention, should be sufficient to effect a beneficial prophylactic or therapeutic immune response, preferably a CMI response in the subject over time.

15 The invention also provides a method for raising an immune response in a mammal comprising the step of administering an effective amount of a composition of the invention. The immune response is preferably protective and preferably involves antibodies and/or cell-mediated immunity. The method may raise a booster response.

20 As used herein, the term "'prophylactically or therapeutically effective dose" means a dose in an amount sufficient to elicit an enhanced immune response, preferably a CMI response to one or more antigens or epitopes and/or to alleviate, reduce, cure or at least partially arrest symptoms and/or complications from a T cell mediated immune disorder.

25 Immunogenic compositions used as vaccines comprise an immunologically effective amount of antigen(s), as well as any other components, as needed. By 'immunologically effective amount', it is meant that the administration of that amount to an individual, either in a single dose or as part of a series, is effective for treatment or prevention. This amount varies depending upon the health and physical condition of the individual to be treated, age, the taxonomic group of individual to be treated
30 (e.g. non-human primate, primate, etc.), the capacity of the individual's immune system to synthesise antibodies, the degree of protection desired, the formulation of the vaccine, the treating doctor's assessment of the medical situation, and other relevant factors. It is expected that the amount will fall in a relatively broad range that can be determined through routine trials.

35 One way of checking efficacy of therapeutic treatment involves monitoring *Chlamydia* infection after administration of the composition of the invention. One way of checking efficacy of prophylactic treatment involves monitoring immune responses against the *Chlamydia* antigen, such as the *Chlamydia pneumoniae* antigen
40 in the compositions of the invention after administration of the composition.

The vaccine compositions of the present invention can be evaluated in *in vitro* and *in vivo* animal models prior to host, e.g., human, administration. For example, *in vitro* neutralization by Peterson *et al* (1988) is suitable for testing vaccine compositions directed toward *Chlamydia*, preferably *Chlamydia pneumoniae*.
45

One example of such an *in vitro* test is described as follows. Hyper-immune antisera is diluted in PBS containing 5% guinea pig serum, as a complement source. *Chlamydia pneumoniae* (10^4 IFU; inclusion forming units) are added to the antisera dilutions. The antigen-antibody mixtures are incubated at 37°C for 45 minutes and
50

5 inoculated into duplicate confluent Hep-2 or HeLa cell monolayers contained in glass vials (e.g., 15 by 45 mm), which have been washed twice with PBS prior to inoculation. The monolayer cells are infected by centrifugation at 1000X g for 1 hour followed by stationary incubation at 37°C for 1 hour. Infected monolayers are incubated for 48 or 72 hours, fixed and stained with *Chlamydia* specific antibody, such as anti-MOMP. Inclusion-bearing cells are counted in ten fields at a magnification of 200X. Neutralization titer is assigned on the dilution that gives 50% inhibition as compared to control monolayers/IFU.

10 The efficacy of vaccine compositions can also be determined *in vivo* by challenging animal models of *Chlamydia pneumoniae* infection, e.g., guinea pigs or mice, with the vaccine compositions. For example, *in vivo* vaccine composition challenge studies in the guinea pig model of *Chlamydia pneumoniae* infection can be performed. A description of one example of this type of approach follows. Female guinea pigs
15 weighing 450 – 500 g are housed in an environmentally controlled room with a 12 hour light-dark cycle and immunized with vaccine compositions via a variety of immunization routes. Post-vaccination, guinea pigs are infected in the genital tract with the agent of guinea pig inclusion conjunctivitis (GPIC), which has been grown in HeLa or McCoy cells (Rank *et al.* (1988)). Each animal receives approximately
20 1.4×10^7 inclusion forming units (IFU) contained in 0.05 ml of sucrose-phosphate-glutamate buffer, pH 7.4 (Schacter, 1980). The course of infection monitored by determining the percentage of inclusion-bearing cells by indirect immunofluorescence with GPIC specific antisera, or by Giemsa-stained smear from a scraping from the genital tract (Rank *et al.* 1988). Antibody titers in the serum is determined by an
25 enzyme-linked immunosorbent assay.

Alternatively, *in vivo* vaccine compositions challenge studies can be performed in the murine model of *Chlamydia pneumoniae* (Morrison *et al.* 1995). A description of one example of this type of approach is as follows. Female mice 7 to 12 weeks of age
30 receive 2.5 mg of depoprovera subcutaneously at 10 and 3 days before vaginal infection. Post-vaccination, mice are infected in the genital tract with 1,500 inclusion-forming units of *Chlamydia pneumoniae* contained in 5ml of sucrose-phosphate-glutamate buffer, pH 7.4. The course of infection is monitored by determining the percentage of inclusion-bearing cells by indirect immunofluorescence with *Chlamydia*
35 *pneumoniae* specific antisera, or by a Giemsa-stained smear from a scraping from the genital tract of an infected mouse. The presence of antibody titers in the serum of a mouse is determined by an enzyme-linked immunosorbent assay

DOSAGE

40 Prophylaxis or therapy can be accomplished by a single direct administration at a single time point or multiple time points. Administration can also be delivered to a single or to multiple sites. Some routes of administration, such as mucosal administration via ophthalmic drops may require a higher dose. Those skilled in the art can adjust the dosage and concentration to suit the particular route of delivery.
45

Dosage treatment can be a single dose schedule or a multiple dose schedule. multiple doses may be used in a primary immunisation schedule and/or in a booster immunisation schedule. in a multiple dose schedule the various doses may be given by the same or different routes *e.g.* a parenteral prime and mucosal boost, a mucosal prime and parenteral boost, *etc.*

HOMOLOGUES

SEQ IDs 1-76 in the compositions of the invention may be supplemented or substituted with molecules comprising sequences homologous (ie. sharing sequence identity) to SEQ ID Nos 1-76.

Proteins (including protein antigens) as used in the invention (as encoded by the NOI) may have homology and/or sequence identity with naturally occurring forms. Similarly coding sequences capable of expressing such proteins will generally have homology and/or sequence identity with naturally occurring sequences. Techniques for determining nucleic acid and amino acid "sequence identity" also are known in the art. Typically, such techniques include determining the nucleotide sequence of the mRNA for a gene and/or determining the amino acid sequence encoded thereby, and comparing these sequences to a second nucleotide or amino acid sequence.

In general, "identity" refers to an exact nucleotide-to-nucleotide or amino acid-to-amino acid correspondence of two polynucleotides or polypeptide sequences, respectively. Two or more sequences (polynucleotide or amino acid) can be compared by determining their "percent identity." The percent identity of two sequences, whether nucleic acid or amino acid sequences, is the number of exact matches between two aligned sequences divided by the length of the shorter sequences and multiplied by 100.

An approximate alignment for nucleic acid sequences is provided by the local homology algorithm of Smith and Waterman, *Advances in Applied Mathematics* 2: 482-489 (1981). This algorithm can be applied to amino acid sequences by using the scoring matrix developed by Dayhoff, *Atlas of Protein Sequences and Structure*, M. O. Dayhoff ed., 5 suppl. 3: 353-358, National Biomedical Research Foundation, Washington, D. C., USA, and normalized by Gribskov, *Nucl. Acids Res.* 14 (6): 6745-6763 (1986). An exemplary implementation of this algorithm to determine percent identity of a sequence is provided by the Genetics Computer Group (Madison, WI) in the "BestFit" utility application. The default parameters for this method are described in the Wisconsin Sequence Analysis Package Program Manual, Version 8 (1995) (available from Genetics Computer Group, Madison, WI). A preferred method of establishing percent identity in the context of the present invention is to use the MPSRCH package of programs copyrighted by the University of Edinburgh, developed by John F. Collins and Shane S.

Sturrok, and distributed by IntelliGenetics, Inc. (Mountain View, CA). From this suite of packages the Smith-Waterman algorithm can be employed where default parameters are used for the scoring table (for example, gap open penalty of 12, gap extension penalty of one, and a gap of six). From the data generated the "Match" value reflects "sequence identity." Other suitable programs for calculating the percent identity or similarity between sequences are generally known in the art, for example, another alignment program is BLAST, used with default parameters. For example,

5 BLASTN and BLASTP can be used using the following default parameters: genetic code = standard; filter = none; strand = both; cutoff= 60; expect = 10; Matrix = BLOSUM62 ; Descriptions = 50 sequences; sort by = HIGH SCORE; Databases = non-redundant, GenBank +EMBL + DDBJ + PDB + GenBank CDS translations + Swiss protein + Spupdate + PIR. Details of these programs can be found at the following internet address: <http://www.ncbi.nlm.gov/cgi-bin/BLAST>.

10 Alternatively, homology can be determined by hybridization of polynucleotides under conditions which form stable duplexes between homologous regions, followed by digestion with single-stranded-specific nuclease (s), and size determination of the digested fragments. Two DNA, or two polypeptide sequences are "substantially homologous" to each other when the sequences exhibit at least about 80%-85%, preferably at least about 90%, and most preferably at least about 95%-98% sequence identity over a defined length of the molecules, as determined using the methods above.

20 As used herein, substantially homologous or homologous also refers to sequences showing complete identity to the specified DNA or polypeptide sequence. DNA sequences that are substantially homologous or homologous can be identified in a Southern hybridization experiment under, for example, stringent conditions, as defined for that particular system. For example, stringent hybridization conditions can include 50% formamide, 5x Denhardt's Solution, 5x SSC, 0.1% SDS and 100 pg/ml denatured salmon sperm DNA and the washing conditions can include 2x SSC, 0.1% SDS at 37 C followed by 1x SSC, 0.1% SDS at 68 C. Defining appropriate hybridization conditions is within the skill of the art.

30 Preferably the degree of identity is preferably greater than 50% (eg. 65%. 80%. 90%. or more) and include mutants and allelic variants. Sequence identity between the proteins is preferably determined by the Smith-Waterman homology search algorithm as implemented in the MPSRCH program (Oxford. Molecular). using an affine gap search with parameters gap open penalty=12 and gap extension penalty=1.

35 SEQ IDs 1-76 in the compositions of the invention may be supplemented or substituted with nucleic acid which can hybridise to the *Chlamydia* nucleic acid. preferably under "high stringency" conditions (c. 65 C in an 0.1 x SSC, 0.5% SDS solution).

Hypothetical Protein

40 As used herein, the term "hypothetical protein" refers to a protein which lacks a known cellular location or a known cellular function. Typically, a hypothetical protein lacks significant homologies with known well characterised proteins.

COMPOSITIONS

45 The invention also provides the compositions of the invention for use as medicaments (eg. as immunogenic compositions or vaccines) or as diagnostic reagents for detecting a *Chlamydia* infection in a host subject. It also provides the use of the compositions in the manufacture of: (i) a medicament for treating or preventing infection due to *Chlamydia pneumoniae* bacteria; (ii) a diagnostic reagent for detecting the presence of *Chlamydia Pneumoniae* bacteria or of antibodies raised against *Chlamydia* Pneumoniae bacteria; and/or (iii) a reagent which can raise antibodies against

50

Chlamydia

pneumoniae

bacteria.

5 The invention also provides a method of treating a patient, comprising administering to the patient a therapeutically effective amount of a composition according to the invention.

10 The present invention provides compositions that are useful for preventing and/or treating T cell mediated immune disorders. In one embodiment, the composition is a pharmaceutical composition. In another preferred embodiment, the composition is an immunotherapeutic composition. In an even more preferred embodiment, the composition is a vaccine composition.. The composition may also comprise a carrier such as a pharmaceutically or immunologically acceptable carrier. Pharmaceutically acceptable carriers or immunologically acceptable carriers are determined in part by the particular composition being administered as well as by the particular method used to administer the composition. Accordingly, there is a wide variety of suitable formulations of pharmaceutical compositions or vaccine compositions or immunotherapeutic compositions of the present invention.

Immunogenic compositions and medicaments

20 Compositions of the invention are preferably immunogenic compositions, and are more preferably vaccine compositions. The pH of the composition is preferably between 6 and 8, preferably about 7. The pH may be maintained by the use of a buffer. The composition may be sterile and/or pyrogen-free. The composition may be isotonic with respect to humans.

25 Vaccines according to the invention may either be prophylactic (*i.e.* to prevent infection) or therapeutic (*i.e.* to treat infection), but will typically be prophylactic. Accordingly, the invention includes a method for the therapeutic or prophylactic treatment of *Chlamydia pneumoniae* infection in an animal susceptible to *Chlamydial* infection comprising administering to said animal a therapeutic or prophylactic amount of the immunogenic compositions of the invention. Preferably, the immunogenic composition comprises a combination of *Chlamydia pneumoniae* antigens, said combination selected from the group consisting of two, three, four, five or all six *Chlamydia pneumoniae* antigens of the first antigen group. Still more preferably, the combination consists of all six *Chlamydia pneumoniae* antigens of the first antigen group.

40 Alternatively, the immunogenic composition comprises a combination of *Chlamydia pneumoniae* antigens, said combination selected from the group consisting of two, three, four, five, six, seven, eight, nine, ten, eleven, or twelve *Chlamydia pneumoniae* antigens selected from the first antigen group and the second antigen group. Preferably, the combination is selected from the group consisting of three, four, or five *Chlamydia pneumoniae* antigens selected from the second antigen group. Still more preferably, the combination consists of five *Chlamydia pneumoniae* antigens selected from the second antigen group.

50 Alternatively, the immunogenic composition comprises a combination of *Chlamydia pneumoniae* antigens, said combination consisting of two, three, four, or five *Chlamydia pneumoniae* antigens of the first antigen group and one, two, three, four, five or six *Chlamydia pneumoniae* antigens of the third antigen group. Preferably, the

combination consists of three, four or five *Chlamydia pneumoniae* antigens of the first antigen group and one, two, three, four, five or six *Chlamydia pneumoniae* antigens of the third antigen group.

5 Alternatively, the immunigenic composition comprises a combination of *Chlamydia pneumoniae* antigens, said combination consisting of two, three, four, five, six, seven, eight, nine, ten, eleven or twelve *Chlamydia pneumoniae* antigens of the first antigen group and the second antigen group and one, two, three, four, five or six *Chlamydia pneumoniae* antigens of the third antigen group. Preferably, the combination is
10 selected from the group consisting of three, four, or five *Chlamydia pneumoniae* antigens from the second antigen group and three, four or five *Chlamydia pneumoniae* from the third antigen group. Still more preferably, the combination consists of five *Chlamydia pneumoniae* antigens from the second antigen group and three, four or five *Chlamydia pneumoniae* antigens of the third antigen group.

15 In certain embodiments, the composition comprises molecules from different *Chlamydia* species. In some embodiments, the composition may comprise molecules from different serogroups and/or strains of the same *Chlamydia* species. Further
20 embodiments comprise mixtures of one or more *Chlamydia* molecules from different strains.

Many proteins are relatively conserved between different species serogroups and strains of *Chlamydia trachomatis* and *Chlamydia pneumoniae*. To ensure maximum cross-strain recognition and reactivity, regions of proteins that are conserved between
25 different *Chlamydia* species, serogroups and strains can be used in the compositions of the present invention. The invention therefore provides proteins which comprise stretches of amino acid sequence that are shared across the majority of *Chlamydia* strains. Preferably, therefore, the composition comprises a protein comprising a
30 fragment of a *Chlamydia pneumoniae* protein (preferably a protein from SEQ ID Nos 1-76 or more preferably SEQ ID Nos 1-41 wherein said fragment consists of n consecutive conserved amino acids.

Further antigens

35 The compositions of the invention may further comprise antigen derived from one or more sexually transmitted diseases in addition to *Chlamydia trachomatis*. Preferably the antigen is derived from one or more of the following sexually transmitted diseases: *N.gonorrhoeae* {e.g. i, ii, iii, iv}; human papilloma virus; *Treponema pallidum*; herpes simplex virus (HSV-1 or HSV-2); HIV (HIV-1 or HIV-2); and
40 *Haemophilus ducreyi*.

A preferred composition comprises: (1) at least t of the *Chlamydia pneumoniae* antigens from either the first antigen group or the second antigen group, where t is 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12 or 13, preferably t is five; (2) one or more antigens from another sexually transmitted disease. Preferably, the sexually transmitted disease is
45 selected from the group consisting of herpes simplex virus, preferably HSV-1 and/or HSV-2; human papillomavirus; *N.gonorrhoeae*; *Treponema pallidum*; and *Haemophilus ducreyi*. These compositions can thus provide protection against the following sexually-transmitted diseases: *Chlamydia*, genital herpes, genital warts, gonorrhoea, syphilis and chancroid (see Stephens et al (1998) Science 282: 754-759).
50

Where a saccharide or carbohydrate antigen is used, it is preferably conjugated to a carrier protein in order to enhance immunogenicity (For example, Ramsay *et al.* (2001) *Lancet* 357(9251):195-196; Lindberg (1999) *Vaccine* 17 Suppl 2:S28-36; Buttery & Moxon (2000) *J R Coll Physicians Lond* 34:163-168; Ahmad & Chapnick (1999) *Infect Dis Clin North Am* 13:113-133; Goldblatt (1998) *J. Med. Microbiol.* 47:563-567; European patent 0 477 508; US Patent No. 5,306,492; International patent application WO98/42721; *Conjugate Vaccines* (eds. Cruse *et al.*) ISBN 3805549326, particularly vol. 10:48-114; and Hermanson (1996) *Bioconjugate Techniques* ISBN: 0123423368 or 012342335).

Preferred carrier proteins are bacterial toxins or toxoids, such as diphtheria or tetanus toxoids. The CRM₁₉₇ diphtheria toxoid is particularly preferred (*Research Disclosure*, 453077 (Jan 2002). Other carrier polypeptides include the *N.meningitidis* outer membrane protein EP-A-0372501), synthetic peptides (EP-A-0378881, EP-A-0427347), heat shock proteins (WO93/17712, WO94/03208) pertussis proteins (WO98/58668, EP-A-0471177) protein D from *H.influenzae* (WO00/56360) cytokines (WO91/01146), lymphokines, hormones, growth factors, toxin A or B from *C.difficile* (WO00/61761) iron-uptake proteins WO01/72337) *etc.* Where a mixture comprises capsular saccharides from both serogroups A and C, it may be preferred that the ratio (w/w) of MenA saccharide:MenC saccharide is greater than 1 (e.g. 2:1, 3:1, 4:1, 5:1, 10:1 or higher). Different saccharides can be conjugated to the same or different type of carrier protein. Any suitable conjugation reaction can be used, with any suitable linker where necessary.

Toxic protein antigens may be detoxified where necessary e.g. detoxification of pertussis toxin by chemical and/or genetic means. Where a diphtheria antigen is included in the composition it is preferred also to include tetanus antigen and pertussis antigens. Similarly, where a tetanus antigen is included it is preferred also to include diphtheria and pertussis antigens. Similarly, where a pertussis antigen is included it is preferred also to include diphtheria and tetanus antigens.

Antigens in the composition will typically be present at a concentration of at least 1µg/ml each. In general, the concentration of any given antigen will be sufficient to elicit an immune response against that antigen. As an alternative to using protein antigens in the composition of the invention, nucleic acid encoding the antigen may be used Robinson & Torres (1997) *Seminars in Immunology* 9:271-283; Donnelly *et al.* (1997) *Annu Rev Immunol* 15:617-648; Scott-Taylor & Dalglish (2000) *Expert Opin Investig Drugs* 9:471-480; Apostolopoulos & Plebanski (2000) *Curr Opin Mol Ther* 2:441-447; Ilan (1999) *Curr Opin Mol Ther* 1:116-120; Dubensky *et al.* (2000) *Mol Med* 6:723-732; Robinson & Pertmer (2000) *Adv Virus Res* 55:1-74; Donnelly *et al.* (2000) *Am J Respir Crit Care Med* 162(4 Pt 2):S190-193 and Davis (1999) *Mt. Sinai J. Med.* 66:84-90. Protein components of the compositions of the invention may thus be replaced by nucleic acid (preferably DNA e.g. in the form of a plasmid) that encodes the protein.

DISEASE STATES

The compositions of the present invention may be used to prevent and/or treat disorders such as but not limited to: pneumonia, cardiovascular diseases, atherosclerosis, bronchitis, pharyngitis, laryngitis, sinusitis, obstructive lung diseases, asthma, chronic obstructive pulmonary disease, reactive arthritis, otitis media,

5 abdominal aortic aneurysm, erythema nodosum, Reiter syndrome, sarcoidosis, Alzheimer's disease, multiple sclerosis, lymphogranuloma venereum, ocular trachoma, pelvic inflammatory disease, inclusion conjunctivitis, genital trachoma, infant pneumonitis, incipient trachoma, keratitis, papillary hypertrophy, corneal infiltration, vulvovaginitis, mucopurulent rhinitis, salpingitis, cervicitis, cervical follicles, prostatitis, proctitis, urethritis, lymphogranule inguinale, climatic bubo, tropical bubo, and/or oresthiomene.

FORMULATIONS

10 *Chlamydial* infections affect various areas of the body and so the compositions of the invention may be prepared in various forms. For example, the compositions may be prepared as injectables, either as liquid solutions or suspensions. Solid forms suitable for solution in, or suspension in, liquid vehicles prior to injection can also be prepared (e.g. a lyophilised composition). The composition may be prepared for topical
15 administration e.g. as an ointment, cream or powder. The composition may be prepared for oral administration e.g. as a tablet or capsule, as a spray, or as a syrup (optionally flavoured). The composition may be prepared for pulmonary administration e.g. as an inhaler, using a fine powder or a spray. The composition may be prepared as a suppository or pessary. The composition may be prepared for nasal,
20 aural or ocular administration e.g. as drops. The composition may be in kit form, designed such that a combined composition is reconstituted just prior to administration to a patient. Such kits may comprise one or more antigens in liquid form and one or more lyophilised antigens.

25 *Further components of the composition*

The composition of the invention will typically, in addition to the components mentioned above, comprise one or more 'pharmaceutically acceptable carriers', which include any carrier that does not itself induce the production of antibodies harmful to the individual receiving the composition. Suitable carriers are typically large, slowly
30 metabolised macromolecules such as proteins, polysaccharides, polylactic acids, polyglycolic acids, polymeric amino acids, amino acid copolymers, and lipid aggregates (such as oil droplets or liposomes). Such carriers are well known to those of ordinary skill in the art. The vaccines may also contain diluents, such as water, saline, glycerol, etc. Additionally, auxiliary substances, such as wetting or
35 emulsifying agents, pH buffering substances, and the like, may be present. A thorough discussion of pharmaceutically acceptable excipients is available in Gennaro (2000) *Remington: The Science and Practice of Pharmacy*. 20th ed., ISBN: 0683306472.

The biological molecules of the present invention be formulated into a pharmaceutical composition or an immunotherapeutic composition or a vaccine composition. Such
40 formulations comprise biological molecules combined with a pharmaceutically acceptable carrier, such as sterile water or sterile isotonic saline. Such formulations may be prepared, packaged, or sold in a form suitable for bolus administration or for continuous administration. Injectable formulations may be prepared, packaged, or sold
45 in unit dosage form, such as in ampoules or in multi-dose containers containing a preservative. Formulations include, but are not limited to, suspensions, solutions, emulsions in oily or aqueous vehicles, pastes, and implantable sustained-release or biodegradable formulations. Such formulations may further comprise one or more additional ingredients including, but not limited to, suspending, stabilizing, or
50 dispersing agents. In one embodiment of a formulation for parenteral administration,

the active ingredient is provided in dry (for eg, a powder or granules) form for reconstitution with a suitable vehicle (e. g., sterile pyrogen-free water) prior to parenteral administration of the reconstituted composition. The pharmaceutical compositions may be prepared, packaged, or sold in the form of a sterile injectable aqueous or oily suspension or solution. This suspension or solution may be formulated according to the known art, and may comprise, in addition to the active ingredient, additional ingredients such as the dispersing agents, wetting agents, or suspending agents described herein. Such sterile injectable formulations may be prepared using a non-toxic parenterally-acceptable diluent or solvent, such as water or 1,3-butane diol, for example. Other acceptable diluents and solvents include, but are not limited to, Ringer's solution, isotonic sodium chloride solution, and fixed oils such as synthetic mono- or di-glycerides. Other parentally-administrable formulations which are useful include those which comprise the active ingredient in microcrystalline form, in a liposomal preparation, or as a component of a biodegradable polymer systems. Compositions for sustained release or implantation may comprise pharmaceutically acceptable polymeric or hydrophobic materials such as an emulsion, an ion exchange resin, a sparingly soluble polymer, or a sparingly soluble salt.

KITS

Also included in the invention is a kit for enhancing a CMI response to the biological molecules of the present invention. Such a kit may comprise an antigenic composition or nucleotide sequence encoding same. The kit may also include an adjuvant, preferably a genetic adjuvant is administered with or as part of the biological molecule and instructions for administering the biological molecule. Other preferred components of the kit include an applicator for administering the biological molecule. As used herein, the term "applicator" refers to any device including but not limited to a hypodermic syringe, gene gun, particle acceleration device, nebulizer, dropper, bronchoscope, suppository, impregnated or coated vaginally-insertable material such as a tampon, douche preparation, solution for vaginal irrigation, retention enema preparation, suppository, or solution for rectal or colonic irrigation for applying the NOI either systemically or mucosally or transdermally to the host subject.

The invention also provides for a kit comprising comprising a combination of *Chlamydia pneumoniae* antigens. The combination of *Chlamydia pneumoniae* antigens may be one or more of the immunogenic compositions of the invention. The kit may further include a second component comprising one or more of the following: instructions, syringe or other delivery device, adjuvant, or pharmaceutically acceptable formulating solution. The invention also provides a delivery device pre-filled with the immunogenic compositions of the invention.

EXAMPLES

The following invention will now be further described only by way of example in which reference is made to the following Figures. The following examples are presented only to illustrate the present invention and to assist one of ordinary skill in making and using the same. The examples are not intended in any way to otherwise limit the scope of the invention. Efforts have been made to ensure accuracy with respect to numbers used (e.g., amounts, temperatures, etc.), but some experimental error and deviation should, of course, be allowed for.

5
10
15
20
Figure 1A. Assay of *in vitro* neutralization of *C.pneumoniae* infectivity for LLC-MK2 cells by polyclonal mouse antisera to recombinant *Chlamydial* proteins. Results are shown as reduction in the number of inclusions obtained when monolayers were infected with antiserum-treated infectious EBs, as compared to inclusion numbers given by untreated EBs. Percent reduction values are plotted against the reciprocal of the corresponding serum dilution. For each dilution inclusion counts were corrected for background inhibition of infectivity observed with the corresponding dilution of the pre-immune serum. The figure shows results obtained with serial dilutions of antibodies raised against a 'neutralizing' antigen (♥), a 'non-neutralizing' FACS- positive antigen (v), and against the GST polypeptide, used in the fusion constructs, alone (σ).

25
30
Figure 1B shows serum titres giving 50% neutralization of infectivity for the 10 *C.pneumoniae* recombinant antigens described in the text. Each titer was assessed in 3 separate experiments (SEM values shown).

35
40
45
Figure 2 shows immunoblot analysis of two dimensional electrophoretic maps of *C.pneumoniae* EBs using the immune sera described in the text. Immunoblots were obtained from either of two EB gels (panels A and B at the top) covering different pH intervals, according to which of the two allowed the best detection of a given antigen. The arrows in the HtrA immunoblot show which of the signals had a corresponding stained spot in the gel (arrows in panel A) which was subjected to MALDI-TOF identification. The two patterns in the HtrA blot are both suggestive of typical electrophoretic 'trains' composed of single charge variants of the same protein.

45
50
Figure 3 shows mean numbers of *C.pneumoniae* IFU recovered from equivalent spleen samples from immunized and mock-immunized hamsters following a systemic challenge. Standard deviation values are shown above the bars. Antigens which induced significant protection are highlighted with an asterisk above the corresponding bar. All antigens were delivered in Freund's adjuvant. n.i. = non immunized controls

55
60
Figure 4 shows flow cytometric analysis of splenocytes from DNA-immunized HLA-A2 transgenic and non transgenic mice. Groups of 4 mice were immunized 3 times i.m. with 50µg of plasmid DNA expressing *C. pneumoniae* Low Calcium Response Protein H. IFN-γ production from splenocytes was monitored following either a 6h (ex-vivo) or a 6 day (restimulated) pulse with peptide CH-6 (10µg/ml). Equal numbers of gated live lymphocyte cells were acquired with a LSRII FACS System (Becton Dickinson) and percentages of IFN-γ producing CD8⁺ T cells were calculated using DIVA Software (Becton Dickinson).

- 5 **Figure 5** shows a flow cytometric analysis of splenocytes from transgenic and non transgenic mice infected with *C. pneumoniae* EBs. (A) HLA-A2 transgenic mice were intranasally infected twice with 5×10^5 *C. pneumoniae* FB/96 EBs and splenocytes were stimulated for 6 days in the presence of relevant peptides before determining IFN- γ production by CD8⁺ T cells as described in the legend of Figure 4. (B) HLA-A2 transgenic and non transgenic mice were infected together with the same EBs preparation and CD8⁺ T cells were subjected to FACS analysis as reported in (A).
- 10 **Table I** shows a summary of data and properties of the *C.pneumoniae* antigens described in the text. The neutralization titer is reported is as the reciprocal of the antiserum dilution causing a 50% reduction in the number of inclusions in the *in vitro* infectivity assay. For the hamster model data the statistical significance of the results was evaluated by a two-tailed Student's t-test: significant data ($p \leq 0.05$) are
- 15 highlighted with an asterisk. ND = not detected.
- Table 2** shows results from hamster mouse model studies for hypothetical proteins.
- 20 **Table 3** shows expressed genes of CPn EB selected by microarray.
- Table 4** shows *C. pneumoniae* selected peptides: protein sources and HLA-A2 stabilization assay.
- 25 **Table 5** shows ELISPOT assay with CD8⁺ T cells from DNA immunised HLA-A2 transgenic mice.
- Table 6** shows IFN- γ production from splenocytes of DNA immunized HLA-A2 transgenic and non transgenic mice.
- 30

METHODS AND MATERIALS (Examples 1-4) (see Reference Section 1)

C.pneumoniae EB purification

5 *C.pneumoniae* FB/96, a clinical isolate obtained from a patient with pneumonia at the Sant'Orsola Polyclinic, Bologna, Italy, was grown in LLC-MK2 cells seeded in individual wells of a six-well plastic plate (7). Cells were harvested 72 hr after infection with a sterile rubber, sonically disrupted and the elementary bodies (EB) purified by gradient centrifugation as described (26). Purified *Chlamydiae* were resuspended in sucrose-phosphate-glutamic acid (SPG) transport buffer, and stored in 0.5 ml aliquots, at -80°C until used. When required, prior to storage, EB infectivity was heat-inactivated by 3 hour incubation at 56°C.

Expression and purification of recombinant proteins

Open reading frames (ORFs), selected from the *C. pneumoniae* CWL029 genome sequence (16), were PCR-cloned into plasmid expression vectors and purified from 15 *E.coli* cultures, essentially as previously described (25). Recombinant *Chlamydial* proteins were obtained as GST fusion proteins by using pGEX-KG derived vectors (12) in *E. coli* BL21 (Novagen). PCR primers were designed so as to amplify genes without the N-terminal signal peptide coding sequence. When a signal peptide or processing site was not clearly predictable, the ORF sequence was cloned as 20 annotated by Kalman and coworkers (16). Recombinant *E.coli* cells were grown in LB medium (500 ml), containing 100 µg/ml Ampicillin, and grown at 37°C until OD₆₀₀ = 0.5, and then induced with 1 mM IPTG. Cells were collected by centrifugation 3 hr after induction and broken in a French Press (SLM Aminco, Rochester, NY). After centrifugation at 30,000 g, the supernatants were loaded onto 25 Glutathione Sepharose 4B columns (Amersham Pharmacia Biotech) and column bound proteins were eluted with 50 mM Tris-HCl, 10 mM reduced glutathione, pH 8.0. Protein concentrations in the samples were determined using the Bradford method.

Preparation of mouse antisera

30 Groups of four 5/6-week old CD1 female mice (Charles River, Como, Italy) were immunized intraperitoneally at day 1 with 20µg of protein in Complete Freund's adjuvant (CFA) and boosted at day 15 and 28 with 20µg of recombinant protein in Incomplete Freund's adjuvant (IFA). Pre-immune and immune sera were prepared 35 from blood samples collected on days 0, 27 and 42. In order to reduce the amount of antibodies possibly elicited by contaminating *E. coli* antigens, the immune sera were incubated overnight at 4°C with nitrocellulose strips adsorbed with a total protein extract from *E. coli* BL21.

Flow cytometry assays

40 Analyses were performed essentially as previously described (25). Gradient purified, heat-inactivated EBs (2x10⁵ cells) from *C.pneumoniae* FB/9, resuspended in phosphate-saline buffer (PBS), 0.1% bovine serum albumin (BSA), were incubated for 30 min. at 4°C with the specific mouse antisera (standard dilution 1:400). After 45 centrifugation and washing with 200 µl of PBS-0.1% BSA, the samples were incubated for 30 minutes at 4°C with Goat Anti-Mouse IgG, F(ab)'₂-specific, conjugated with R-Phycoerythrin (Jackson Immunoresearch Laboratories Inc.). The samples were washed with PBS-0.1%BSA, resuspended in 150 µl of PBS-0.1%BSA and analysed by Flow Cytometry using a FACSCalibur apparatus (Becton Dickinson, 50 Mountain View, CA). Control samples were similarly prepared. Positive control

antibodies were: i), a commercial anti-*C.pneumoniae* specific monoclonal antibody (Argene Biosoft, Varilhes, France) and, ii), a mouse polyclonal serum prepared by immunizing mice with gradient purified *C.pneumoniae* EBs. Background control sera were obtained from mice immunized with the purified GST peptide used in the fusion constructs (GST-fusions control). FACS data were analysed using the Cell Quest Software (Becton Dickinson, Mountain View, CA). The shift between the background control histogram and the immune serum testing histogram was taken as a measure of antibody binding to the EB cell surface. The Kolmorov-Smirnov (K-S) two-sample test (44) was performed on the two overlapped histograms. The D/s(n) values (an index of dissimilarity between the two curves) are reported as "K-S score" in Table 1.

2D Western Blot analysis of immune sera, and Mass Spectrometry

Gradient purified *C. pneumoniae* EBs were washed with 5 mM Tris-HCl pH 7.5, 0.1 mM EDTA, 10% glycerol, centrifuged 15 min. at 13 000 x g and pellets were resuspended in reswelling solution (7 M urea, 2 M thiourea, 2% (w/v) CHAPS, (2%w/v) ASB14, 2% (v/v) IPG buffer pH 3-10 NL, or pH 4-7, 2 mM TBP, 65 mM DTT). Protein samples (200 or 20 µg of protein for Coomassie Blue stained reference gels, or gels to be processed for immunoblotting, respectively) were adsorbed overnight on Immobiline DryStrips (7 cm, pH 3-10 NL, or pH 4-7). Electrofocusing was performed in an IPGphor Isoelectric Focusing Unit (Amersham Biosciences, Uppsala, Sweden). The focused strips were equilibrated as described (15) and loaded on linear 9-16.5 % acrylamide gradients (7x 4 cm, 1.5 mm thick), for SDS-PAGE separation in a Mini Protean III Cell (Bio-Rad, Hercules, CA). Gels were stained with colloidal Coomassie Blue (Novex, San Diego, CA) (4) and the protein maps so obtained were scanned with a Personal Densitometer SI (Molecular Dynamics) at 12 bits and 50 mm per pixel.

For Western Blot analyses, the proteins separated in the 2DE maps were transferred onto nitrocellulose membranes, overnight at 30 Volts, using a Protean III apparatus (BioRad, Hercules, CA). Membranes were stained with a 0.05% (w/v) CPTS (Copper(II) phthalocyanine-3,4',4'',4'''-tetrasulfonic acid tetrasodium salt) in 12 mM HCl, and marked peripherally with 8 India-ink dots to provide anchors for subsequent image superimposition and matching. After scanning and image acquisition, the membranes were destained with 0.5 M NaHCO₃, incubated with the mouse sera to be analyzed (either pre-immune or specific immune sera, diluted 1:1000), and then with a peroxidase-conjugated anti-mouse antibody (Amersham Biosciences, Uppsala, Sweden). After washing with PBS, 0.1% Tween-20, blots were developed using the Opti-4CN Substrate Kit (Biorad, Hercules, CA), and the images of the immunostained blots again acquired as above. Images were analysed with the computer program Image Master 2D Elite, version 4.01 (Amersham Biosciences, Uppsala, Sweden). Superimposition and matches between Western-blot membranes and Coomassie stained gels were performed as follow. First, the CPTS-stained membrane image and the immunostained blot image were superimposed using the peripheral dot marks. Then, the sum image so obtained was superimposed to the Coomassie stained protein map using the CPTS stained CPn proteins as anchors. The areas on the Coomassie stained map corresponding to immunostained spots on the blot were excised from the preparative gel for protein identification. Protein sample were dried in a vacuum centrifuge, and in-gel digested, for 2h at 37°C, with an excess of porcine Trypsin (Promega, Madison, WI), in 100 mM ammonium bicarbonate. Tryptic peptides were desalted and concentrated using Zip-Tip (Millipore, Bedford, MA). Peptides were

5 directly eluted and loaded onto a SCOUT 384 Anchor Chip multiprobe plate (400 μ m, Bruker Daltonics, Bremen, Germany) with a solution of 2-5 dihydroxybenzoic acid (5g/l), in 50% acetonitrile, 0.1% trifluoroacetic acid. Spectra were acquired on a Bruker Biflex III matrix-assisted laser desorption ionization-time of flight (MALDI-TOF) apparatus. Resulting values for monoisotopic peaks were used for database searches using the Mascot software (32), as available at the website <http://www.matrixscience.com/>.

In vitro neutralization assays

10 *In vitro* neutralization assays were performed on LLC-MK2 (Rhesus monkey kidney) epithelial cell cultures. Serial four-fold dilutions of mouse immune and corresponding preimmune sera were prepared in sucrose-phosphate-glutamic acid buffer (SPG). Mouse polyclonal sera to whole EBs were used as positive control of neutralization, whereas SPG buffer alone was used as negative control of neutralization (control of
15 infection). Purified infectious EBs from the *C.pneumoniae* FB/96 were diluted in SPG buffer to contain 2.5×10^7 IFU/ml, and 10ul of EBs suspension were added to each serum dilution in a final volume of 100ul. Antibody-EB interaction was allowed to proceed for 30 min at 37°C on a slowly rocking platform. The 100ul of reaction mix of each sample was used to inoculate PBS-washed LLC-MK2 confluent
20 monolayers (in triplicate for each serum dilution), in a 24-well tissue culture plate, and centrifuged at 805 x g for 1 hour at 37°C. After centrifugation Eagle's minimal essential medium containing Earle's salts, 20% fetal bovine serum and 1ug/ml cycloheximide was added. Infected cultures were incubated at 37°C in 5%CO₂ for 72 hours. The monolayers were fixed with methanol and the *Chlamydial* inclusions were
25 detected by staining with mouse anti-*Chlamydia* fluorescein-conjugated monoclonal antibody (Merifluor *Chlamydia*, Meridian Diagnostics, Inc.) and quantified by counting 10 fields per well at a magnification of 40X. The inhibition of infectivity due to EBs interaction with the immune sera was calculated as percentage reduction in mean IFU number as compared to the SPG (buffer only)/EBs control. In this
30 calculation the IFU counts obtained with immune sera were corrected for background inhibition of infection due to the corresponding pre-immune mouse serum. According to common practice, the sera were considered as "neutralizing" if they could cause a 50% or greater reduction in infectivity. The corresponding neutralizing titer was defined as the serum dilution at which a 50% reduction of infectivity was observed.
35 Experimental variability was evaluated by calculating the standard error of measurement (SEM), from three titration experiments for each recombinant antigen, as shown in Fig.1B.

In vivo screening

40 *In vivo* evaluation was performed using a hamster model of systemic infection, as recently described (34). Essentially, adult (10-11 week old) Syrian hamsters (Morini, S. Polo D'Enza, Italy), previously immunized with the recombinant vaccine candidates were challenged systemically with infectious Cpn elementary bodies (EB). Protection was assessed by determining the levels of viable EB recovered from the
45 spleen, as compared to non-immunized animals. Statistical significance of the results was evaluated by a two-tailed Student's t-test.

Groups of 8 hamsters were immunized subcutaneously with recombinant antigens, or only injected with buffer for the control group, at days 0, 7, and 21. For each
50 immunization 20 ug protein 1:1 diluted with Freund's complete adjuvant (first dose)

and Freund's incomplete adjuvant (booster doses) was injected. At day 35 post-infection the hamsters were anaesthetised with Ketamine and inoculated intraperitoneally and intranasally with 0.1 ml of *C.pneumoniae* EB suspension (1.0×10^8) at each site. Animals were sacrificed seven days after infection. The spleen
5 was weighed, and homogenized in a mortar to obtain a 10% (wt/vol) suspension in cold SPG buffer. Tissue suspensions were centrifuged at 300 x g for 10 min at 4°C to remove coarse debris. The clarified homogenates (0.2 ml) were inoculated in duplicate onto LLC-MK2 cells seeded in plastic individual well of a 24 well plate, incubated at 37°C for 72 h and fixed in acetone before detection and counting of
10 numbers of *Chlamydial* inclusions per well by immunofluorescence microscopy. The protocol was approved by the ethical committee of the University of Bologna.

Example 1 (in vitro studies)

Screening antisera for in vitro neutralizing properties

Following a genome-wide screening for proteins likely to be localized on the cell surface of *C. pneumoniae*, we recently reported (25) that antisera to 53 recombinant *Chlamydial* antigens were capable to bind in a FACS assay, the surface of *Chlamydial* cells. In order to check whether some of the FACS positive antigens were capable of interfering with EB *in vitro* infectivity, we raised mouse antisera against the recombinant FACS positive antigens and assessed the effect of each antiserum on the infectivity of purified EBs with respect to monolayers of LLC-MK2 cells. Infectious EB were first incubated with the antiserum and then used to infect cell monolayers in 24-well multititer plates. In parallel, control samples were similarly processed in which the EBs were: i), either treated with buffer only, or, ii), treated with the same dilutions of the corresponding preimmune mouse sera.

Results I

Using this assay, 10 sera have so far proved to effectively neutralize *in vitro* infectivity to an extent greater than 50%, a property that common practice qualifies such antigens as "neutralising" (Figure 1). These 10 sera were obtained by mouse immunization with recombinant proteins derived from the following *C.pneumoniae* genes:

- *pmp10* and *pmp2*, encoding two members of the heterogeneous *Chlamydial* PMP family of polymorphic membrane proteins;
- *artJ*, encoding a putative extracellular solute (possibly Arginine) binding protein of an aminoacid transport system;
- *eno*, encoding a protein homologous to bacterial enolases, glycolytic enzymes which can be found also on the bacterial surface;
- *htrA*, encoding a putative chaperone with heat-shock inducible protease activity;
- the Cpn0301 "hypothetical" gene, encoding a protein homologous to the ompH family of bacterial proteins, some members of which have been shown to be chaperones involved in outer membrane biosynthesis;
- two Cpn-specific "hypothetical" genes Cpn0795 and Cpn0042;
- *omcA* encoding a 7-9 kDa protein annotated as an outer membrane protein; and
- *atoS* a putative sensor member of a transport system.

As shown in Figure 1 and summarized in Table I, OmpH, enolase and Cpn0795 appeared to induce the highest neutralizing sera, with titers above 400. By contrast, Pmp2, ArtJ and Cpn0042 induced titers equal or less than 100, while the remaining 4 antigens, Pmp10, HtrA, AtoS and OmcA showed intermediate titers.

Example 2 (in vivo studies)

Evaluation of antisera specificity by 2D immunoblot analysis of Cpn protein extracts

5 In order to investigate if the neutralizing activity observed in the *in vitro* infection of LLC-MK2 monolayers was actually due to the binding of the antibodies to the selected *C.pneumoniae* proteins, rather than to possible cross-reactions with other antigens, we assessed the specificity of the antisera by immunoblot analysis of two dimensional electrophoretic maps of EB proteins.

10 In particular, this analysis was carried out on six antigens (Pmp2, Pmp10, Eno, ArtJ, HtrA and OmpH-like) known to be visible in the 2D maps of EB total proteins (Montigiani *et al.*, 2002 Infection and Immunity 70: 368-379). Total EB proteins were resolved by 2D-electrophoresis using two different pH intervals (pH 3-10 non
15 linear, and pH 4-7, respectively) since previous experiments had shown that some of the proteins under study were better detected using one rather than the other of the above pH intervals. For each pH interval four gels were run in parallel. One gel was stained with Coomassie Blue to visualize the protein spots, while the other gels were blotted on nitrocellulose filters and stained with one of the selected sera at 1000-fold
20 dilution. Subsequently, the images of the immunostained blots (Fig.2, panels c to h) were superimposed to the corresponding Coomassie Blue-stained gel to identify the spots which had reacted with a given antiserum. The matching protein spots were excised and processed for peptide identification by MALDI-TOF analysis.

Results 2

25 In all six maps the immunoreactive protein species in the excised gel area were found to contain peptides from the expected *Chlamydial* protein. Even when the serum reacted with more than one electrophoretic protein species, the mass spectra of all spots which could be detected in the Coomassie Blue stains 2DE map were always consistent with the same polypeptide being present as multiple electrophoretic
30 species.

Interestingly, the immunoblot obtained with the HtrA antiserum showed two sets of 4 spots arranged as two typical electrophoretic 'trains' at two different molecular weights. On the Coomassie Blue stained gel it was possible to identify 4
35 corresponding spots, 3 in the upper train and 1 in the lower Mw set. MS analysis identified all of them as products of Cpn HtrA gene. Interestingly the lower Mwt species missed 3 N-terminal tryptic peptides, detected in the higher Mw spot series, and mapping within the first 100 aa of the ORF. These results suggest that HtrA was present in the EB protein sample both as a full *htrA* product, and as a discrete species missing a short N-terminal peptide, possibly as a result of some post-translational
40 processing.

Discussion of Results 2

In the analysis of data which are based on polyclonal antibody reactivity one should consider that cross-reactions due to epitope mimicry are always difficult to exclude.
45 The problem of antisera specificity was addressed in this work by 2D immunoblotting and identification of the reacting electrophoretic species by mass spectrometry analysis. This approach was possible for 6 of the 10 antisera, i.e. those corresponding to proteins previously identified by mass spectrometry (MALDI-TOF) analysis on 2D electrophoretic maps of *C.pneumoniae* EB proteins (25, 42) (Table 1, and Figure 2).
50 The probability of fortuitous cross-reactions between unrelated *Chlamydial* protein

species was minimized by the results of the immunoblotting analyses which showed that out of ca 300 protein spots in a map, all those reacting with the tested antisera were consistent with the expected antiserum specificity. Obviously, since during 2-D electrophoresis conformational epitopes are generally lost, structure-dependent cross-reactions cannot be ruled out in this type of analysis.

Example 3

In vivo evaluation of the in vitro neutralizing antigens in a hamster model of systemic infection

We have recently described a new hamster model of systemic *Chlamydia pneumoniae* infection in which replicating *Chlamydia* disseminate through macrophages and accumulate in the spleen (34). We therefore asked the question whether the *in vitro* neutralizing antigens we identified would also have protective activity *in vivo* using this model. To this aim, the 10 *in vitro* neutralizing recombinant antigens were used to immunize 8 hamsters with 3 subcutaneous injections over a three-week period, and challenged with 2×10^8 Cpn EBs two weeks later. Spleen infection was assessed 7 days after challenge. The difference between the mean number of infectious *Chlamydiae* recovered from control animals and the mean number of *Chlamydiae* recovered from animals immunized with the recombinant *Chlamydial* antigens, was taken as a measure of protection specifically induced by the putative vaccine candidate.

Results 3

The results of spleen protection observed for the various antigens in repeated experiments are shown in Figure 3 and reported as percentage values in Table 1. Six out of ten antigens, Pmp2, Pmp10, Enolase, the OmpH-like protein, and the products of the *C.pneumoniae*-specific genes Cpn0759 and Cpn0042, showed a statistically significant protective activity, with a reduction in IFU recovered from the spleens of immunized animals higher than 80% with respect to mock-immunized controls.

A limit of the hamster model is that, because of the absence of immunological reagents, the relative contribution of humoral and cell-mediated immunity cannot be assessed. However, we asked the question whether recombinant antigens could elicit also in the hamster neutralizing antibodies with sufficiently high titers. Therefore we tested the sera from hamsters immunized with Pmp2 and enolase, two of the most protective antigens, in the *in vitro* neutralization assay. Both antigens had a neutralizing titer of approximately 100 (data not shown).

Summary of Results 3

In conclusion, a considerable proportion (60%) of the *in vitro* neutralizing antigens were also protective in the hamster *in vivo* model and our data suggest that antibody-mediated neutralization could play a role at least in this model of systemic infection.

Discussion of Results 3

Beside assaying the *in vitro* neutralization properties of the selected subset of 10 FACS-positive antigens, we also assessed the performance of these antigens in protecting against *C. pneumoniae* infection in an animal model of systemic infection recently described in the hamster (34). This evaluation addressed the capability that the recombinant antigens would have of inducing a protective response against naturally replicating *Chlamydiae* (rather than EB's purified from *in vitro* cultures grown under artificial conditions) and in the context of a systemic infection. In fact

the hamster model we used, while it does not model the typical respiratory infection considered to be the predominant route by which *C. pneumoniae* infects humans, it nevertheless simulates a situation of systemic invasion which could be preliminary to the establishment of *C. pneumoniae* chronic infection considered by several authors as being associated to the development or progression of cardiovascular disease, and other chronic degenerative diseases. Notably, a limit of any hamster model is the current lack of hamster-specific immunological reagents which would allow the analysis of cell mediated immune responses. However, in the case of systemic infections, by common wisdom, neutralizing antibodies are likely to have a protective action. The finding that 6 of the 10 'in vitro neutralizing' antigens had also a >80% protective action *in vivo*, and that a measurable neutralizing activity was also found in the sera of immunized hamsters, suggests that a specific antibody mediated immunity could at least partially contribute to the animal protection here described.

Example 4

Two 'hypothetical' proteins 6784 and 6814 (encoded by the ORFs Cpn0498 and Cpn0525) yielded FACS-positive sera which, however, were not able to neutralize host cell infection *in vitro*. However, these antigens performed remarkably well in the hamster-spleen test.

Table 2

Gene/ORF ID in CWL029	Protein ID	Recombinant Fusion Type	Annotation	Reciprocal of 50% neutralisation titre	% Protection in the hamster spleen test (ref 34)
Cpn0498	4376784	GST	Hypothetical protein	0	94
CPn0525	4376814	GST	Hypothetical protein (similarity to CT398)	0	97
CPn0324		HIS	Low Calcium Response Element (LcrE)		Completely protected 8 of 16 animals and reduced the infectivity titres of the eight positive animals

Discussion of Results 4

Interestingly, whilst antiserum against CPn0525 gave negative *in vitro* results (ie no neutralising activity), the CPn0525 protein gave 97 per cent protection from spleen infection in an *in vivo* hamster immunisation assay (ie a positive *in vivo* result).
5 Likewise, whilst antiserum against Cpn0498 gave negative *in vitro* results (ie no neutralising activity), the CPn0498 protein gave 94 per cent protection from spleen infection in an *in vivo* hamster immunisation assay. Thus a positive signal obtained in the FACS assay does not guarantee a corresponding positive *in vitro* neutralization activity and conversely a negative neutralization activity does not mean that a positive
10 *in vivo* result can be excluded.

General Discussion of Results 1-4

Strategy for identification of *Chlamydia pneumoniae* antigens of interest

Our strategy was based on the following experimental steps: 1) analysis of
15 *Chlamydia* genome sequence to select putative membrane-associated antigens, 2) cloning, expression and purification selected antigens, 3) preparation of antigen-specific sera by mouse immunization with the purified antigens, 4) FACS analysis of *Chlamydia* EBs using the mouse sera to identified surface-exposed antigens, 5) "*in vitro* neutralization" assay to test whether antibodies elicited by a given antigen can
20 interfere with the process of eukaryotic cell infection, and 6) use of appropriate animal model to test the capacity of selected antigens to confer protection against *Chlamydia* challenge.

As recently described by Montigiani *et al* ((2002) Infection and Immunity 70: 368-
25 379) from the initial screening of the *C.pneumoniae* genome, a panel of mouse sera was prepared against over 170 recombinant His-tagged or GST-fusion proteins encoded by genes or "open reading frames" somehow predicted to be periferally located in the *Chlamydial* cell. When these antibodies were tested in a FACS assay for their ability to bind the surface of purified *C.pneumoniae* EBs, a list of 53 "FACS-
30 positive" sera was obtained. The corresponding putative surface antigens were then further assessed for their capability of inducing neutralizing antibodies. This part of the work involved testing which of the sera contained antibodies capable of interfering with the process of *in vitro* infection of epithelial cell cultures. In the *in vitro* "neutralization" assay purified infectious EBs are incubated with progressive
35 dilutions of the immune sera and, in parallel, dilutions of the corresponding pre-immune sera, and of sera against non *Chlamydia* control antigens.

Cell cultures are infected in the presence of cycloheximide, which inhibits host cell protein synthesis and favours *Chlamydial* intracellular growth with the consequent
40 formation of typical cytoplasmic inclusions which can be stained with *Chlamydia* specific fluorescence-labeled monoclonal antibodies and counted with an UV light microscope. Working with appropriate pathogen-to-host cell ratios, it can be reasonably assumed that the number of detected cytoplasmic inclusion is proportional to the number of infectious *Chlamydiae* in the original sample. So a reduction in
45 inclusion numbers caused by the presence of an antigen-specific antiserum, as compared to the numbers obtained with control sera, gives a measure of the capability of a given antigen to elicit antibodies which can inhibit some stage of the *Chlamydial* infection process. According to common convention, an anti-serum is labelled as 'neutralizing' when the reduction of infectivity is equal or greater than 50%, and the

serum dilution yielding a 50% reduction in infectivity is referred to as the 50% end-point neutralization titer.

5 Some of the results obtained by screening the panel of recombinant antigens with the
10 *C.pneumoniae* *in vitro* neutralization assay confirm that some of the listed antigens,
like the members of the family of heterogeneous polymorphic membrane proteins
(PMP), which, on the basis of published literature data, could be reasonably expected
to be surface-exposed and possibly induce neutralizing antibodies. However, there
15 are also proteins which could be considered so far only hypothetical, and proteins
which just on the basis of their current functional annotation could not be at all
expected to be found on the bacterial surface. Using an *in vitro* neutralising assay, it
was found that sera to 10 CPn antigens have so far proved to effectively neutralize *in*
vitro infectivity to an extent greater than 50%, a property that common practice
qualifies such antigens as "neutralising" (Figure 1). These 10 sera were obtained by
mouse immunization with recombinant proteins derived from the *C.pneumoniae*
genes listed below.

20 Using a recently described *in vivo* model of systemic infection (hamster model),
hamsters immunised with 6 of the *in vitro* neutralising antigens, when challenged with
CPn EBs, showed a greater than 80% reduction of spleen infection as compared with
non-immunised controls.

Characterisation of 10 CPn proteins

The proteins identified by the present work can be divided in 3 groups:

- 25 • proteins which have an annotation compatible with (could be reasonably expected
to have) an expected/predicted exposure on the *Chlamydial* cell surface and with the
possibility that antibodies binding to them may actually interfere with host cell
attachment and entry (ie proteins which could possibly induce neutralising antibodies)
- 30 • proteins which by homology with other gram-negative bacteria could be expected
to have a periplasmic exposure (ie would not be expected at all to be found on the
bacterial cell surface); and
- proteins which are still labelled as 'hypothetical' (ie cellular location and/or
cellular function not known)

35 Group 1

(Pmp proteins (*pmp2* and *pmp10*), *OmcA* and *OmpH*)

***Pmp* proteins (*pmp2* and *pmp10*)**

The first group includes the 2 polymorphic outer membrane proteins (Pmp's) Pmp2
and Pmp10 (10, 11, 14, 30), the outer membrane protein OmpH-like, and OmcA,
40 which is annotated (*Chlamydia* Genome Project at [http://Chlamydia-
www.berkeley.edu:4231/](http://Chlamydia-
www.berkeley.edu:4231/)) as "predicted 9-kD cysteine-rich, outer membrane protein,
lipoprotein". The Pmp family of *Chlamydia*-specific proteins is generally thought to
comprise probable pathogenicity factors, with an autonomous secretion capacity
(autotransporters), important for adhesion to host cells and are generally considered as
45 promising vaccine candidates. However, apart from very recent unpublished results
on Pmp21, this is the first time that antisera to recombinant Pmp's are reported to
have neutralizing properties.

OmcA

OmcA is the product of a gene co-transcribed in the same operon with the 60 kDa OmcB cystein-rich protein which is a major structural component of the *Chlamydial* outer membrane and a major immunogen in human *C. trachomatis* infections. OmcB and OmcA are likely to interact in some as yet unknown outer membrane structure, so it is possible that antibodies to OmcA can interfere with EB infectivity.

OmpH

Finally, the *Chlamydial* OmpH is probably a member of the OmpH (Skp) family of proteins which have been reported to have chaperonin activities in other bacteria very important for the correct biosynthesis of the outer membrane. These proteins appear to cooperate in this task with HtrA (see below). In fact, in *E.coli* single KO mutants of either OmpH (Skp) or HtrA (DegP) are still viable, but double mutants do not grow (37). It should be pointed out that even if the amino acid sequences of the ompH-like proteins of *Chlamydia* (all *C.pneumoniae* and *C.trachomatis* or *C.caviae* variants) line-up very well with the rest of the bacterial OmpH proteins, they are the only ones to be acidic, whereas the rest of the family comprises mostly very basic proteins (including some with histone like behaviour, at least *in vitro*). One could speculate that if the chaperone activity is maintained also in the ompH like *Chlamydial* proteins, this may be related to some *Chlamydial* peculiarity.

Second Group of Selected Proteins (ArtJ, AtoS, HtrA and Enolase)

The second group, which represents a somehow surprising finding, includes ArtJ, AtoS, HtrA and Enolase. If the current annotation (justified by analogy with homologous genes in other bacteria) is correct, all these proteins would be expected to have a periplasmic location in gram-negative bacteria, and to be surface-exposed only in a gram-positive bacterium. It is possible that owing to their atypical life cycle, requiring an efficient passage from a dormant spore-like status (the EB) to an active form needing to adapt quickly to host-cell responses to invasion, *Chlamydiae* in fact display some sensors directly on the outer surface of their infectious form.

ArtJ

In the case ArtJ – for which we have data supporting both antigen expression and serum specificity – the hypothesis of an atypical situation peculiar to *Chlamydia* is supported by the anomalous gene set-up resulting from the present analysis of the *Chlamydia* genomes. ArtJ is so annotated by analogy with the ART transport systems of *E.coli* which has 5 genes organized in two operons (24) : artPIQM and artJ which are responsible for the arginine transport. In Cpn however the artPIQM genes are absent and therefore it appears that *Chlamydial* ArtJ operates in a molecular context which is different from its *E.coli* model and must be peculiar to this species.

HtrA

HtrA (DegP), which in other bacteria has a complex hexameric structure, has been described as having multiple functions (3, 5, 18, 19, 27, 38) : a chaperonin assisting a correct outer membrane biogenesis, inducible protease for the elimination of misfolded membrane proteins, and also a sensor of 'stress' conditions. In *Chlamydia* none of these properties has been demonstrated yet, however we find that in purified EB HtrA is present in two forms one of which appears to be processed by being deprived of the N-terminal fragment. This fragment, if aligned with the homologous

HtrA sequence from *Thermotoga maritima* (18), would comprise a predicted loop acting as a structural lid controlling the access to the protease active. So it appears tempting to speculate that HtrA could have a similar protease activity and the two forms identified on the 2-D map represent the active and inactive species. Interestingly, the *C. trachomatis* HtrA ortholog is recognized by human sera from patients who had a *Chlamydial* genital infection (35), and a similarly HtrA is one of the antigens in the immunoproteome of *Helicobacter pylori* (13). Furthermore the homologue protein in *Haemophilus influenzae* is a protective antigen in both a passive infant rat model of bacteremia and the active chinchilla model of otitis media (23).

Enolase

Also in the second group of proteins expected to be located elsewhere than the cell surface, is Cpn enolase. This protein aligns with the well known family of conserved glycosylases, which are essentially cytoplasmic enzymes, but in *Streptococci* enolase has been shown to have also a cell surface location, and extracellular matrix binding properties (1, 28, 29)). Interestingly, Gaston and colleagues (8) also showed that in patients with reactive arthritis induced by *C. trachomatis*, enolase induces specific CD4⁺ T-cell responses. Furthermore, a clone responding to the enolase *C. trachomatis* ortholog, responded also to *C. pneumoniae* EBs, and, since no proliferative response could be observed using a fungal or a mammalian enolase, the authors of this study concluded that the CD4 T-cell stimulating epitope must be *Chlamydia* specific.

Third Group of Proteins

(unknown cellular location and/or cellular function, Cpn0759, Cpn0042)

The third of the 3 groups in which we propose to divide, just for the sake of discussion, the 10 neutralizing antigens above described, comprises two proteins which are still annotated in public *Chlamydial* databases as the hypothetical products of two CPn-specific genes: Cpn0759 and Cpn0042. The Cpn0759 gene is the second gene in a cluster of 6 Cpn-specific hypothetical genes (from Cpn0794 to Cpn0799) immediately upstream of the enolase gene. With the exception of Cpn0759 the products of all the other genes in the cluster share similarities of 30 to 40% over long stretches of amino acids. The Cpn0042 gene encodes a hypothetical protein, with 4 coiled-coil regions, which has been described as a member of a new family of hypervariable outer membrane proteins (33). Interestingly, the hypervariability of these proteins could be due to a strand-slippage mechanism induced by the presence of a poly(C) stretch within the coding region of the corresponding genes, a mechanism already described in the Pmp's family for the *pmp10* gene (30). However, as indicated by their annotation, the function of these proteins is still unknown, and our observations provide the first experimental indication of a possible function related to the *Chlamydial* infection process.

Fourth Group of Proteins

Cpn0498

So in this case the triple parallel-screening evaluation, with two positive and one negative result, brought to the identification of a previously unknown antigen (ie an antigen with unknown biological function) having, according to current views, just the desirable basic properties in terms of antigenic function of a vaccine candidate.

Example 5

Background

The main stages in the *Chlamydial* life cycle are:

- 5 (i) the binding to the host cell surface and entry into the cytoplasm through a specialised vacuole (the *Chlamydial* inclusion) by an extracellular spore-like infective form, called the elementary body (EB); and
- (ii) the conversion of the EB to a non-infective replicative form called a reticulate body (RB) that replicates by binary fission a number of times within the inclusion to form a microcolony.

10 The sets of genes which are expressed in the various stages of the *Chlamydial* life cycle and the signals that trigger the passage from one stage to the next are largely unknown and still need investigation.

15 Current DNA microarray techniques permit profiling of gene expression at the mRNA level as a function of the cellular state. This can lead to the identification of genes or clusters of genes whose up- or down-regulation is associated to a particular state of the cell and to the identification of therapeutically relevant targets. Using this technology, DNA fragments representing specific portions of all genes belonging to a given organism (the fragments can be derived from cDNA libraries or can be obtained
20 by PCR amplification and chemical synthesis) are chemically bound to the surfaces of solid supports (chips) at high densities and in an ordered manner. Currently up to 10,000 DNA fragments or 250,000 oligonucleotides can be spotted onto a single squared centimetre of chip surface. The DNA chips are then utilised to define which of the spotted genes are transcriptionally active in a particular cellular population. To do so,
25 RNA is prepared, labelled with fluorescent dyes and finally hybridised to the DNA fragments fixed to the surface of the chip. By using an appropriate computer-assisted fluorescence detector, the fluorescence signals emitted by each spot upon excitation with a laser beam is carefully quantified to define the transcription activity of all the arrayed genes.

30 CPn DNA microarrays have been developed to look at the transcriptional events which occur when a given CPn pathogen gets into contact with the host cells, both in *in vivo* and *in vitro* settings. DNA chips carrying the entire genome of a particular bacterium, such as the CPn bacterium can be prepared in a very short period of time
35 so that whole genome expression analysis can be determined.

Experimental Methodology

Specifically, a genomic DNA (open reading frame probes) microarray approach for gene expression in CPn bacteria was adopted. In this respect, an array was prepared
40 for the analysis of the CPn life cycle on the basis of the published annotation of the complete genome. The *Chlamydia* DNA chips carry about 1000 PCR-derived DNA fragments, which have an average size of 400-700bp and correspond to internal portions of all CPn annotated genes.

Results 5

45 Table 2 shows transcriptional activity for expressed genes for CPn EB selected by microarray. The data in Table 2 shows different mRNAs in order of abundance from cells in their infectious "spore-like" (EB) form. The cells were used at the end of their cycle where gene expression is likely to be at its highest. As values less than 300 is

likely to be background, the top set of proteins (approx top 30) with more intense signals are likely to be the most interesting proteins

A review of the values for the expressed genes indicates that three of the FACS positive CPn antigens (CPn0331 (hypothetical), CPn0234 (hypothetical) and CPn0572 (hypothetical) are all highly expressed genes.

Materials and Methods (Examples 6-8) (Reference Section II)

T cell Epitope prediction and peptide synthesis

T cell epitope prediction was carried out on the genomic sequence of *C. pneumoniae* CWL029 strain (Accession numbers NC 000922 or AE001363) using the BIMAS algorithm [24]. Synthetic peptides (purity > 80%) were synthesized by Primm Srl (Milan, Italy), suspended in 100% DMSO and kept at -20° C before use.

RMA-S/A2 cell line and HLA-A2 transgenic and non transgenic mice

The T cell lymphoma murine cell line RMA-S stably transfected with HLA-A2 (RMA-S/A2, H-2^b, TAP2⁺), was kindly provided by Dr. Barnaba, Università degli Studi "La Sapienza", Rome, Italy, and cultured at 37° C in RPMI-1640 (GIBCO) supplemented with heat inactivated 10% FCS, 100 IU/ml penicillin/streptomycin, 2 mM Lglutamine (GIBCO) and 5×10⁻⁵ M 2-ME (Sigma). H2-b HLA-A2 transgenic mice [35] were housed in a pathogen-free environment and screened for HLA-A2 expression by FCM carried out on total blood samples using the BB7.2 anti-A2 mAb [48]. Only mice with percentages of A2 expressing cells higher than 70-80 % were used for DNA immunization and *C. pneumoniae* infection experiments. Animals which showed no HLA-A2 expression were mated in order to obtain an HLA-A2 non transgenic population, to be used as a control in the experiments.

Epitope stabilization assay

RMA-S/A2 cells (3-5 x 10⁵/well) were seeded in serum-free RPMI medium, supplemented with human β2 microglobulin (3 μg/ml, Sigma), without or with the test peptide (10μM). Following overnight incubation at 26°C in humidified 5% CO₂ atmosphere, cells were shifted to 37° C for 2 h before determining the HLA-A2 expression level at the cell surface using the BB7.2 anti-A2 mAb and a PE-conjugated anti-mouse IgG (Jackson ImmunoResearch). Fluorescence intensity on living cells, which did not incorporate propidium iodide, was analyzed by FCM. As controls, corresponding samples without peptide and samples with peptide but treated only with the anti-mouse secondary antibody, were used.

Infection and DNA immunization of HLA-A2 transgenic and non transgenic mice

Transgenic mice were intranasally infected twice with a month interval, using 5x10⁵ *C. pneumoniae* FB/96 EBs [4] diluted in 50 μl of PBS. *C. pneumoniae* antigen coding genes were amplified by PCR using FB/96 genomic DNA, cloned into plasmid pcmvKaSF2120 [49] and verified by DNA sequence analysis. Fifty μg of endotoxin free recombinant plasmid DNA, diluted in Dulbecco's phosphate buffer (GIBCO), were injected into the tibialis muscle of mice at days 0, 21 and 35.

CD8⁺ T cells isolation and IFN-γ determination by ELISpot assay

Splenocytes from DNA immunized mice were prepared one week after the third immunization using Cell Strainer (Falcon) filters. Following red blood cells lysis,

CD8⁺ T cells from spleen cells suspensions were enriched by positive selection using magnetic activated cell sorting (MACS-Miltenyi Biotec) with CD8a (Ly-2) microbeads. CD8⁺ T cells purity was higher than 90%, as determined by FMC. Multiscreen 96-well nitrocellulose plates (Millipore) were coated with 5 µg/ml of the anti-mouse IFN-γ antibody (R4-6A2, PharMingen) in 100 µl of carbonate buffer, pH 9.2. After overnight incubation at 4°C, plates were saturated at 37°C with 200 µl of BSA (1%) in PBS for 2 h. Purified CD8⁺ (5x10⁴) were added in a total volume of 200 µl/well in the presence of irradiated (3,000 rad) spleen cells from non immunized HLA-A2 transgenic mice as a source of antigen-presenting cells (2x10⁵/well), 10 µg/ml of peptide and 10U/ml of human r-IL-2 (Chiron Corporation). After incubation for 20 h at 37° C, 5% CO₂, plates were washed and developed for bound IFN-γ using sequential incubations with biotinylated antimouse IFN-γ (XMB1.2, PharMingen), ExtrAvidin-alkaline phosphatase and substrate BCIP/NBT (Sigma) dissolved in water. Spots were enumerated in each well using a dissecting microscope. Medium containing an irrelevant peptide or without peptide was used as negative controls, while positive controls were represented by CD8⁺ T cells treated with ConA (5 µg/ml).

In vitro* cultures and flow cytometric analysis of splenocytes from transgenic and non transgenic mice infected with *C. pneumoniae

Splenocytes from infected mice were isolated one week after the second infection with *C. pneumoniae* Ebs. For *ex vivo* analysis of IFN-γ production, 2x10⁶ splenocytes were seeded in the presence of the test peptide (10µg/ml) and anti-mouse CD28 antibody (1µg/ml, PharMingen) as co-stimulus. After a two h incubation at 37° C, 5 % CO₂, Brefeldin A (10 µg/ml, Sigma) was added and the incubation was extended for 4 additional hours. Following two washes with PBS, cells were permeabilized, fixed and stained both with anti-murine-IFN-γ-(PE), anti-murine CD8 (APC) and anti-murine-CD69 (FITC) and the corresponding isotypes. Positive controls for cytokine production were performed on cells stimulated with anti-mouse CD3 and CD28 antibodies (1 µg/ml, PharMingen) . Cells cultured either in the absence of peptide or pulsed with the HepB negative control peptide were used as negative controls. All samples were analyzed using a FACS LSRII flow cytometer (Becton Dickinson). For analysis of IFN-γ production by short term T cell lines, 5-10x10⁶ splenocytes from infected mice were cultured for 6 days in the presence of the test peptide (20 µg/ml), with rIL-2 (10 µg/ml) being added after the first two days. At the end of the incubation period, cells were washed twice in RPMI, pulsed again for 6 h in the presence of the test peptide (10µg/ml), 1x10⁵ freshly prepared CD8 depleted antigen presenting cells from HLA-A2 transgenic mice (irradiated at 3000 rad) and anti-mouse CD28 antibody (1µg/ml, PharMingen) as co-stimulus. After a two h incubation at 37° C, 5 % CO₂, Brefeldin A (10 µg/ml, Sigma) was added, the incubation was extended for 4 additional hours and IFN-γ production was analyzed by FCM.

45

Example 6

In silico analysis of *Chlamydia pneumoniae* genome and prediction of HLA-A2 T cell epitopes

5 The genome of the *Chlamydia pneumoniae* CWL029 strain was used to predict 9mer peptide sequences with high probability to bind class I HLA-A2 molecules. The analysis was carried out using the predictive algorithm available at the NIH Bioinformatics & Molecular Analysis Section Web server (<http://bimas.cit.nih.gov/>), which ranks potential MHC binders according to the predictive half-time dissociation of peptide/MHC complexes [24]. Since some *Chlamydial* proteins have been reported to induce autoimmune responses [25-28], we restricted our search to a subset of proteins, distinctive of the *Chlamydia* genus, consisting of 13 protein identified as members of the type III secretion system, 17 Polymorphic Membrane Proteins (PMP) and 19 additional proteins, 5 of which already identified as EB surface antigens [4]. The predicted binding score of 157.22, obtained for the well characterized HIV-1 p17 gag epitope ⁷⁷SLYNTVATL⁸⁵ [29], was taken as an arbitrary cut-off for peptide selection. A total of 55 potential *C. pneumoniae*-derived T cell epitopes, belonging to 31 different proteins, were identified (Table I), which had predicted binding scores ranging from 156.77 to 42,485.263

20 **In vitro binding of peptides to HLA-A2**

The capacity of the selected peptides to bind to HLA-A2 was assessed using an *in vitro* MHC class I stabilization assay, carried out with the murine transporter associated with antigen processing (TAP)-deficient cell line RMA-S/A2, stably transfected with the human class I A2 gene. MHC class I molecules, cultured at 37° C, are unstably expressed on the cell surface of TAP-deficient cells [30-32]. Culturing the cells at 37° C in the presence of binding peptides, results in formation of a more stable MHC/peptide complex which can be monitored by flow cytometric analysis. RMA-S/A2 cells were therefore cultured overnight at 26° C in the presence of the test peptides, shifted to 37° C for 2 hours and the surface level of stabilized A2 molecules was quantified by direct staining with an anti-HLA-A2 specific mAb. Two known HLA-A2 restricted CTL epitopes were used as positive controls for binding to A2, the HIV-1 p17 gag peptide [29] and the influenza matrix M1 protein peptide FluMP58 [33], while the Hepatitis B virus envelope antigen peptide HbenvAg125 (HepB) was used as a negative control [34].

35 **Results 6**

The binding results obtained are shown in Table 4 and allowed the identification of 15 peptides with a net mean fluorescence intensity (Net MFI) higher than 92.3, corresponding to the value obtained with the HIV-1 p17 gag positive control peptide, 8 peptides with a Net MFI intermediate between the values 92.3 and 63.1, obtained with the two positive control peptides, and 12 peptides with an Net MFI ranging between 29.6 and 63. Fifteen of the *in silico* predicted peptides (27.2 %) did not confer stabilization to the A2 molecules, showing a Net MFI lower than 14, obtained with the HepB negative control peptide.

Example 7

Some HLA-A2 binders are recognized by CD8⁺ T cells from DNA-immunized transgenic mice

5 The *in vitro* assay with RMA-S/A2 cells allowed the definition of a set of peptides which were able to bind and stabilize the HLA-A2 molecules on the cell surface. To gain information about the possibility that the predicted epitopes could indeed be generated by *in vivo* processing of the antigens from which they were derived, peptide recognition by CD8⁺ T cells was studied under conditions in which the related complete antigen was intracellularly expressed and presented *in vivo*. The
10 full-length ORF sequences coding for 13 *Chlamydial* proteins, including a total of 24 predicted peptides, were cloned into a suitable DNA expression vector and each recombinant plasmids was used to immunize distinct groups of transgenic mice expressing a chimeric class I molecule composed of the $\alpha 1$ and $\alpha 2$ domains of HLA-0201 and the $\alpha 3$ domains, transmembrane and cytoplasmic, of H-2K^b [35].

15 The ORF sequences were selected among those containing either one or more epitopes positive in the *in vitro* assay or a combination of positive and negative epitopes. The ORF sequence corresponding to the outer membrane protein A (OMPA, CPn 0695) was included in this analysis, since human MHC-I-restricted epitopes have
20 already been reported for this protein in *C. trachomatis* [18;36]. One coding sequence, related to gene CPn 0131 was chosen, which included four epitopes, all negative in the *in vitro* stabilization assay. After three immunization cycles, transgenic mice were sacrificed, spleen CD8⁺ T cells were isolated, stimulated for 20 hour with the corresponding peptide and *ex vivo* IFN- γ production was assessed using an enzyme-linked immunospot (ELISpot) assay.
25

Results 7

DNA-mediated expression of the ORFs including peptides CH-6 (CPn 0811), CH-7 (CPn 0623), CH-10 (CPn 0828), CH-13 (CPn 0695, OMPA) and CH-37 (CPn 0210)
30 were associated with numbers of spot forming cells (SFC) significantly higher than those obtained with the HepB unrelated peptide, whereas some peptides related to antigens coded by genes CPn 0131, CPn 0323 and CPn 0062 induced SFC values only 2-3 times higher than the HepB control peptide (Table 5). Peptides related to antigens coded by genes CPn 0132, CPn 0322, CPn 0325, CPn 0415 and CPn 0728
35 did not induce any IFN- γ production (data not shown).

Example 8

To test the capacity of peptides to amplify specific CD8⁺ T cell populations *in vitro*, some of these plasmids were used to repeat the DNA immunization experiment and to
40 determine by flow cytometry the intracellular IFN- γ production by CD8⁺ T cells, both *ex vivo* and after a 6 day stimulation in the presence of the relevant peptides. In the attempt to establish a direct correlation between IFN- γ production by CD8⁺ T cells and HLA-A2 specific restriction, the experiment was carried out with both transgenic and non transgenic syngenic mice. The plasmids used contained genes CPn 0695, CPn
45 0811 and CPn 0823, including peptides CH-13, CH-6 and CH-7 respectively, which were highly positive in the *in vitro* binding and in the ELISpot assays and gene CPn 0323, including six different peptides, all of them with ELISpot values slightly higher than background

Results 8

The results of the experiment are summarized in Table 6, while representative dot plots from flow cytometric analysis of splenocytes stimulated with peptide CH-6 are shown in Fig. 4. When fresh spleen cells of DNA-immunized transgenic mice were pulsed with the tested peptides, only CH-6 or CH-7 induced relative fold increase (RFI) values about 5 times higher than those obtained pulsing the same cells with the HepB negative control peptide (Table 6, 4.58 and 5.2 RFI respectively).

When short term T cell lines (TCLs) instead of fresh splenocytes were used, a larger panel of peptides were able to trigger a significantly higher IFN- γ production by CD8⁺ T cells (Table 6). In fact, in addition to peptides CH-6 and CH-7, also peptides CH-13, CH-44, CH-45 and CH-46 were recognized by CD8⁺ T cell populations significantly larger than those induced by pulsing the same cells with the HepB peptide (RFI > 5). Importantly, peptide-induced IFN- γ production, appeared to be largely HLA-A2-dependent, since when the same experiments were carried out with non transgenic mice, the RFI values obtained were reliably lower (Table 6). The fact that non-transgenic and transgenic spleen cells were both efficiently activated using the polyclonal stimulus (anti-CD3/anti-CD28), reinforced the hypothesis that the lower CD8⁺ T cells induction in non transgenic mice was due to the absence of specific interactions between the peptides and the human HLA-A2 molecules.

CD8⁺ T cells of transgenic mice infected with *C. pneumoniae* recognize HLA-A2 binders *in vivo*

It has been recently shown that infection of mice with *C. pneumoniae* elicits a pathogen-specific murine class I-restricted immune response [22]. Therefore, we asked whether any of the A2 *in vitro* binders could be recognized by specific CD8⁺ T cells that are clonally selected during the immune response raised against the corresponding native antigen in *C. pneumoniae* infected cells.

To address this issue, HLA-A2 transgenic mice were intranasally infected with a non lethal dose of *C. pneumoniae* EBs and challenged with an equal dose of bacteria one month later, before being sacrificed to obtain splenocytes that were used to measure IFN- γ production by CD8⁺ T cells. Since no appreciable IFN- γ -production could be observed if splenocytes from infected mice were tested directly *ex vivo* (data not shown), spleen cells were cultured with each individual peptide or with the HepB irrelevant peptide for 6 days. The resulting short-term TCLs were then pulsed again for 6 hours with the same peptides and intracellular IFN- γ production by CD8⁺ T cells was assessed. The results obtained with 40 tested peptides are shown in Fig. 5A. Sixteen peptides (CH-2, CH-7, CH-8, CH-10, CH-13, CH-15, CH-20, CH-21, CH-28, CH-35, CH-37, CH-45, CH-46, CH-47, CH-50 and CH-55) elicited the strongest CD8⁺ responses (1 to 7.1 % of IFN- γ -producing CD8⁺ T cells), while 19 peptides elicited low but consistent responses (percentages of CD8⁺/IFN- γ ⁺ T cells between 0.3 and 0.9). Five peptides did not induce percentages of IFN- γ -producing CD8⁺ T cells significantly higher than those observed in response to the HepB control peptide.

When eight among the most reactive peptides were used again to pulse splenocytes of both transgenic and non transgenic mice infected with *C. pneumoniae*, seven of them were recognized by specific CD8⁺/IFN- γ ⁺ T cell populations present only in the transgenic mice, while peptide CH-7 was recognized by CD8⁺ T cells present in both mice groups (Fig. 5B).

General Discussion of Results in Examples 6-8

In this work we have described peptides derived from *C. pneumoniae* antigens identified as putative T cell epitopes recognized by the common human class I MHC A2 haplotype.

Understanding *C. pneumoniae*-specific CD8⁺ T cell-mediated immune response and designing protective vaccines rely on the possibility of identifying bacterial antigens or epitopes recognized by CD8⁺ T cells. Whereas the induction of a CTL-dependent immune response is predictable in response to pathogens which replicate in the cellular cytosol, providing antigens which can enter the cellular MHC-I presentation pathway, in the case of *Chlamydiae* it is not immediately obvious which antigens are made available to the proteasome and how they reach the cytosol, since these bacteria have a stringent intravacuolar localization inside the infected cell.

We have chosen an *in vivo* system based on HLA-A2 transgenic mice to test which of the predicted peptides could be recognized by specific CD8⁺ T cells following either DNA immunization with individual antigen coding genes or infection with *C. pneumoniae*. Our choice of a murine model is supported by previously published data. Wizel *et al.* [22], recently reported the first evidence that CD8⁺ T cells specific for *C. pneumoniae* antigens are induced in infected mice, and identified bacterial-derived murine MHC-I-restricted T cell epitopes able to trigger either lysis of *C. pneumoniae* infected cells or *in vitro* inhibition of the pathogen intracellular growth. These findings seem to confirm that some *C. pneumoniae* antigens can indeed reach the cytosol of infected cells and enter the MHC-I presentation pathway, i.e. during remodeling that occurs during *Chlamydia* replication or following autolysis of developing bacterial particles [22].

Furthermore, Kuon *et al.* [42] recently reported the identification of 11 *C. trachomatis*-derived HLA-B27-restricted peptides, capable of stimulating CD8⁺ T cells obtained from patients with *Chlamydia*-induced reactive arthritis. Importantly, 8 of them overlapped those selected by analyzing splenocytes of HLA-B27 transgenic mice infected with *C. trachomatis*, indicating that antigen processing can be closely reproduced using the murine animal model, although differences between murine and human antigen processing and T cell repertoires have been hypothesized [43].

The experiment which we have performed with *C. pneumoniae* infected A2 transgenic mice revealed that at least 16 peptides were recognized by IFN- γ -positive CD8⁺ T cell populations, which were actually expanded as a consequence of bacterial infection, since we could not detect IFN- γ production pulsing spleen cells from non infected transgenic mice with the same peptides (data not shown). These results suggest that the corresponding *Chlamydial* antigen may be able to enter the MHC-I presentation pathway. The finding that a number of these peptides can also be recognized by specific CD8⁺ T cells when the corresponding protein is separately expressed by DNA immunization, strongly reinforces the hypothesis that the responses observed with the infected mice are indeed specific for the *in silico* predicted peptide epitopes and their corresponding antigens. Importantly, the comparisons of peptide-induced IFN- γ -positive CD8⁺ T cells in A2 transgenic and non transgenic mice, either immunized with DNA or infected with *C. pneumoniae*, indicate that this recognition event is largely A2-specific.

Though, we cannot rule out the possibility that some of the tested peptides are also able to interact with the murine class-I MHC molecules, as suggested by the result obtained with CH-7 in infected non transgenic mice (Fig. 5) and by the RFI values obtained with CH-7, CH-8 and CH-13 in DNA-immunized non transgenic mice (Table 6).

Both with DNA immunization and bacterial infection, we were able to show that the OMPA-derived CH-13 peptide induces a specific CD8⁺ T cell response in A2 transgenic mice. These results appear to validate the choice of this animal model, since our observation that OMPA can enter the MHC-I presentation pathway correlates with the previous identification of HLA-A2-restricted and of murine MHC-I-restricted epitopes in OMPA proteins of *C. trachomatis* [18] and of *C. pneumoniae* [23] respectively. With the exception of CH-13 and CH-17, all the other peptides recognized by CD8⁺ T cells of infected mice belong to *C. pneumoniae* antigens for which neither human nor murine T cell epitopes have been identified [22;23]. Interestingly, a couple of positively reacting peptides (CH-50 and CH-55) belong to the group of polymorphic outer membrane proteins [44;45], while most of the others are part of the group of Type III secretion system-related proteins [45;46]. Peptides CH-7 and CH-8, both included in protein T of the *Yersinia* outer protein (Yop) system [47] and CH-10, included in protein J, which is part of the same translocation system, appear to be particularly reactive in the assay with the infected mice (Fig. 5A).

This is also true for other peptides included in antigens which are again related to the type III secretion system, such as CH-45, CH-46, and CH-47, all present in the low calcium response protein D. Intriguingly, CH-8, which is the most reactive peptide in the assay with the infected mice, does not seem to be recognized by a specific T cell population when the corresponding antigen is expressed by DNA immunization (Tables 5 and 6). This may be due to different factors, i.e. low *in vivo* expression level of the injected DNA or altered protein conformation.

On the other hand, we should also consider the possibility that, following infection of mice with *C. pneumoniae*, this peptide is recognized by a CD8⁺ T cell population which is instead specific for an epitope derived from an unidentified *C. pneumoniae* antigen having a closely related sequence. Contrarily to CH-8, stimulation of spleen cells from infected transgenic mice with peptide CH-6 did not allow the detection of IFN- γ ⁺/CD8⁺ T cells (Fig. 5A), but the same peptide was clearly reactive in the DNA immunization experiments (Tables 5 and 6). This may suggest that Low Calcium Response Protein H is not available for the cellular proteasome, but we could also assume either that the amount of peptide available to the MHC-presenting machinery is not sufficient to induce a cell response which is detectable with our assay, or that the reacting CD8⁺ T cell population does not expand over the detection limit of our assay.

On the whole, the results presented here allowed the identification of a number of antigens which may be available for proteasome-mediated processing in the course of *C. pneumoniae* infections, proposing them as possible targets for a HLA-A2-dependent cellular immune response. Further analysis will be required to verify if the specifically induced CD8⁺ T cells are able to recognize and induce lysis of peptide pulsed or *C. pneumoniae* infected mammalian cells and, possibly, to correlate the

- identified T cell epitopes with CD8⁺ T cell populations naturally induced in *C. pneumoniae* infected patients. Importantly, the results obtained with DNA-mediated expression of distinct antigens, can represent an initial step towards the definition of a significant set of *C. pneumoniae* HLA-A2-restricted epitopes, which could eventually be combined in DNA minigenes in the attempt to induce a CTL-dependent anti-*Chlamydia* protective immune response

Example 9

Immunizations with Combinations of the First Antigen Group

- The five antigens of the first antigen group (OmpH-like protein, pmp10, pmp2, Enolase, OmpH-like, CPn0042 and CPn00795) were prepared as described in the Materials and Methods Section above for Examples 1-4. The antigens are expressed and purified. Compositions of antigen combinations are then prepared comprising five antigens per composition (and containing 15 µg of each antigen per composition). CD1 mice are divided into seven groups (5-6 mice per group for groups 1 through 4; 3 to 4 mice for groups 5, 6 and 7), and immunized as follows:

Group	Immunizing Composition	Route of Delivery
1	Mixture of 5 antigens (15 µg/each) + CFA	Intra-peritoneal
2	Mixture of 5 antigens (15 µg/each) + AIOH (200µg)	Intra-peritoneal
3	Mixture of 5 antigens (15 µg/each) + AIOH (200µg) + CpG (10µg)	Intra-peritoneal
4	Complete Freund's Adjuvant (CFA)	Intra-peritoneal
5	Mixture of 5 antigens (5 µg/each) + LTK63 (5µg)	Intranasal
6	AIOH (200µg) + CpG (10µg)	Intra-peritoneal
7	LTK63 (5µg)	Intranasal

- Mice are immunized at two week intervals. Two weeks after the last immunization, all mice are challenged by intravaginal infection with *Chlamydia pneumoniae* serovar D.

- Experiment 9 was repeated with another group of CPn antigens. These were: CPn0385 (PepA), CPn0324 (LcrE), CPn0503 (DnaK), CPn0525 (Hypothetical) and CPn0482 (ArtJ). These antigens are combined and administered with and without alum and CpG as described in Experiment 9.

Summary

Applicants have identified a number of CPn proteins with desirable immunological and/or biological properties. Specifically, at least twelve CPn proteins have been identified which are capable of inducing the production of antibodies, which can neutralise, in a dose-dependent manner, the infectivity of *C. pneumoniae* in *in vitro* cell cultures. The induction of neutralising antibodies is important because it prevents infectious EBs from invading human tissues. Furthermore, at least six of these CPn proteins were also capable of attenuating *Chlamydial* (*C. pneumoniae*) infection in a *in vivo* hamster model. In addition, some of these CPn proteins were also capable of inducing not only adequate T-cell responses but also high serum levels of neutralising antibodies.

Apart from very recent unpublished results on pmp21, this is the first time that antisera to recombinant pmpps (pmp2 and pmp10) are reported to have neutralising properties. Interestingly, whilst antiserum against CPn0525 gave negative *in vitro* results (ie no neutralising activity), the CPn0525 protein gave 97 per cent protection from spleen infection in an *in vivo* hamster immunisation assay (see Table 2) (ie a positive *in vivo* result). Likewise, whilst antiserum against CPn0498 gave negative *in vitro* results (ie no neutralising activity), the CPn0498 protein gave 94 per cent protection from spleen infection in an *in vivo* hamster immunisation assay (ie a positive *in vivo* result). Thus a positive signal obtained in the FACS assay does not guarantee a corresponding positive *in vitro* neutralization activity and conversely a negative neutralization activity does not mean that a positive *in vivo* result can be excluded (also look for an example that a negative FACS assay result for a given antibody does not mean that a neutralizing activity can be excluded).

Some of the results obtained by screening the panel of recombinant antigens with the *C. pneumoniae in vitro* neutralization assay are shown in Table 2. Just by a cursory look at the 'current annotation' column it can be seen that both in Table 1 and 2 are listed antigens, like the members of the family of heterogeneous polymorphic membrane proteins (PMP), which, on the basis of published literature data, could be reasonably expected to be surface-exposed and possibly induce neutralizing antibodies, but there are also proteins which could be considered so far only hypothetical, and proteins which just on the basis of their current functional annotation could not be at all expected to be found on the bacterial surface.

The characterisation for the first time of some of these CPn proteins in terms of not only neutralising properties but also different score profiles in a panel of screening tests is an important contribution to the art because it facilitates the selective combination of CPn antigens with particular immunological and biological properties.

In conclusion, this paper describes a group of recombinant antigens which can induce antibodies inhibiting the infectivity of *C. pneumoniae in vitro* and have protective effects *in vivo*.

All publications mentioned in the above specification are herein incorporated by reference. Various modifications and variations of the described methods and system of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with specific preferred embodiments, it should be understood that the

invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the described modes for carrying out the invention which are obvious to those skilled in molecular biology or related fields are intended to be covered by the present invention.

5

REFERENCE SECTION I (Examples 1-4)

1. **Bergmann, S., M. Rohde, G. S. Chhatwal, and S. Hammerschmidt.** 2001. alpha-Enolase of *Streptococcus pneumoniae* is a plasmin(ogen)-binding protein displayed on the bacterial cell surface. *Mol Microbiol* 40:1273-87.
- 5 2. **Bush, R. M., and K. D. Everett.** 2001. Molecular evolution of the *Chlamydiaceae*. *Int J Syst Evol Microbiol* 51:203-20.
3. **Clausen, T., C. Southan, and M. Ehrmann.** 2002. The HtrA family of proteases: implications for protein composition and cell fate. *Mol Cell* 10:443-55.
- 10 4. **Doherty, N. S., B. H. Littman, K. Reilly, A. C. Swindell, J. M. Buss, and N. L. Anderson.** 1998. Analysis of changes in acute-phase plasma proteins in an acute inflammatory response and in rheumatoid arthritis using two-dimensional gel electrophoresis. *Electrophoresis* 19:355-63.
5. **Elzer, P. H., R. W. Phillips, G. T. Robertson, and R. M. Roop, 2nd.** 1996. The HtrA stress response protease contributes to resistance of *Brucella abortus* to killing by murine phagocytes. *Infect Immun* 64:4838-41.
- 15 6. **Everett, K. D., R. M. Bush, and A. A. Andersen.** 1999. Emended description of the order *Chlamydiales*, proposal of *ParaChlamydiaceae* fam. nov. and *Simkaniaceae* fam. nov., each containing one monotypic genus, revised taxonomy of the family *Chlamydiaceae*, including a new genus and five new species, and standards for the identification of organisms. *Int J Syst Bacteriol* 49 Pt 2:415-40.
- 20 7. **Farencena, A., M. Comanducci, M. Donati, G. Ratti, and R. Cevenini.** 1997. Characterization of a new isolate of *Chlamydia trachomatis* which lacks the common plasmid and has properties of biovar trachoma. *Infect Immun* 65:2965-9.
- 25 8. **Goodall, J. C., G. Yeo, M. Huang, R. Raggiaschi, and J. S. Gaston.** 2001. Identification of *Chlamydia trachomatis* antigens recognized by human CD4+ T lymphocytes by screening an expression library. *Eur J Immunol* 31:1513-22.
- 30 9. **Granoff, D. M., G. R. Moe, M. M. Giuliani, J. Adu-Bobie, L. Santini, B. Brunelli, F. Piccinetti, P. Zuno-Mitchell, S. S. Lee, P. Neri, L. Bracci, L. Lozzi, and R. Rappuoli.** 2001. A novel mimetic antigen eliciting protective antibody to *Neisseria meningitidis*. *J Immunol* 167:6487-96.
- 35 10. **Grimwood, J., L. Olinger, and R. S. Stephens.** 2001. Expression of *Chlamydia pneumoniae* polymorphic membrane protein family genes. *Infect Immun* 69:2383-9.
11. **Grimwood, J., and R. S. Stephens.** 1999. Computational analysis of the polymorphic membrane protein superfamily of *Chlamydia trachomatis* and *Chlamydia pneumoniae*. *Microb Comp Genomics* 4:187-201.
- 40 12. **Guan, K. L., and J. E. Dixon.** 1991. Eukaryotic proteins expressed in *Escherichia coli*: an improved thrombin cleavage and purification procedure of fusion proteins with glutathione S-transferase. *Anal Biochem* 192:262-7.
13. **Haas, G., G. Karaali, K. Ebermayer, W. G. Metzger, S. Lamer, U. Zimny-Arndt, S. Diescher, U. B. Goebel, K. Vogt, A. B. Roznowski, B. J. Wiedenmann, T. F. Meyer, T. Aebischer, and P. R. Jungblut.** 2002. Immunoproteomics of *Helicobacter pylori* infection and relation to gastric disease. *Proteomics* 2:313-24.
- 45 14. **Henderson, I. R., and A. C. Lam.** 2001. Polymorphic proteins of *Chlamydia* spp.--autotransporters beyond the Proteobacteria. *Trends Microbiol* 9:573-8.

15. Herbert, B. R., M. P. Molloy, A. A. Gooley, B. J. Walsh, W. G. Bryson, and K. L. Williams. 1998. Improved protein solubility in two-dimensional electrophoresis using tributyl phosphine as reducing agent. *Electrophoresis* 19:845-51.
- 5 16. Kalman, S., W. Mitchell, R. Marathe, C. Lammel, J. Fan, R. W. Hyman, L. Olinger, J. Grimwood, R. W. Davis, and R. S. Stephens. 1999. Comparative genomes of *Chlamydia pneumoniae* and *C. trachomatis*. *Nat Genet* 21:385-9.
- 10 17. Kawa, D. E., and R. S. Stephens. 2002. Antigenic topology of *Chlamydial* PorB protein and identification of targets for immune neutralization of infectivity. *J Immunol* 168:5184-91.
18. Kim, D. Y., D. R. Kim, S. C. Ha, N. K. Lokanath, C. J. Lee, H. Y. Hwang, and K. K. Kim. 2002. Crystal structure of the protease domain of a heat-shock protein HtrA from *Thermotoga maritima*. *J Biol Chem* 27:27.
- 15 19. Krojer, T., M. Garrido-Franco, R. Huber, M. Ehrmann, and T. Clausen. 2002. Crystal structure of DegP (HtrA) reveals a new protease-chaperone machine. *Nature* 416:455-9.
20. Kubo, A., and R. S. Stephens. 2000. Characterization and functional analysis of PorB, a *Chlamydia* porin and neutralizing target. *Mol Microbiol* 38:772-80.
- 20 21. Kuo, C. C., J. T. Grayston, L. A. Campbell, Y. A. Goo, R. W. Wissler, and E. P. Benditt. 1995. *Chlamydia pneumoniae* (TWAR) in coronary arteries of young adults (15-34 years old). *Proc Natl Acad Sci U S A* 92:6911-4.
22. Kuo, C. C., A. Shor, L. A. Campbell, H. Fukushi, D. L. Patton, and J. T. Grayston. 1993. Demonstration of *Chlamydia pneumoniae* in atherosclerotic lesions of coronary arteries. *J Infect Dis* 167:841-9.
- 25 23. Loosmore, S. M., Y. P. Yang, R. Oomen, J. M. Shortreed, D. C. Coleman, and M. H. Klein. 1998. The *Haemophilus influenzae* HtrA protein is a protective antigen. *Infect Immun* 66:899-906.
24. Makarova, K. S., A. A. Mironov, and M. S. Gelfand. 2001. Conservation of the binding site for the arginine repressor in all bacterial lineages. *Genome Biol* 2:RESEARCH0013.
- 30 25. Montigiani, S., F. Falugi, M. Scarselli, O. Finco, R. Petracca, G. Galli, M. Mariani, R. Manetti, M. Agnusdei, R. Cevenini, M. Donati, R. Nogarotto, N. Norais, I. Garaguso, S. Nuti, G. Saletti, D. Rosa, G. Ratti, and G. Grandi. 2002. Genomic approach for analysis of surface proteins in *Chlamydia pneumoniae*. *Infect Immun* 70:368-79.
- 35 26. Moroni, A., G. Pavan, M. Donati, and R. Cevenini. 1996. Differences in the envelope proteins of *Chlamydia pneumoniae*, *Chlamydia trachomatis*, and *Chlamydia psittaci* shown by two-dimensional gel electrophoresis. *Arch Microbiol* 165:164-8.
- 40 27. Pallen, M. J., and B. W. Wren. 1997. The HtrA family of serine proteases. *Mol Microbiol* 26:209-21.
28. Pancholi, V., and V. A. Fischetti. 1998. alpha-enolase, a novel strong plasmin(ogen) binding protein on the surface of pathogenic streptococci. *J Biol Chem* 273:14503-15.
- 45 29. Pancholi, V., and V. A. Fischetti. 1997. A novel plasminogen/plasmin binding protein on the surface of group A streptococci. *Adv Exp Med Biol* 418:597-9.

30. Pedersen, A. S., G. Christiansen, and S. Birkelund. 2001. Differential expression of Pmp10 in cell culture infected with *Chlamydia pneumoniae* CWL029. *FEMS Microbiol Lett* 203:153-9.
- 5 31. Perez Melgosa, M., C. C. Kuo, and L. A. Campbell. 1994. Isolation and characterization of a gene encoding a *Chlamydia pneumoniae* 76-kilodalton protein containing a species-specific epitope. *Infect Immun* 62:880-6.
32. Perkins, D. N., D. J. Pappin, D. M. Creasy, and J. S. Cottrell. 1999. Probability-based protein identification by searching sequence databases using mass spectrometry data. *Electrophoresis* 20:3551-67.
- 10 33. Rocha, E. P., O. Pradillon, H. Bui, C. Sayada, and E. Denamur. 2002. A new family of highly variable proteins in the *Chlamydia pneumoniae* genome. *Nucleic Acids Res* 30:4351-60.
34. Sambri, V., M. Donati, E. Storni, K. Di Leo, M. Agnusdei, R. Petracca, O. Finco, G. Grandi, G. Ratti, and R. Cevenini. 2004. Experimental infection by *Chlamydia pneumoniae* in the hamster. *Vaccine* in press.
- 15 35. Sanchez-Campillo, M., L. Bini, M. Comanducci, R. Raggiaschi, B. Marzocchi, V. Pallini, and G. Ratti. 1999. Identification of immunoreactive proteins of *Chlamydia trachomatis* by Western blot analysis of a two-dimensional electrophoresis map with patient sera. *Electrophoresis* 20:2269-79.
- 20 36. Schachter, J., R. S. Stephens, P. Timms, C. Kuo, P. M. Bavoil, S. Birkelund, J. Boman, H. Caldwell, L. A. Campbell, M. Chernesky, G. Christiansen, I. N. Clarke, C. Gaydos, J. T. Grayston, T. Hackstadt, R. Hsia, B. Kaltenboeck, M. Leinonen, D. Ocjius, G. McClarty, J. Orfila, R. Peeling, M. Puolakkainen, T. C. Quinn, R. G. Rank, J. Raulston, G. L. Ridgeway, P. Saikku, W. E. Stamm, D. T. Taylor-Robinson, S. P. Wang, and P. B. Wyrick. 2001. Radical changes to *Chlamydial* taxonomy are not necessary just yet. *Int J Syst Evol Microbiol* 51:249, 251-3.
- 25 37. Schafer, U., K. Beck, and M. Muller. 1999. Skp, a molecular chaperone of gram-negative bacteria, is required for the formation of soluble periplasmic intermediates of outer membrane proteins. *J Biol Chem* 274:24567-74.
- 30 38. Sebert, M. E., L. M. Palmer, M. Rosenberg, and J. N. Weiser. 2002. Microarray-based identification of htrA, a *Streptococcus pneumoniae* gene that is regulated by the CiaRH two-component system and contributes to nasopharyngeal colonization. *Infect Immun* 70:4059-67.
- 35 39. Siscovick, D., R. Alexander, M. Davidson, M. Leinonen, S. O'Connor, P. Ewald, C. Meier, M. Puolakkainen, J. Hughes, and J. Nieto. 2000. Collaborative multidisciplinary workshop report: the role of epidemiology studies in determining a possible relationship between *Chlamydia pneumoniae* infection and atherothrombotic diseases. *J Infect Dis* 181:S430-1.
- 40 40. Siscovick, D. S., S. M. Schwartz, M. Caps, S. P. Wang, and J. T. Grayston. 2000. *Chlamydia pneumoniae* and atherosclerotic risk in populations: the role of seroepidemiology. *J Infect Dis* 181:S417-20.
41. Stephens, R. S. (ed.). 1999. *Chlamydia: Intracellular Biology, Pathogenesis and Immunology*. American Society for Microbiology Press, Washington.
- 45 42. Vandahl, B. B., S. Birkelund, H. Demol, B. Hoorelbeke, G. Christiansen, J. Vandekerckhove, and K. Gevaert. 2001. Proteome analysis of the *Chlamydia pneumoniae* elementary body. *Electrophoresis* 22:1204-23.
43. Wolf, K., E. Fischer, D. Mead, G. Zhong, R. Peeling, B. Whitmire, and H. D. Caldwell. 2001. *Chlamydia pneumoniae* major outer membrane protein is a
- 50

surface-exposed antigen that elicits antibodies primarily directed against conformation-dependent determinants. Infect Immun 69:3082-91.

44. **Young, I. T.** 1977. Proof without prejudice: use of the Kolmogorov-Smirnov test for the analysis of histograms from flow systems and other sources. J Histochem Cytochem 25:935-41.

5

REFERENCE SECTION II (EXAMPLES 6-8)

1. Pizza,M., Scarlato,V., Massignani,V., Giuliani,M.M., Aricò, ,B., Comanducci,M., Jennings,G.T., Baldi,L., Bartolini,E., Capecchi,B.,
5 Galeotti,C.L., Luzzi,E., Manetti,R., Marchetti,E., Mora,M., Nuti,S., Ratti,G., Santini,L., Savino,S., Scarselli,M., Storni,E., Zuo,P., Broeker,M., Hundt,E., Knapp,B., Blair,E., Mason,T., Tettelin,H., Hood,D.W., Jeffries,A.C., Saunders,N.J., Granoff,D.M., Venter,J.C., Moxon,E.R., Grandi,G., and Rappuoli,R., Identification of Vaccine Candidates Against Serogroup B
10 Meningococcus by Whole-Genome Sequencing. *Science* 2000. 287: 1816-1820.
2. Rappuoli,R., Reverse vaccinology. *Current Opinion in Microbiology* 2000. 3: 445-450.
- 15 3. Rappuoli,R., Reverse vaccinology, a genome-based approach to vaccine development. *Vaccine* 2001. 19: 2688-2691.
4. Montigiani,S., Falugi,F., Scarselli,M., Finco,O., Petracca,R., Galli,G., Mariani,M., Manetti,R., Agnusdei,M., Cevenini,R., Donati,M., Nogarotto,R.,
20 Norais,N., Garaguso,I., Nuti,S., Saletti,G., Rosa,D., Ratti,G., and Grandi,G., Genomic Approach for Analysis of Surface Proteins in *Chlamydia pneumoniae*. *Infect.Immun.* 2002. 70: 368-379.
5. Ross,B.C., Czajkowski,L., Hocking,D., Margetts,M., Webb,E., Rothel,L.,
25 Patterson,M., Agius,C., Camuglia,S., and Reynolds,E., Identification of vaccine candidate antigens from a genomic analysis of *Porphyromonas gingivalis*. *Vaccine* 2001. 19: 4135-4142.
6. Wizemann,T.M., Heinrichs,J.H., Adamou,J.E., Erwin,A.L., Kunsch,C.,
30 Choi,G.H., Barash,S.C., Rosen,C.A., Masure,H.R., Tuomanen,E., Gayle,A., Brewah,Y.A., Walsh,W., Barren,P., Lathigra,R., Hanson,M., Langermann,S., Johnson,S., and Koenig,S., Use of a Whole Genome Approach To Identify Vaccine Molecules Affording Protection against *Streptococcus pneumoniae* Infection. *Infect.Immun.* 2001. 69: 1593-1598.
- 35 7. De Groot,A.S., Bosma,A., Chinai,N., Frost,J., Jesdale,B.M., Gonzalez,M.A., Martin,W., and Saint-Aubin,C., From genome to vaccine: in silico predictions, ex vivo verification. *Vaccine* 2001. 19: 4385-4395.
- 40 8. Martin,W., Sbail,H., and De Groot,A.S., Bioinformatics tools for identifying class I-restricted epitopes. *Methods* 2003. 29: 289-298.
9. Hill, A. V. and Davenport, M. P., Reverse Immunogenetics: from HLA-disease associations to vaccine candidates. *Mol.Med.Today* 1996. 2: 38-45.
- 45 10. Pamer, E. G., Harty, J. T., and Bevan, M. J., Precise prediction of a dominant class I MHC-restricted epitope of *Listeria monocytogenes*. *Nature* 1991. 353 (6347): 852-855.

11. Kuo,C.C., Jackson,L.A., Campbell,L.A., and Grayston,J.T., *Chlamydia pneumoniae* (TWAR). *Clin.Microbiol.Rev.* 1995. 8: 451-461.
- 5 12. Grayston, J. T., Background and current knowledge of *Chlamydia pneumoniae* and atherosclerosis. *The Journal of Infectious Diseases* 2000. 181: S402-S410.
13. Campbell,L. and Kuo,C.C., *Chlamydia pneumoniae* and atherosclerosis. *Seminars in Respiratory Infections* 2003. 18: 48-54.
- 10 14. Hammerschlag, M. R., The intracellular life of chlamidia. *Semin.Pediatr.Infect.Dis.* 2002. 13 (4): 239-248.
- 15 15. Rottenberg,M.E., Gigliotti Rothfuchs,A.C., Gigliotti,D., Svanholm,C., Bandholtz,L., and Wigzell,H., Role of Innate and Adaptive Immunity in the Outcome of Primary Infection with *Chlamydia pneumoniae*, as Analyzed in Genetically Modified Mice. *J Immunol* 1999. 162: 2829-2836.
- 20 16. Penttila, Anttila, Varkila, Puolakkainen, Sarvas, Makela, and Rautonen, Depletion of CD8⁺ cells abolishes memory in acquired immunity against *Chlamydia pneumoniae* in BALB/c mice. *Immunology* 1999. 97: 490-496.
- 25 17. Halme,S., Latvala,J., Karttunen,R., Palatsi,I., Saikku,P., and Surcel,H.M., Cell-Mediated Immune Response during Primary *Chlamydia pneumoniae* Infection. *Infect.Immun.* 2000. 68: 7156-7158.
- 30 18. Kim,S.K., Angevine,M., Demick,K., Ortiz,L., Rudersdorf,R., Watkins,D., and DeMars,R., Induction of HLA Class I-Restricted CD8⁺ CTLs Specific for the Major Outer Membrane Protein of *Chlamydia trachomatis* in Human Genital Tract Infections. *J Immunol* 1999. 162: 6855-6866.
- 35 19. Read,T.D., Brunham,R.C., Shen,C., Gill,S.R., Heidelberg,J.F., White,O., Hickey,E.K., Peterson,J., Utterback,T., Berry,K., Bass,S., Linher,K., Weidman,J., Khouri,H., Craven,B., Bowman,C., Dodson,R., Gwinn,M., Nelson,W., DeBoy,R., Kolonay,J., McClarty,G., Salzberg,S.L., Eisen,J., and Fraser,C.M., Genome sequences of *Chlamydia trachomatis* MoPn and *Chlamydia pneumoniae* AR39. *Nucl.Acids.Res.* 2000. 28: 1397-1406.
- 40 20. Shirai,M., Hirakawa,H., Kimoto,M., Tabuchi,M., Kishi,F., Ouchi,K., Shiba,T., Ishii,K., Hattori,M., Kuhara,S., and Nakazawa,T., Comparison of whole genome sequences of *Chlamydia pneumoniae* J138 from Japan and CWL029 from USA. *Nucl.Acids.Res.* 2000. 28: 2311-2314.
- 45 21. Kalman, S., Mitchell, W., Maranthé, R., Lammel, C., Fan, J., Hyman, R. W., Olinger, R., Grimwood, J., Davis, R. W., and Stephens, R. S., Comparative genomes of *Chlamydia pneumoniae* and *C. trachomatis*. *Nat.Genet.* 1999. 21 (4), 385-389.
22. Wizel,B., Starcher,B.C., Samten,B., Chroneos,Z., Barnes,P.F., Dzuris,J., Higashimoto,Y., Appella,E., and Sette,A., Multiple *Chlamydia pneumoniae*

Antigens Prime CD8+ Tc1 Responses That Inhibit Intracellular Growth of This Vacuolar Pathogen. *J Immunol* 2002. 169: 2524-2535.

23. Saren,A., Pascolo,S., Stevanovic,S., Dumrese,T., Puolakkainen,M.,
5 Sarvas,M., Rammensee,H.G., and Vuola,J.M., Identification of *Chlamydia pneumoniae*-Derived Mouse CD8 Epitopes. *Infect.Immun.* 2002. 70: 3336-3343.
24. Parker, K. C., Bednarek, M. A., and Coligan, J. E., Scheme for ranking
10 potential HLA-A2 binding peptides based on independent binding of individual peptide side-chains. *J.Immunol.* 1994. 152: 163-175.
25. Yi,Y., Yang,X., and Brunham,R.C., Autoimmunity to heat shock protein 60 and antigen-specific production of interleukin-10. *Infect.Immun.* 1997. 65: 1669-1674.
- 15 26. Lamb,D.J., El Sankary,W., and Ferns,G.A.A., Molecular mimicry in atherosclerosis: a role for heat shock proteins in immunisation. *Atherosclerosis* 2003. 167: 177-185.
- 20 27. Stephens, R. S., The cellular paradigm of *Chlamydial* pathogenesis. *Trends Microbiol.* 2003. 11 (1), 44-51.
28. Morrison, RP., *Chlamydial* hsp60 and the immunopathogenesis of *Chlamydial* disease. *Semin.Immunol.* 2003. 3 (1): 25-33.
- 25 29. Nixon, DF and McMichael, A. J., Cytotoxic T-cell recognition of HIV proteins and peptides. *AIDS* 1991. 5 (9): 1049-1059.
- 30 30. Anderson,K.S., Alexander,J., Wei,M., and Cresswell,P., Intracellular transport of class I MHC molecules in antigen processing mutant cell lines. *J.Immunol.* 1993. 151: 3407-3419.
31. Ljunggren,H.G., Stam,N.J., Ohlen,C., Neefjes,J.J., Hoglund,P., Heemels,M.T., Bastin,J., Schumacher,T.N., Townsend,A., Karre,K., and ., Empty MHC class I molecules come out in the cold. *Nature* 1990. 346: 476-480.
- 35 32. Ljunggren,H.G., Ohlen,C., Hoglund,P., Franksson,L., and Karre,K., The RMA-S lymphoma mutant; consequences of a peptide loading defect on immunological recognition and graft rejection. *Int.J.Cancer Suppl* 1991. 6: 38-44.
- 40 33. Bednarek,M.A., Sauma,S.Y., Gammon,M.C., Porter,G., Tamhankar,S., Williamson,A.R., and Zweerink,H.J., The minimum peptide epitope from the influenza virus matrix protein. Extra and intracellular loading of HLA-A2. *J Immunol* 1991. 147: 4047-4053.
- 45 34. Gagliardi,M.C., De Petrillo,G., Salemi,S., Boffa,L., Longobardi,M.G., Dellabona,P., Casorati,G., Tanigaki,N., Harris,R., and Lanzavecchia,A., Presentation of peptides by cultured monocytes or activated T cells allows specific priming of human cytotoxic T lymphocytes *in vitro*. *Int.Immunol.* 1995. 7: 1741-1752.

35. Vitiello, A., Marchesini, D., Furze, J., Sherman, L. A., and Chesnut, R. W., Analysis of HLA-restricted influenza-specific cytotoxic T lymphocyte response in transgenic mice carrying a chimeric human-mouse class I major histocompatibility complex. *J.Exp.Med.* 1991. 173 (4): 1007-1015.
- 5 36. Kim, S.K., Devine, L., Angevine, M., DeMars, R., and Kavathas, P.B., Direct Detection and Magnetic Isolation of *Chlamydia trachomatis* Major Outer Membrane Protein-Specific CD8+ CTLs with HLA Class I Tetramers. *J Immunol* 2000. 165: 7285-7292.
- 10 37. Meister, G.E., Roberts, C.G.P., Berzofsky, J.A., and De Groot, A.S., Two novel T cell epitope prediction algorithms based on MHC-binding motifs; comparison of predicted and published epitopes from *Mycobacterium tuberculosis* and HIV protein sequences. *Vaccine* 1995. 13: 581-591.
- 15 38. Schafer, J.R., Jesdale, B.M., George, J.A., Kouttab, N.M., and De Groot, A.S., Prediction of well-conserved HIV-1 ligands using a matrix-based algorithm, EpiMatrix. *Vaccine* 1998. 16: 1880-1884.
- 20 39. Reche, P.A., Glutting, J.P., and Reinherz, E.L., Prediction of MHC class I binding peptides using profile motifs. *Human Immunology* 2002. 63: 701-709.
- 25 40. Rammensee, H. G., Bachmann, J., Emmerich, N. P., Bachor, O. A., and Stevanovic, S., SYFPEITHI: database for MHC ligands and peptide motifs. *Immunogenetics* 1999. 50 (3-4): 213-219.
- 30 41. Cerundolo, V., Alexander, J., Anderson, K., Lamb, C., Cresswell, P., McMichael, A., Gotch, F., and Townsend, A., Presentation of viral antigen controlled by a gene in the major histocompatibility complex. *Nature* 1990. 345: 449-452.
- 35 42. Kuon, W., Holzhutter, H.G., Appel, H., Grolms, M., Kollnberger, S., Traeder, A., Henklein, P., Weiss, E., Thiel, A., Lauster, R., Bowness, P., Radbruch, A., Kloetzel, P.M., and Sieper, J., Identification of HLA-B27-Restricted Peptides from the *Chlamydia trachomatis* Proteome with Possible Relevance to HLA-B27-Associated Diseases. *J Immunol* 2001. 167: 4738-4746.
- 40 43. Endert van, P. M., Riganelli, D., Greco, G., Fleischhauer, K., Sidney, J., Sette, A., and Bach, J. F., The peptide-binding motif for the human transporter associated with antigen processing. *J.Exp.Med.* 1995. 182 (6): 1883-1895.
- 45 44. Grimwood, J. and Stephens, R. S., Computational analysis of the polymorphic membrane protein superfamily of *Chlamydia trachomatis* and *Chlamydia pneumoniae*. *Microb.Comp.Genomics* 1999. 4: 187-201.
- 46 45. Rockey, D. D., Lenart, J., and Stephens, R. S., Genome sequencing and our understanding of *Chlamydiae*. *Infection and Immunity* 2000. 68 (10): 5473-5479.
- 46 46. Kim, J.F., Revisiting the *Chlamydial* type III protein secretion system: clues to the origin of type III protein secretion. *Trends in Genetics* 2001. 17: 65-69.

47. **Cornelis, G. R.,** The *Yersinia* Ysc-Yop virulence apparatus. *Int.J.Med.Microbiol.* 2002. 291 (6-7): 455-462.
- 5 48. **Parham, P. and Brodsky, FM.,** Partial purification and some properties of BB7.2. A cytotoxic monoclonal antibody with specificity for HLA-A2 and a variant of HLA-A28. *Hum.Immunol.* 2003. 3 (4): 277-299.
- 10 49. **Grifantini, R., Finco, O., Bartolini, E., Draghi, M., Del Giudice, G., Kocken, C., Thomas, A., Abrignani, S., and Grandi, G.,** Multi-plasmid DNA vaccination avoids antigenic competition and enhances immunogenicity of a poorly immunogenic plasmid. *Eur.J.Immunol.* 1998. 28 (4): 1225-1232.

Figure 1A





Figure 1B



BEST AVAILABLE COPY




Figure 2

BEST AVAILABLE COPY





Figure 3. Mean numbers of *C.pneumoniae* IFU recovered from equivalent spleen samples from immunized and mock-immunized hamsters following a systemic challenge. Standard deviation values are shown above the bars. Antigens which induced significant protection are highlighted with an asterisk above the corresponding bar. All antigens were delivered in Freund's adjuvant. n.i. = non immunized controls


BEST AVAILABLE COPY

Figure 4



BEST AVAILABLE COPY

Figure 5



BEST AVAILABLE COPY

APPLICATION DATA SHEET**Application Information**

Application number:	To Be Assigned
Filing Date:	March 2, 2004
Application Type:	Provisional
Subject Matter:	Utility
Suggested classification:	To Be Assigned
Suggested Group Art Unit:	To Be Assigned
CD-ROM or CD-R?	None
Number of CD disks:	
Number of copies of CDs:	
Sequence submission?	None
Computer Readable Form (CRF)?	None
Number of copies of CRF:	
Title:	Immunogenic Compositions For Chlamydia Pneumoniae
Attorney Docket Number:	21431.001
Request for Early Publication?	No
Request for Non-Publication?	No
Suggested Drawing Figure:	
Total Drawing Sheets:	6
Small Entity?	No
Petition included?	No
Petition Type:	
Licensed U.S. Gov't Agency:	No
Contract or Grant No:	
Secrecy Order in Parent Appl.?	No

First Applicant Information

Applicant Authority Type:	Inventor
Primary Citizenship Country:	
Status:	Full Capacity
Given Name:	Guido
Middle Name:	
Family Name:	Grandi
Name Suffix:	
City of Residence:	Milano
State or Province of Residence:	
Country of Residence:	Italy
Street of mailing address:	c/o Chiron Corporation, P.O. Box 8097
City of mailing address:	Emeryville
State or Province of mailing address:	CA
Country of mailing address:	US
Postal or Zip Code of mailing address:	94662-8097

Second Applicant Information

Applicant Authority Type:	Inventor
Primary Citizenship Country:	
Status:	Full Capacity
Given Name:	Giulio
Middle Name:	
Family Name:	Ratti
Name Suffix:	
City of Residence:	
State or Province of Residence:	

Country of Residence:

Street of mailing address:

c/o Chiron Corporation, P.O. Box 8097

City of mailing address:

Emeryville

State or Province of mailing address:

CA

Country of mailing address:

US

Postal or Zip Code of mailing address:

94662-8097

Correspondence Information

Correspondence Customer Number:

27476

Name:

Street of mailing address:

City of mailing address:

State or Province of mailing address:

Country of mailing address:

Postal or Zip Code of mailing address:

Phone number:

Fax Number:

E-Mail address:

Representative Information

Representative Customer Number:		
---------------------------------	--	--

Domestic Priority Information

Application :	Continuity Type:	Parent Application:	Parent Filing Date:

Application :	Continuity Type:	Parent Application:	Parent Filing Date:

Foreign Priority Information

Country:	Application number:	Filing Date:	Priority Claimed:

Assignee Information

Assignee name:	Chiron S.r.l.
Street of mailing address:	4560 Horton Street
City of mailing address:	Emeryville
State or Province of mailing address:	CA
Country of mailing address:	US
Postal or Zip Code of mailing address:	94608-2916